

1. Consider the function

$$f(x, y) = x^3 - y^2 - 3xy + 1 \quad \text{for } (x, y) \in \mathbb{R}^2.$$

a) Which of the following statements are true?

4 Punkte

Mark your answers directly on this sheet by ticking the correct circle. Each statement gives one point and no justification is needed. Wrong or multiple answers give zero points.

true	false	
<input type="radio"/>	<input type="radio"/>	The gradient of f is $\nabla f(x, y) = \begin{pmatrix} 3x^2 - 3y \\ -3x - 2y \end{pmatrix}$.
<input type="radio"/>	<input type="radio"/>	At the point $(-1, 1)$ the function f has the biggest decrease in direction $\begin{pmatrix} 0 \\ 1 \end{pmatrix}$.
<input type="radio"/>	<input type="radio"/>	The function f has three critical points.
<input type="radio"/>	<input type="radio"/>	The function f has a local minimum at the point $(-\frac{3}{2}, \frac{9}{4})$.

b) Determine the slope of the level set of f at the point $(-1, 0)$.

Hint: Use the tangent to that level set.

4 Punkte

2. a) Consider the surface S given by

$$z = x^2 + y^2, \quad 0 \leq z \leq 1.$$

Use Stokes' Theorem in order to compute the downward flux of the rotation of the vectorfield

$$\vec{F}(x, y, z) := \begin{pmatrix} y + 1 \\ z - x \\ x - 2 \end{pmatrix}$$

through S . That is, compute

$$\iint_S \operatorname{rot}(\vec{F}) \cdot \vec{n} \, dA$$

where \vec{n} points into the negative z -direction.

7 Punkte

- b) Let \vec{G} be a divergence-free vector field on the whole of \mathbb{R}^3 .

Which of the following statements are true in general?

3 Punkte

Mark your answers directly on this sheet by ticking the correct circle. Each statement gives one point and no justification is needed. Wrong or multiple answers give zero points.

true	false	
<input type="radio"/>	<input type="radio"/>	For the three components of \vec{G} we have everywhere $\frac{\partial G_1}{\partial x} + \frac{\partial G_2}{\partial y} + \frac{\partial G_3}{\partial z} = 0.$
<input type="radio"/>	<input type="radio"/>	The line integral $\oint_C \vec{G} \cdot d\vec{r}$ along a closed curve C is always equal to 0.
<input type="radio"/>	<input type="radio"/>	The flux $\iint_A \vec{G} \cdot \vec{n} \, dA$ out of a closed surface A is always equal to 0.

3. Solve the following wave equation

$$\begin{cases} u_{tt} = u_{xx} , \\ u_x(0, t) = 0 , \\ u_x(1, t) = 0 , \\ u(x, 0) = 2 \cos^2(3\pi x) - \cos(7\pi x) , \\ u_t(x, 0) = \cos(4\pi x). \end{cases}$$

You may use here the following basis solutions without proof


$$1, \quad \cos(n\pi t) \cdot \cos(n\pi x), \quad \sin(n\pi t) \cdot \cos(n\pi x), \quad n = 1, 2, \dots$$

Hint: $\cos^2 \theta = \frac{1}{2} + \frac{\cos(2\theta)}{2}$.

7 Punkte

For exercises 7-31: Each question gives two points. Wrong or multiple answers give zero points. **Only** answers **on the answer sheet** count.

4. Which is a parametrisation of the following ellipse, with $t \in [0, 2\pi]$?



(a) $\begin{pmatrix} x(t) \\ y(t) \end{pmatrix} = \begin{pmatrix} 2 \sin t \\ 1 + 4 \cos t \end{pmatrix}.$

(c) $\begin{pmatrix} x(t) \\ y(t) \end{pmatrix} = \begin{pmatrix} \frac{\sin t}{2} \\ \frac{\cos t - 1}{4} \end{pmatrix}.$

(b) $\begin{pmatrix} x(t) \\ y(t) \end{pmatrix} = \begin{pmatrix} 2 \sin t \\ \cos t - 1 \end{pmatrix}.$

(d) $\begin{pmatrix} x(t) \\ y(t) \end{pmatrix} = \begin{pmatrix} \frac{1}{2} \sin t \\ \frac{1}{4} + \cos t \end{pmatrix}.$

8. For which b is the double integral

$$\int_0^b \int_0^\pi y \sin x \, dx \, dy$$

equal to 2?

(a) $b = \frac{\sqrt{2}}{2}$.

(c) $b = 2\sqrt{2}$.





(b) $b = \sqrt{2}$.

(d) $b = \frac{1}{2}$.

9. Consider the integral

$$\int_0^1 \int_0^2 \int_0^{2-2y} f(x, y, z) \, dz \, dx \, dy.$$

What is the corresponding region of integration?



10. The integral

$$\iint_B \sqrt{4 - x^2 - y^2} \, dx \, dy$$


over the region

$$B = \{(x, y) \mid x, y \leq 0, 1 \leq x^2 + y^2 \leq 4\}$$

is equal to

- (a) $\frac{\pi\sqrt{3}}{4}$. (b) $\frac{3\pi}{2}$. (c) $\frac{3\pi}{4}$. (d) $\frac{\pi\sqrt{3}}{2}$.

11. Which vector field corresponds to this drawing?



- (a) $\vec{F} = \begin{pmatrix} x + y \\ x^2 \end{pmatrix}$ (c) $\vec{F} = \begin{pmatrix} -x \\ 2y \end{pmatrix}$
 (b) $\vec{F} = \begin{pmatrix} x^2 - y^2 \\ x - y \end{pmatrix}$ (d) $\vec{F} = \begin{pmatrix} y \\ x^2 + y^2 \end{pmatrix}$

12. Which of the following vector fields \vec{F} has a potential on \mathbb{R}^2 ?

(a) $\vec{F}(x, y) = \begin{pmatrix} y - x \\ x - y \end{pmatrix}$.

(c) $\vec{F}(x, y) = \begin{pmatrix} x^2 - y \\ 2xy \end{pmatrix}$.

(b) $\vec{F}(x, y) = \begin{pmatrix} x - y \\ x + y \end{pmatrix}$.

(d) $\vec{F}(x, y) = \begin{pmatrix} y - x^2 \\ x^2 + y \end{pmatrix}$.

13. On which of the following regions is the vector field


$$\vec{G}(x, y) = \frac{1}{(x-2)^2 + (y-3)^2} \begin{pmatrix} -y+3 \\ x-2 \end{pmatrix}$$

a gradient field?

Hint: The vorticity of \vec{G} is zero for all $(x, y) \neq (2, 3)$.

- (a) In the first quadrant without $(2, 3)$.
- (b) In the halfplane $y \leq 0$.
- (c) In the annulus $1 \leq (x-2)^2 + (y-3)^2 \leq 2$.
- (d) In the plane \mathbb{R}^2 without the line segment from the origin to $(2, 3)$.

14. Which of the following parametrisations (where (u, v) varies on a square) describes the depicted surface?



- (a) $\vec{r}(u, v) = \begin{pmatrix} u \\ v \\ e^{-u^2-v^2} \end{pmatrix}$
- (b) $\vec{r}(u, v) = \begin{pmatrix} u \\ v \\ \sqrt{u^2 + v^2} \end{pmatrix}$
- (c) $\vec{r}(u, v) = \begin{pmatrix} u \\ v \\ \sin(u^2 + v^2) \end{pmatrix}$
- (d) $\vec{r}(u, v) = \begin{pmatrix} u \\ v \\ 1 - u^2 - v^2 \end{pmatrix}$

15. What is the outward flux of the vector field

$$\vec{F}(x, y) = \begin{pmatrix} 2x - x \cos(xy) \\ x + y \cos(xy) \end{pmatrix}$$

through the circle $x^2 + y^2 = 1$?

- (a) π (b) -2π (c) 2π (d) $-\pi$