

COMPLETE CLASSIFICATION OF GLOBAL SOLUTIONS TO THE OBSTACLE PROBLEM

SIMON EBERLE¹, ALESSIO FIGALLI², AND GEORG S. WEISS³

ABSTRACT. The characterization of global solutions to the obstacle problems in \mathbb{R}^N , or equivalently of null quadrature domains, has been studied for more than 90 years. In this paper, we give a conclusive answer to this problem by proving the following long-standing conjecture: *The coincidence set of a global solution to the obstacle problem is either a half-space, an ellipsoid, a paraboloid, or a cylinder with an ellipsoid or a paraboloid as base.*

CONTENTS

1. Introduction	1
1.1. Null quadrature domains and the obstacle problem	1
1.2. Classification results	2
1.3. Structure of the paper	4
2. Notation and preliminaries	5
3. The Newtonian potential expansion	9
4. Improved estimate on the asymptotic behavior of the coincidence set \mathcal{C}	13
5. Linear and almost-linear behavior of $V_{\mathcal{C}}$	22
6. Constructing matching paraboloid solutions	27
7. Proof of Theorem 1.2: the case $N \geq 4$	32
8. Proof of Theorem 1.2: the case $N = 3$	33
8.1. Linear rescaling and ACF dichotomy	34
8.2. Fine adjustment of u_{∞} at large scales	36
8.3. Ordering of solutions and conclusion	40
9. Proof of Theorem 1.1	41
Appendix A. Proof of Lemma 6.1	43
References	45

1. INTRODUCTION

1.1. Null quadrature domains and the obstacle problem. In 1678, Newton stated his famous *no gravity in the cavity theorem*: spherical shells do not exert gravitational force inside the cavity of the shell. This result was later extended to ellipsoidal shells (homoeoid) first by Laplace, and soon after by Ivory using a more geometric approach.

¹BASQUE CENTER OF APPLIED MATHEMATICS, SPAIN

²DEPARTMENT OF MATHEMATICS, ETH ZÜRICH, SWITZERLAND

³FACULTY OF MATHEMATICS, UNIVERSITY OF DUISBURG-ESSEN, GERMANY

E-mail addresses: seberle@bcamath.org, alessio.figalli@math.ethz.ch, georg.weiss@uni-due.de.

In modern terms, these results can be stated in terms of null quadrature domains. We recall that an open set $\Omega \subset \mathbb{R}^N$ is called a *null quadrature domain* if

$$\int_{\Omega} h \, dx = 0$$

for every harmonic function $h \in L^1(\Omega) \cap C^0(\overline{\Omega})$. With this terminology, the results of Newton, Laplace, and Ivory can be stated saying the complement of a ball/ellipsoid is a null quadrature domain. In greater generality, one can prove that null quadrature domains include:

- half-spaces;
- exteriors of ellipsoids;
- exterior of paraboloids;
- cylinders over domains of the types listed above.

A major question, which has been investigated over the last 90 years, is to understand whether this list includes all the possible null quadrature domains. Before discussing it, it is important to point out that null quadrature domains are related to solutions to the obstacle problem. More precisely, as discussed for instance in [4, Theorem II] and [14, Theorem 4.1],

Ω is a null quadrature domain \Leftrightarrow

$$\Omega = \{u > 0\} \text{ for some non-negative solution } u \in C_{\text{loc}}^{1,1}(\mathbb{R}^N) \text{ of } \Delta u = \chi_{\{u>0\}}.$$

In other words, characterizing null quadrature domains is equivalent to characterizing the coincidence set $\{u = 0\}$ for global solutions to the obstacle problem

$$\Delta u = \chi_{\{u>0\}}, \quad u \geq 0, \quad \text{in } \mathbb{R}^N. \quad (1.1)$$

It is well-known that global solutions to the obstacle problem are convex (see for instance [17, Theorem 5.1]). In particular, the coincidence set $\{u = 0\}$ is convex.

1.2. Classification results. The first partial classification of global solutions with *compact* coincidence sets has been achieved more than 90 years ago: in 1931, Dives [6] showed that, for $N = 3$, if $\{u = 0\}$ has non-empty interior and is bounded then it is an ellipsoid. Many years later, in 1979, Lewy gave a new proof of this result [16].

In 1981, Sakai gave a full classification of global solutions in *two dimensions* using complex analysis (cf. [18]).

The higher dimensional analogue to Dive's result, i.e., if $\{u = 0\}$ is bounded and has non-empty interior then it is an ellipsoid, was proved shortly after in two steps. First, in [5], DiBenedetto and Friedman proved the result in 1986 under the additional assumption that $\{u = 0\}$ is symmetric with respect to the hyperplanes $\{x_j = 0\}$ for all $j \in \{1, \dots, N\}$. Then, in the same year, Friedman and Sakai [10] removed the symmetry assumption. Very recently, in [9], two of the authors gave a concise proof of the characterization of compact coincidence sets.

Hence, while global solutions with compact coincidence sets had been completely classified, the structure of solutions with *unbounded* coincidence sets remained largely open and is related to the following conjecture (here, one is implicitly assuming that $\{u = 0\}$ has non-empty interior, as otherwise solutions are trivially classified, see Remark 2.5 below):

Conjecture: *The coincidence set of a global solution to the obstacle problem is either a half-space, an ellipsoid, a paraboloid, or a cylinder with an ellipsoid or a paraboloid as base.*

This conjecture, which has been investigated over more than 30 years, has been officially raised in several papers: first by Shahgholian in [19, conjecture on p. 10], then by Karp

and Margulis in [13, Conjecture 4.5], and recently in the monograph ‘Research Problems in Function Theory’ [11, §3.1 pp. 63-64, and Problem 3.28].

In the recent work [7] the first and third author, together with Shahgholian, have been able to solve the conjecture in the case of “ x_N -monotone” solutions when $N \geq 6$ (see Definition 2.8 and Remark 1.4 below).

Here we are able to fully characterize global solutions in the remaining dimensions (actually, our proof gives a complete characterization for all dimensions $N \geq 3$), allowing us to prove the conjecture above in full generality. Here is our main result:

Theorem 1.1 (Complete characterization of global solutions to the obstacle problem). *Let $N \geq 2$, and let u be a solution of (1.1) such that the coincidence set $\{u = 0\}$ has non-empty interior. Then the coincidence set is either a half-space, an ellipsoid, a paraboloid, or a cylinder with an ellipsoid or a paraboloid as base.*

As we shall explain later, this result is a rather direct consequence of the classification of x_N -monotone solutions. More precisely, the core of this paper is the following:

Theorem 1.2 (Characterization of monotone solutions). *Let $N \geq 3$ and let u be an x_N -monotone solution according to Definition 2.8 below. Then $\{u = 0\}$ is a paraboloid.*

Remark 1.3. Thanks to [8, Main Theorem], Theorem 1.1 implies a fine result on the behavior of the regular part of the free boundary close to singularities.

Remark 1.4. As mentioned before, Theorem 1.2 has already been proved for $N \geq 6$ in [7]. A reason for this dimensional restriction comes from the fact that, in the proof in [7], a key role is played by the Newtonian potential associated to $\{u = 0\}$, defined (up to a multiplicative constant) as $\frac{1}{|x|^{N-2}} * \chi_{\{u=0\}}$. However, if $\{u = 0\}$ is a paraboloid then the above convolution converges only for $N \geq 6$.

As we shall see later, this definition of Newtonian potential can be “corrected” to obtain a convergent expression also in lower dimensions (see Definition 3.1 and Lemma 3.3). However, the positivity of the Newtonian potential is important for the arguments in [7], while our generalized potential loses this property.

At a more “fundamental” level, the role of the dimension can be seen as follows: if $p(x) = \lim_{r \rightarrow \infty} \frac{u(rx)}{r^2}$ denotes the blow-down polynomial appearing in Definition 2.8(iv), then the behaviour of $u - p$ changes considerably with the dimension. In particular, if $\{u = 0\}$ is a paraboloid (this is a particular case of x_N -monotone solution) then one can check by explicit computations that, for $N \geq 4$, there exists a linear function ℓ such that:

- $\int_{B_R} |u - p - \ell| dx$ is bounded for $N \geq 6$;
- $\int_{B_R} |u - p - \ell| dx \simeq \log(R)$ for $N = 5$;
- $\int_{B_R} |u - p - \ell| dx \simeq \sqrt{R}$ for $N = 4$.

This different behavior is the reason for the dimensional restriction $N \geq 6$ in [7]. In this paper, instead, we develop a new approach that only requires $\int_{B_R} |u - p - \ell| dx = o(R)$, giving a unified proof of Theorem 1.2 for $N \geq 4$. Unfortunately, in the “critical” dimension $N = 3$, $\int_{B_R} |u - p| dx \simeq R \log R$. In particular, there is no affine function that dictates the behaviour at infinity of $u - p$. As the reader will see, this fact is a source of major difficulties for proving Theorem 1.2 in dimension $N = 3$.

We note that also for $N = 2$ the behavior of $u - p$ is superlinear: $\int_{B_R} |u - p| dx \simeq R^{3/2}$. However, when $N = 2$ one can rely on the Riemann mapping theorem to obtain a short proof of Theorem 1.1 (see [18]).

1.3. Structure of the paper. In Section 2 we introduce some notation and collect a series of useful preliminary estimates on solutions to the obstacle problem and on the Alt-Caffarelli-Friedman (ACF) functional that will play a crucial role in our proof.

In Section 3 we prove that, for an x_N -monotone solution u , one can define a Newtonian-type potential $V_{\mathcal{C}}$ associated with its coincidence set $\mathcal{C} = \{u = 0\}$ so that the expansion $u = p + V_{\mathcal{C}}$ holds.

Thanks to the Newtonian expansion proved in Section 3, in Section 4 we show that the coincidence set \mathcal{C} is asymptotically contained inside a paraboloid. The proof of this result is rather easy in dimension $N \geq 4$, while the case $N = 3$ requires an extremely delicate argument.

In Section 5, we use the result from Section 4 to analyze the asymptotic behavior of $V_{\mathcal{C}}$. In particular: for $N \geq 4$ we can show that $V_{\mathcal{C}}$ behaves at infinity like a linear function (up to sublinear corrections); for $N = 3$, on each large ball B_R , $V_{\mathcal{C}}$ is at most CR -away from an affine function whose slope behaves like $\log R$. In other words, while for $N \geq 4$ the gradient of $V_{\mathcal{C}}$ is bounded, for $N = 3$ it has a BMO-type behavior (see Lemma 5.2).

In Section 6, exploiting the information on $V_{\mathcal{C}}$ obtained in the previous section, we can construct matching paraboloid solutions (i.e., solutions that have paraboloids as coincidence sets). More precisely, for $N \geq 4$ we can find a paraboloid solution u_P such that $u - u_P$ grows sublinearly at infinity. Instead, for $N = 3$, for each R we construct a paraboloid solution u_{P_R} such that $\frac{1}{|B_R|} \|u - u_{P_R}\|_{L^1(B_R)} \leq CR$. With all this preparatory work, we can then prove our main result.

More precisely, in Section 7 we focus on the case $N \geq 4$. In that case, applying the ACF formula to the difference between u and suitable translations of the paraboloid solution constructed in Section 6, and exploiting the sublinear growth at infinity, we are able to prove that such solutions are ordered. Once this is achieved, we conclude easily.

Then, in Section 8 we focus on the case $N = 3$. In this case, due to the lack of a sublinear approximation of u via paraboloid solutions, we cannot directly apply the ACF formula to deduce that u and some suitable paraboloid solutions are ordered. Instead, we apply the ACF formula to the functions $\frac{1}{R}(u - u_{P_R})(R \cdot)$ to construct a comparison solution u_{∞} whose coincidence set is a paraboloid. Then, by a delicate ACF-type dichotomy, we show that one-homogeneous blow-down limits of $u - u_{\infty}$ exist and:

- either they have constant sign (so u and u_{∞} are “ordered at infinity”);
- or they are linear functions.

While in the first case we can conclude similarly to the case $N \geq 4$, the second case requires a refined analysis. More precisely, exploiting the information that $u - u_{\infty}$ behaves as a linear function at infinity, we can construct fine adjustments of the paraboloid solution u_{∞} to show that, for some suitable translations of u_{∞} , the ACF energy vanishes. Then, we conclude similarly to the first case.

For completeness, in Section 9 we provide a new self-contained argument showing how Theorem 1.1 follows from Theorem 1.2.

Acknowledgements. The second author has received funding from the European Research Council under the Grant Agreement No. 721675 “Regularity and Stability in Partial Differential Equations (RSPDE)”. We thank Lili Du and the anonymous referees for several useful comments on a preliminary version of this manuscript.

2. NOTATION AND PRELIMINARIES

Throughout this work, \mathbb{R}^N will be equipped with the Euclidean inner product $x \cdot y$ and the induced norm $|x|$. Due to the nature of the problem, we will often write $x \in \mathbb{R}^N$ as $x = (x', x_N) \in \mathbb{R}^{N-1} \times \mathbb{R}$. Also, we denote by $(e^i)_{1 \leq i \leq N}$ the elements of the canonical base of \mathbb{R}^N .

In our estimates, C denotes a generic positive constant that may change from line to line. We shall use C_N whenever the constant depends only on the dimension.

We write $B_r(x)$ to denote the open N -dimensional ball of center x and radius r , while $B'_r(x')$ is the open $(N-1)$ -dimensional ball of center $x' \in \mathbb{R}^{N-1}$ and radius r . Whenever the center is omitted, it is assumed to be the origin 0 .

When considering a set A , χ_A shall denote the characteristic function of A . With \mathcal{H}^k we refer to k -dimensional Hausdorff measure. If A and B are two sets, we denote their symmetric difference by $A \Delta B := (A \setminus B) \cup (B \setminus A)$. Given a function $f : \mathbb{R}^N \rightarrow \mathbb{R}$, we define $f_+ := \max\{f, 0\}$ and $f_- := \max\{-f, 0\}$. Furthermore, we define the differential operator $\nabla' f := (\partial_1 f, \dots, \partial_{N-1} f)$.

Definition 2.1 (Coincidence set). *Given a solution u to the obstacle problem (1.1), we define its coincidence set \mathcal{C} to be*

$$\mathcal{C} := \{u = 0\}.$$

Remark 2.2. As already mentioned before, global solutions to the obstacle problem are convex (see e.g. [17, Theorem 5.1]). In particular, the coincidence set \mathcal{C} of a global solution is convex.

To get compactness of solutions, it is useful to recall that they are uniformly $C^{1,1}$ -regular. Also, as shown by Caffarelli, their blow-down limits with respect to quadratic rescaling are either half-space solutions or quadratic polynomials (see [3]). We summarize these results in the following two lemmas:

Lemma 2.3 (Characterization of blow-down limits). *Let $u : \mathbb{R}^N \rightarrow [0, \infty)$ be a global solution to the obstacle problem. Then the following convergence holds in $C_{\text{loc}}^{1,\alpha}(\mathbb{R}^N)$ for each $\alpha \in (0, 1)$:*

$$\lim_{r \rightarrow \infty} \frac{u(rx)}{r^2} = \begin{cases} \frac{1}{2} \max(x \cdot e, 0)^2 & \text{for some } e \in \partial B_1, \\ \frac{1}{2} x^T Q x & \text{for some } Q \in \mathbb{R}^{N \times N} \text{ symmetric,} \\ & \text{non-negative definite, satisfying } \text{tr}(Q) = 1. \end{cases}$$

A global solution of the form $\frac{1}{2} \max(x \cdot e, 0)^2$ is called half-space solution.

Lemma 2.4 (Uniform regularity and compactness). *The following regularity and compactness properties hold:*

- (i) *Let u be a global solution to the obstacle problem in \mathbb{R}^N . Then $\|D^2 u\|_{L^\infty(\mathbb{R}^N)} \leq C_N$.*
- (ii) *Let $(u_k)_{k \in \mathbb{N}}$ be a sequence of global solutions to the obstacle problem in \mathbb{R}^N that vanish at the origin. Then there exists a subsequence $(u_{k_j})_{j \in \mathbb{N}}$ converging to a global solution u_0 in $C_{\text{loc}}^{1,\alpha}(\mathbb{R}^N)$ for each $\alpha \in (0, 1)$. In addition, $\chi_{\{u_{k_j}=0\}} \rightarrow \chi_{\{u_0=0\}}$ a.e. in \mathbb{R}^N .*

Proof. Since u is convex it follows that $0 \leq \partial_{ee} u \leq \Delta u = \chi_{\{u>0\}} \leq 1$ for all $e \in \partial B_1$. This proves (i).

Concerning (ii), we note that the compactness in $C_{\text{loc}}^{1,\alpha}(\mathbb{R}^N)$ is a direct consequence of (i) and Ascoli-Arzelà Theorem. The a.e. convergence of the characteristic functions of the contact sets follows from [17, Proposition 3.17(i)-(ii)]. \square

Remark 2.5. As noted in the previous proof, global solutions grow at most quadratically at infinity (cf. [17, Theorem 2.1]). Also, if the convex set $\{u = 0\}$ has empty interior, then $\Delta u \equiv 1$. Hence, Liouville's theorem implies that the only global solutions whose coincidence sets have empty interior are quadratic polynomials.

Within the class of global solutions to the obstacle problem, we now introduce some terminology for denoting some special solutions

Definition 2.6 (Cylindrical solutions). *We say that a global solution to the obstacle problem is cylindrical if there exists $e \in \partial B_1$ such that*

$$\nabla u \cdot e \equiv 0 \quad \text{in } \mathbb{R}^N.$$

A useful criterion for being a cylindrical solution is contained in the following:

Lemma 2.7. *Let u be a global solution, and assume that its coincidence set \mathcal{C} contains an infinite line. Then u is constant in the direction of that line.*

Proof. First of all we may assume that \mathcal{C} has non-empty interior, as otherwise u is a non-negative quadratic polynomial (see Remark 2.5) and the result follows easily.

Since \mathcal{C} is convex, the assumption of containing a line implies that \mathcal{C} is a product, namely there exists a system of coordinates such that $\mathcal{C} = \mathcal{K} \times \mathbb{R}$ for some convex set $\mathcal{K} \subset \mathbb{R}^{N-1}$.¹ Hence, given $\sigma \in \mathbb{R}$, the global solution $u_\sigma(x) := u(x + \sigma e^N)$ has the same contact set as u , and therefore $\Delta(u - u_\sigma) \equiv 0$. Since $u - u_\sigma$ vanishes on \mathcal{C} which has non-empty interior, it follows by unique continuation that $u - u_\sigma \equiv 0$. Since σ is arbitrary, this shows that u is invariant in the e^N -direction, proving the result. \square

Definition 2.8 (x_N -monotone solutions). *We say that a global solution to the obstacle problem (1.1) is x_N -monotone if:*

- (i) \mathcal{C} has non-empty interior;
- (ii) $\mathcal{C} \subset \{x_N \geq 0\}$ and $\mathcal{C} \cap \{x_N = 0\} = \{0\}$;
- (iii) $\partial_N u \leq 0$ in \mathbb{R}^N ;
- (iv) $\lim_{r \rightarrow \infty} \frac{u(rx)}{r^2} = \frac{1}{2} x'^T Q x' =: p(x')$ in $C_{\text{loc}}^{1,\alpha}(\mathbb{R}^N)$ for each $\alpha \in (0, 1)$, where $x = (x', x_N)$, and $Q \in \mathbb{R}^{(N-1) \times (N-1)}$ is symmetric, positive definite, and satisfies $\text{tr}(Q) = 1$.

Remark 2.9. Thanks to Definition 2.8(ii)-(iii), if u is x_N -monotone then $\{te^N : t \geq 0\} \subset \mathcal{C}$. Also, since the matrix $Q \in \mathbb{R}^{(N-1) \times (N-1)}$ in Definition 2.8(iv) is positive definite, there exists a constant $c_p > 0$ such that

$$p(x') \geq c_p |x'|^2 \quad \text{for all } x' \in \mathbb{R}^{N-1}. \quad (2.1)$$

The following important result on x_N -monotone solutions is proved in [7, Proposition 5.1] and will be used in Lemma 3.3 to prove that the generalized Newtonian potential associated to the contact set of an x_N -monotone solution is well-defined.

¹This classical fact can be proved as follows. Assume that the line ℓ is parallel to e^N , say $\ell = \{\bar{x} + se^N : s \in \mathbb{R}\}$ for some $\bar{x} \in \mathbb{R}^N$, and define $K_\tau := \mathcal{C} \cap \{x_N = \tau\}$. Let $\text{conv}(A)$ denote the convex hull of the set A . Then, by convexity of \mathcal{C} , $\text{conv}(K_\tau \cup \ell) \subset \mathcal{C}$ for any $\tau \in \mathbb{R}$. Since $\text{conv}(K_\tau \cup \ell) = K_\tau \times \mathbb{R}$, it follows that $\mathcal{C} \supset \mathcal{K} \times \mathbb{R}$ with $\mathcal{K} \times \mathbb{R} := \bigcup_\tau (K_\tau \times \mathbb{R})$. On the other hand, it is clear by construction that $\mathcal{C} \subset \mathcal{K} \times \mathbb{R}$, so the result follows.

Lemma 2.10 (\mathcal{C} is “almost contained” in a paraboloid). *Let $N \geq 3$, and let u be an x_N -monotone solution. Fix $\delta \in (0, 1)$, and define $T_\delta := \{(y', y_N) \in \mathbb{R}^N : |y'|^2 < y_N^{1+\delta}\}$. Then there exists a radius $\hat{r} > 1$ such that*

$$\mathcal{C} \setminus B_{2\hat{r}} \subset \{y_N > \hat{r}\} \cap T_\delta. \tag{2.2}$$

Definition 2.11 (Ellipsoids and Paraboloids). *We call a set $E \subset \mathbb{R}^N$ ellipsoid if, after a translation and a rotation,*

$$E = \left\{ x \in \mathbb{R}^N : \sum_{j=1}^N \frac{x_j^2}{a_j^2} \leq 1 \right\}$$

for some $a = (a_1, \dots, a_N) \in (0, \infty)^N$. We call a set $P \subset \mathbb{R}^N$ a paraboloid, if, after a translation and a rotation,

$$P = \{(x', x_N) \in \mathbb{R}^N : x_N \geq 0, x' \in \sqrt{x_N} E'\},$$

where E' is an $(N - 1)$ -dimensional ellipsoid.

An important role in this paper will be played by the Alt-Caffarelli-Friedman (ACF) functional originally introduced in [1]: given a function $v : \mathbb{R}^N \rightarrow \mathbb{R}$ with $N \geq 2$, we define

$$\Phi(v, r) := \frac{1}{r^4} \int_{B_r} \frac{|\nabla v_+|^2}{|x|^{N-2}} dx \int_{B_r} \frac{|\nabla v_-|^2}{|x|^{N-2}} dx. \tag{2.3}$$

We recall in the following lemma some useful facts about the ACF functional.

Lemma 2.12 (Properties of the Alt-Caffarelli-Friedman monotonicity functional). *Let $N \geq 2$, and let $v : \mathbb{R}^N \rightarrow \mathbb{R}$ be a continuous $W_{\text{loc}}^{1,2}$ function such that both v_+ and v_- are subharmonic. Then:*

(i) *The functional $\Phi(v, r)$ is finite for each $r > 0$, and*

$$r \mapsto \Phi(v, r) \quad \text{is non-decreasing.}$$

(ii) *The following bound holds for any $r > 0$:*

$$\int_{B_r} \frac{|\nabla v_\pm|^2}{|x|^{N-2}} dx \leq C_N \left(\int_{B_{4r}} v_\pm dx \right)^2.$$

(iii) *The following bound holds for any $r > 0$:*

$$\Phi(v, r) \leq \frac{C_N}{r^4} \left(\int_{B_{4r}} v_+ dx \right)^2 \left(\int_{B_{4r}} v_- dx \right)^2.$$

(iv) *Assume that $\Phi(v, R) \rightarrow 0$ as $R \rightarrow \infty$. Then either $v \geq 0$ in \mathbb{R}^N or $v \leq 0$ in \mathbb{R}^N .*

Proof. Usually (i) is stated and proved under the extra assumption $v(0) = 0$. However, as noted in [17, Theorem 2.4], this extra condition is not needed and therefore (i) holds in our setting.

By Hölder’s inequality and subharmonicity of v_\pm , we can estimate

$$\|v_\pm\|_{L^2(B_{2r})}^2 \leq \|v_\pm\|_{L^\infty(B_{2r})} \|v_\pm\|_{L^1(B_{2r})} \quad \text{and} \quad \|v_\pm\|_{L^\infty(B_{2r})} \leq \frac{C_N}{r^N} \|v_\pm\|_{L^1(B_{4r})}. \tag{2.4}$$

Also, as noted in [17, Section 2.2.3], the bound

$$\int_{B_r} \frac{|\nabla w|^2}{|x|^{N-2}} dx \leq C_N \left(\int_{B_{2r}} w^2 dx \right)$$

holds for any non-negative subharmonic function w . Applying this inequality to $w = v_{\pm}$ and using (2.4), we obtain (ii).

Multiplying the two estimates in (ii) (the one for v_+ with the one for v_-), (iii) follows.

To prove (iv) we can assume that there is a point $y \in \mathbb{R}^N$ such that $v(y) = 0$ as otherwise, by continuity of v , either $v > 0$ or $v < 0$ and the result is trivially true. By the monotonicity and non-negativity of Φ , our assumption implies that $\Phi(v, r) \equiv 0$ for all $r > 0$. Hence, by the definition of the ACF functional (cf. (2.3)), for each $r \in (0, \infty)$,

$$\text{either } \nabla v_+ \equiv 0 \text{ in } B_r \text{ or } \nabla v_- \equiv 0 \text{ in } B_r.$$

Since by assumption $v(y) = 0$ we deduce that, for all $r > |y|$,

$$\text{either } v_+ \equiv 0 \text{ in } B_r \text{ or } v_- \equiv 0 \text{ in } B_r.$$

Therefore, by continuity,

$$\text{either } v_+ \equiv 0 \text{ in } \bigcup_{r>|y|} B_r = \mathbb{R}^N \text{ or } v_- \equiv 0 \text{ in } \bigcup_{r>|y|} B_r = \mathbb{R}^N,$$

which proves (iv). \square

We conclude this section with a couple of simple but important results on the difference of two global solutions. These results will play a crucial role in the proof of Theorem 1.2 where we will apply the ACF functional to the difference of two global solutions.

Lemma 2.13 (Subharmonicity properties and Caccioppoli estimate). *Let $u_1, u_2 : \mathbb{R}^N \rightarrow \mathbb{R}$ be global solutions to the obstacle problem. Then the following hold:*

- (i) *The functions $(u_1 - u_2)_+$, $(u_1 - u_2)_-$, and $|u_1 - u_2|$ are subharmonic.*
- (ii) *The following bound holds for any $r > 0$:*

$$\int_{B_r} |\nabla(u_1 - u_2)|^2 dx \leq \frac{C_N}{r^2} \int_{B_{2r}} (u_1 - u_2)^2 dx.$$

Proof. Set $w := u_1 - u_2$ and note that, since $\Delta u_i = \chi_{\{u_i > 0\}}$,

$$\Delta w = \chi_{\{u_1 > 0\}} - \chi_{\{u_2 > 0\}} \geq 0 \quad \text{inside } \{u_1 > u_2\}. \quad (2.5)$$

Choosing a sequence of smooth, convex, non-decreasing functions $\varphi_\varepsilon : \mathbb{R} \rightarrow \mathbb{R}$ such that $\varphi_\varepsilon|_{(-\infty, 0)} \equiv 0$ and $\varphi_\varepsilon(s) \rightarrow s_+$ as $\varepsilon \rightarrow 0$ locally uniformly, we see that

$$\Delta[\varphi_\varepsilon(w)] = \varphi'_\varepsilon(w) [\chi_{\{u_1 > 0\}} - \chi_{\{u_2 > 0\}}] + \varphi''_\varepsilon(w) |\nabla w|^2 \geq 0.$$

Letting $\varepsilon \rightarrow 0$, we conclude that $(u_1 - u_2)_+$ is subharmonic. Since $(u_1 - u_2)_- = (u_2 - u_1)_+$, by symmetry between u_1 and u_2 we deduce that $(u_1 - u_2)_-$ is subharmonic. Finally, since $|u_1 - u_2| = (u_1 - u_2)_+ + (u_1 - u_2)_-$, the subharmonicity of $|u_1 - u_2|$ follows. This proves (i).

To prove (ii) we define $w_r(x) := \frac{w(rx)}{r^2}$ and we note that, as a consequence of (2.5), it holds $w_r \Delta w_r \geq 0$, or equivalently

$$\Delta(w_r^2) \geq 2|\nabla w_r|^2. \quad (2.6)$$

Now, fix $\eta \in C_c^\infty(B_2)$ a non-negative cut-off function satisfying $\eta \equiv 1$ in B_1 . Integrating the inequality (2.6) against η we obtain

$$\int_{B_1} |\nabla w_r|^2 dx \leq \int_{B_2} |\nabla w_r|^2 \eta dx \leq \frac{1}{2} \int_{B_2} w_r^2 \Delta \eta dx \leq C \int_{B_2} w_r^2 dx,$$

as desired. □

Remark 2.14. As a direct consequence of Lemma 2.13(i) we recover the well-known fact that, given a global solution u , $(\partial_e u)_+$ and $(\partial_e u)_-$ are subharmonic for each $e \in \partial B_1$. Indeed, given $h > 0$ it suffices to apply Lemma 2.13(i) to u and $u(\cdot + he)$ to deduce that both $\left(\frac{u(\cdot + he) - u}{h}\right)_+$ and $\left(\frac{u(\cdot + he) - u}{h}\right)_-$ are subharmonic, and then the result follows by letting $h \rightarrow 0$.

Lemma 2.15 (Strong convergence). *Let $\varrho \in (0, \infty)$ and let $(u_k)_{k \in \mathbb{N}}, (v_k)_{k \in \mathbb{N}}$ be two sequences of global solutions to the obstacle problem in \mathbb{R}^N such that*

$$w_k := u_k - v_k \rightharpoonup w \quad \text{weakly in } W^{1,2}(B_\varrho) \text{ as } k \rightarrow \infty,$$

for some harmonic function $w : \mathbb{R}^N \rightarrow \mathbb{R}$. Then, for each $\delta \in (0, 1)$,

$$w_k = u_k - v_k \rightarrow w \quad \text{strongly in } W^{1,2}(B_{\delta\varrho}) \text{ as } k \rightarrow \infty.$$

Proof. First of all note that, for all $k \in \mathbb{N}$,

$$w_k \Delta w_k = (u_k - v_k)(\chi_{\{u_k > 0\}} - \chi_{\{v_k > 0\}}) \geq 0.$$

Hence, given $\eta \in C_c^\infty(B_\varrho; [0, \infty))$ satisfying $\eta \equiv 1$ in $B_{\delta\varrho}$, integrating by parts twice we get

$$\begin{aligned} \int_{B_{\delta\varrho}} |\nabla w_k|^2 &\leq \int_{B_\varrho} |\nabla w_k|^2 \eta = - \int_{B_\varrho} (w_k \nabla w_k \cdot \nabla \eta + \eta w_k \Delta w_k) \leq - \int_{B_\varrho} w_k \nabla w_k \cdot \nabla \eta \\ &\rightarrow - \int_{B_\varrho} w \nabla w \cdot \nabla \eta = \int_{B_\varrho} |\nabla w|^2 \eta \quad \text{as } k \rightarrow \infty, \end{aligned}$$

where the last equality follows by the harmonicity of w .

Now, choosing a sequence $(\eta_j)_{j \in \mathbb{N}} \subset C_c^\infty(B_\varrho; [0, \infty))$ such that $\eta_j \equiv 1$ in $B_{\delta\varrho}$ for all $j \in \mathbb{N}$ and $\eta_j \rightarrow \chi_{B_{\delta\varrho}}$ pointwise in B_ϱ , we conclude that

$$\limsup_{k \rightarrow \infty} \int_{B_{\delta\varrho}} |\nabla w_k|^2 \leq \int_{B_{\delta\varrho}} |\nabla w|^2.$$

Therefore, by the lower-semicontinuity of the Dirichlet energy we deduce that $\|\nabla w_k\|_{L^2(B_{\delta\varrho})} \rightarrow \|\nabla w\|_{L^2(B_{\delta\varrho})}$. This convergence of the L^2 -norm of the gradients together with the weak convergence implies the desired strong convergence. □

3. THE NEWTONIAN POTENTIAL EXPANSION

As mentioned in Remark 1.4, in [7] a very important role is played by the Newtonian potential associated to the coincidence set of a solution, defined (up to a multiplicative constant) as $\frac{1}{|x|^{N-2}} * \chi_{\mathcal{C}}$. Unfortunately, if \mathcal{C} is a paraboloid then the above convolution converges only for $N \geq 6$. For this reason we will introduce a generalized Newtonian potential in the spirit of [12], which will be shown in Lemma 3.3 to be well-defined and to have subquadratic growth.

Definition 3.1 (Generalized Newtonian potential). *Let $N \geq 3$, and define the function*

$$G(x, y) := \frac{1}{|x - y|^{N-2}} - \frac{1}{|y|^{N-2}} - (N - 2) \frac{x \cdot y}{|y|^N} \quad \text{for all } x, y \in \mathbb{R}^N.$$

Given $M \subset \mathbb{R}^N$ measurable, assume that $G(x, \cdot)\chi_M \in L^1(\mathbb{R}^N)$ for each $x \in \mathbb{R}^N$. Then we define the generalized Newtonian potential associated to M as

$$V_M(x) := \alpha_N \int_M G(x, y) \, dy, \quad \text{where } \alpha_N := \frac{1}{N(N-2)|B_1|}. \quad (3.1)$$

Lemma 3.2 (Scaling of the generalized Newtonian potential). *Let $M \subset \mathbb{R}^N$ be a measurable set for which V_M is well-defined. Then V_M satisfies the following scaling law:*

$$V_M(\gamma x) = \gamma^2 V_{\frac{1}{\gamma}M}(x) \quad \text{for all } \gamma > 0.$$

Proof. The proof follows from a direct calculation: since

$$G(\gamma x, y) = \gamma^{2-N} G\left(x, \frac{y}{\gamma}\right) \quad \text{for all } \gamma > 0, x, y \in \mathbb{R}^N,$$

$$\begin{aligned} V_M(\gamma x) &= \alpha_N \int_M G(\gamma x, y) \, dy = \gamma^{2-N} \alpha_N \int_M G\left(x, \frac{y}{\gamma}\right) \, dy \\ &= \gamma^{2-N} \alpha_N \int_{\frac{1}{\gamma}M} G(x, z) \gamma^N \, dz = \gamma^2 V_{\frac{1}{\gamma}M}(x). \end{aligned}$$

□

Lemma 3.3 (Generalized Newtonian potential of \mathcal{C}). *Let $N \geq 3$, and let u be an x_N -monotone solution in the sense of Definition 2.8. Then*

(i) *The generalized potential $V_{\mathcal{C}}$ of \mathcal{C} is well-defined and locally bounded.*

(ii) *$V_{\mathcal{C}}(x)$ grows subquadratically as $|x| \rightarrow \infty$. More precisely, there exists a constant C such that*

$$\frac{|V_{\mathcal{C}}(x)|}{(1 + |x|)^{7/4}} \leq C \quad \text{for all } x \in \mathbb{R}^N.$$

(iii) *$V_{\mathcal{C}} \in W_{\text{loc}}^{2,p}(\mathbb{R}^N)$ for each $p \in [1, \infty)$, $\Delta V_{\mathcal{C}} = -\chi_{\mathcal{C}}$, and $V_{\mathcal{C}}(0) = |\nabla V_{\mathcal{C}}(0)| = 0$.*

Proof. Fix $\delta \in (0, 1)$, let T_{δ} be as in Lemma 2.10, and recall that (2.2) holds.

To prove the estimate, we first note that the trivial bound

$$|G(x, y)| \leq \frac{1}{|x - y|^{N-2}} + \frac{1}{|y|^{N-2}} + (N - 2) \frac{|x|}{|y|^{N-1}}$$

holds. Also, by the Taylor expansion $f(1) = f(0) + f'(0) + \int_0^1 (1 - \tau) f''(\tau) \, d\tau$ applied to $f(\tau) := \frac{1}{|\tau x - y|^{N-2}}$, we get

$$|G(x, y)| \leq C \frac{|x|^2}{|y|^N} \quad \text{for } |y| > 2|x|. \quad (3.2)$$

Using these two bounds and (2.2), we obtain (here $\hat{r} > 1$ is the radius provided by Lemma 2.10)

$$\begin{aligned}
 \int_{\mathcal{C}} |G(x, y)| \, dy &\leq \int_{\mathcal{C} \cap B_{2\hat{r}}} \left(\frac{1}{|x-y|^{N-2}} + \frac{1}{|y|^{N-2}} + (N-2) \frac{|x|}{|y|^{N-1}} \right) dy \\
 &\quad + \int_{\mathcal{C} \cap (B_{2|x|} \setminus B_{2\hat{r}})} \left(\frac{1}{|x-y|^{N-2}} + \frac{1}{|y|^{N-2}} + (N-2) \frac{|x|}{|y|^{N-1}} \right) dy \\
 &\quad + C \int_{\mathcal{C} \setminus (B_{2|x|} \cup B_{2\hat{r}})} \frac{|x|^2}{|y|^N} dy \\
 &\leq \int_{B_{2\hat{r}}} \left(\frac{1}{|x-y|^{N-2}} + \frac{1}{|y|^{N-2}} + (N-2) \frac{|x|}{|y|^{N-1}} \right) dy \\
 &\quad + \int_{T_\delta \cap (B_{2|x|} \setminus B_{2\hat{r}})} \left(\frac{1}{|x-y|^{N-2}} + \frac{1}{|y|^{N-2}} + (N-2) \frac{|x|}{|y|^{N-1}} \right) dy \\
 &\quad + C \int_{T_\delta \setminus (B_{2|x|} \cup B_{2\hat{r}})} \frac{|x|^2}{|y|^N} dy =: I_1 + I_2 + I_3.
 \end{aligned} \tag{3.3}$$

Since the integral of $\frac{1}{|x-y|^{N-2}}$ over a ball is maximized when x coincides with the center of the ball,² to bound the first integral I_1 we note that

$$\int_{B_{2\hat{r}}} \left(\frac{1}{|x-y|^{N-2}} + \frac{1}{|y|^{N-2}} \right) dy \leq 2 \int_{B_{2\hat{r}}} \frac{1}{|y|^{N-2}} dy \leq C \quad \text{and} \quad \int_{B_{2\hat{r}}} \frac{1}{|y|^{N-1}} dy \leq C,$$

therefore

$$I_1 \leq C(1 + |x|). \tag{3.4}$$

About I_2 , we note that this integral is nonzero only if $|x| > \hat{r}$. In such a case, we observe that, provided that \hat{r} is large enough,

$$T_\delta \cap (B_{2|x|} \setminus B_{2\hat{r}}) \subset T_\delta \cap \{\hat{r} < y_N < 2|x|\}.$$

Hence, noticing that $\frac{(N-1)(1+\delta)}{2} + 3 - N \leq 1 + \delta$ for $N \geq 3$, since $\frac{1}{|y|} \leq \frac{1}{y_N}$ we get

$$\begin{aligned}
 \int_{T_\delta \cap (B_{2|x|} \setminus B_{2\hat{r}})} \frac{1}{|y|^{N-2}} dy &\leq \int_{T_\delta \cap \{\hat{r} < y_N < 2|x|\}} \frac{1}{y_N^{N-2}} dy = \int_{\hat{r}}^{2|x|} \mathcal{H}^{N-1}(T_\delta \cap \{y_N = t\}) \frac{1}{t^{N-2}} dt \\
 &\leq C \int_{\hat{r}}^{2|x|} t^{\frac{(N-1)(1+\delta)}{2} + 2 - N} dt \leq C(1 + |x|)^{\frac{(N-1)(1+\delta)}{2} + 3 - N} \leq C(1 + |x|)^{1+\delta}, \tag{3.5}
 \end{aligned}$$

and analogously

$$\begin{aligned}
 \int_{T_\delta \cap (B_{2|x|} \setminus B_{2\hat{r}})} \frac{1}{|y|^{N-1}} dy &\leq C \int_{\hat{r}}^{2|x|} t^{\frac{(N-1)(1+\delta)}{2} + 1 - N} dt \\
 &\leq C(1 + |x|)^{\frac{(N-1)(1+\delta)}{2} + 2 - N} \leq C(1 + |x|)^\delta. \tag{3.6}
 \end{aligned}$$

²This follows, for instance, from the Hardy-Littlewood inequality

$$\int_{\mathbb{R}^n} f(y)g(y) \, dy \leq \int_{\mathbb{R}^n} f^*(y)g^*(y) \, dy, \quad f^*, g^* \text{ symmetric decreasing rearrangement of } f, g \geq 0,$$

applied with $f(y) = \frac{1}{|x-y|^{N-2}}$ and $g(y) = \chi_{B_r}(y)$. Indeed, in this case, $f^*(y) = \frac{1}{|y|^{N-2}}$ and $g^* = g$.

Therefore, combining (3.5) and (3.6),

$$\int_{T_\delta \cap (B_{2|x|} \setminus B_{2\hat{r}})} \left(\frac{1}{|y|^{N-2}} + (N-2) \frac{|x|}{|y|^{N-1}} \right) dy \leq C(1 + |x|)^{1+\delta}.$$

Finally, to estimate the integral of $\frac{1}{|x-y|^{N-2}}$, we use again that the integral of $\frac{1}{|x'-y'|^{N-2}}$ over a ball is maximized when x' coincides with the center of the ball. This yields

$$\int_{T_\delta \cap \{y_N = t\}} \frac{1}{|x' - y'|^{N-2}} dy' = \int_{B'_{t^{\frac{1+\delta}{2}}}} \frac{1}{|x' - y'|^{N-2}} dy' \leq \int_{B'_{t^{\frac{1+\delta}{2}}}} \frac{1}{|y'|^{N-2}} dy' = Ct^{\frac{1+\delta}{2}},$$

and therefore, since $\frac{1}{|x-y|} \leq \frac{1}{|x'-y'|}$,

$$\int_{T_\delta \cap (B_{2|x|} \setminus B_{2\hat{r}})} \frac{1}{|x-y|^{N-2}} dy \leq C \int_{\hat{r}}^{2|x|} t^{\frac{1+\delta}{2}} dt \leq C|x|^{\frac{3+\delta}{2}}.$$

Overall, this proves that $I_2 \leq C(1 + |x|)^{\frac{3+\delta}{2}}$.

Finally, for I_3 , we simply note that

$$T_\delta \setminus (B_{2|x|} \cup B_{2\hat{r}}) \subset T_\delta \cap \{y_N > \max\{|x|, \hat{r}\}\},$$

hence

$$\begin{aligned} \int_{T_\delta \setminus (B_{2|x|} \cup B_{2\hat{r}})} \frac{|x|^2}{|y|^N} dy &\leq |x|^2 \int_{\max\{|x|, \hat{r}\}}^\infty \mathcal{H}^{N-1}(T_\delta \cap \{y_N = t\}) \frac{1}{t^N} dt \\ &= C|x|^2 \int_{\max\{|x|, \hat{r}\}}^\infty t^{\frac{(N-1)(1+\delta)}{2} - N} dt \leq C|x|^2(1 + |x|)^{\frac{(N-1)(1+\delta)}{2} + 1 - N} \leq C(1 + |x|)^{1+\delta}. \end{aligned} \quad (3.7)$$

Combining all these bounds, we have shown that

$$\int_{\mathcal{C}} |G(x, y)| dy \leq C(1 + |x|)^{\frac{3+\delta}{2}},$$

where $\delta \in (0, 1)$ is arbitrary. This proves that $V_{\mathcal{C}}$ is well-defined and locally bounded. Also, choosing $\delta = 1/2$, we obtain that

$$\frac{|V_{\mathcal{C}}(x)|}{(1 + |x|)^{7/4}} \leq C_{\hat{r}, N} \quad \text{for all } x \in \mathbb{R}^N \quad (3.8)$$

where the constant $C_{\hat{r}, N}$ depends only on the dimension N and the radius \hat{r} defined in (2.2) for $\delta = 1/2$.

To prove the $W^{2,p}$ -regularity of $V_{\mathcal{C}}$ we note that, for $\varrho > 2 \max\{\hat{r}, |x|\}$,

$$|V_{\mathcal{C}}(x) - V_{\mathcal{C} \cap B_\varrho}(x)| \leq C \int_{T_\delta \setminus B_\varrho} \frac{|x|^2}{|y|^N} dy \leq C|x|^2 \int_\varrho^\infty t^{\frac{(N-1)(1+\delta)}{2} - N} dt \leq C|x|^2 \varrho^{\delta-1}.$$

This implies that $V_{\mathcal{C}}$ is the locally uniform limit of the sequence of the continuous functions $V_{\mathcal{C} \cap B_\varrho}$ as $\varrho \rightarrow \infty$. Also, since

$$\alpha_N \Delta_x \left(\frac{1}{|x-y|^{N-2}} - \frac{1}{|y|^{N-2}} - (N-2) \frac{x \cdot y}{|y|^N} \right) = -\delta_x \quad \text{in the sense of distributions,}$$

one easily deduces that $\Delta V_{\mathcal{C} \cap B_\varrho} = -\chi_{\mathcal{C} \cap B_\varrho} \in L^\infty(\mathbb{R}^N)$ for each $\varrho > 0$. Thus, by elliptic regularity, the functions $V_{\mathcal{C} \cap B_\varrho}$ are locally uniformly bounded in $W^{2,p}$ for each $p < \infty$. In

particular because of the compact embedding $W^{2,p}(B_\varrho) \hookrightarrow C^{1,\alpha}(B_\varrho)$ for $p > N$ and $\alpha = 1 - \frac{N}{p}$, we deduce that

$$V_{\mathcal{C} \cap B_\varrho} \rightarrow V_{\mathcal{C}} \quad \text{and} \quad \nabla V_{\mathcal{C} \cap B_\varrho} \rightarrow \nabla V_{\mathcal{C}} \quad \text{locally uniformly in } \mathbb{R}^N, \text{ as } \varrho \rightarrow \infty.$$

Since $G(0, \cdot) \equiv 0$ and $\nabla_x G(0, \cdot) \equiv 0$, we obtain for each $\varrho > 0$ that

$$V_{\mathcal{C} \cap B_\varrho}(0) = \int_{\mathcal{C} \cap B_\varrho} G(0, y) \, dy = 0, \quad \nabla V_{\mathcal{C} \cap B_\varrho}(0) = \int_{\mathcal{C} \cap B_\varrho} \nabla_x G(0, y) \, dy = 0,$$

so we conclude that $V_{\mathcal{C}}(0) = \nabla V_{\mathcal{C}}(0) = 0$. □

As a consequence of the previous lemma, we can now show the following important result.

Proposition 3.4 (Newtonian potential expansion). *Let $N \geq 3$, let u be an x_N -monotone solution in the sense of Definition 2.8, and let p be the blow-down limit in Definition 2.8(iv). Then the expansion*

$$u = p + V_{\mathcal{C}}$$

holds.

Proof. Recall that, thanks to Lemma 3.3(iii), $V_{\mathcal{C}}$ is a strong $W_{\text{loc}}^{2,p}(\mathbb{R}^N)$ solution of $\Delta V_{\mathcal{C}} = -\chi_{\mathcal{C}}$. Moreover, if we set $v := u - p$, then $v \in C_{\text{loc}}^{1,1}(\mathbb{R}^N)$ (see Lemma 2.4(i)) and it solves the same equation as $V_{\mathcal{C}}$, i.e., $\Delta v = -\chi_{\mathcal{C}}$. Hence $v - V_{\mathcal{C}}$ is harmonic in \mathbb{R}^N , and it follows from Definition 2.8(iv) and Lemma 3.3(ii) that $v - V_{\mathcal{C}}$ has subquadratic growth. This allows us to apply Liouville's theorem to obtain that

$$v - V_{\mathcal{C}} = \ell + c,$$

where ℓ is a linear function and c is a constant. Thus

$$u = p + \ell + c + V_{\mathcal{C}} \quad \text{in } \mathbb{R}^N.$$

Now, since $0 \in \partial\mathcal{C}$, it follows from Lemma 3.3(iii) that

$$\begin{aligned} 0 &= u(0) = p(0) + \ell(0) + c + V_{\mathcal{C}}(0) = c, \\ 0 &= \nabla u(0) = \nabla p(0) + \nabla \ell(0) + \nabla V_{\mathcal{C}}(0) = \nabla \ell. \end{aligned}$$

This proves that both ℓ and c vanish, concluding the proof. □

As we shall see in the next section, this potential expansion allows us to obtain a very precise control on the asymptotic behavior of the coincidence set \mathcal{C} .

4. IMPROVED ESTIMATE ON THE ASYMPTOTIC BEHAVIOR OF THE COINCIDENCE SET \mathcal{C}

The goal of this section is to prove that \mathcal{C} is contained in some paraboloid. While for $N \geq 4$ there is a very simple argument to prove this result, the proof for $N = 3$ is amongst the most delicate of this paper (see in particular the proof of Lemma 4.3 below).

Proposition 4.1 (\mathcal{C} is contained in a paraboloid). *Let $N \geq 3$, and let u be an x_N -monotone solution in the sense of Definition 2.8. Then there are constants $a_0, \gamma_0 \in (0, \infty)$ such that:*

- (i) $\mathcal{C} \cap \{x_N > a_0\} \subset \{|x'|^2 < \gamma_0 x_N\}$;
- (ii) $\mathcal{C} \cap \{x_N \leq a_0\}$ is bounded.

Proof. Thanks to Lemma 2.10 it follows that $\mathcal{C} \cap \{x_N \leq a\}$ is bounded for each $a > 0$, so (ii) holds for each $a_0 > 0$. In particular, it suffices to prove (i) for a_0 sufficiently large.

We first prove the result in the case $N \geq 4$ (since the proof is very simple), and then focus on the delicate case $N = 3$.

• **The case $N \geq 4$.** Arguing as in the proof of Lemma 3.3, given $\delta \in (0, 1)$, for $|x| > \hat{r}$ we have

$$\begin{aligned} \frac{1}{\alpha_N} V_{\mathcal{C}}(x) &= \int_{\mathcal{C}} G(x, y) \, dy \geq \int_{\mathcal{C} \cap B_{2\hat{r}}} \left(\frac{1}{|x-y|^{N-2}} - \frac{1}{|y|^{N-2}} - (N-2) \frac{|x|}{|y|^{N-1}} \right) \, dy \\ &\quad + \int_{\mathcal{C} \cap (B_{2|x|} \setminus B_{2\hat{r}})} \left(\frac{1}{|x-y|^{N-2}} - \frac{1}{|y|^{N-2}} - (N-2) \frac{|x|}{|y|^{N-1}} \right) \, dy \\ &\quad - C \int_{T_{\delta} \setminus (B_{2|x|} \cup B_{2\hat{r}})} \frac{|x|^2}{|y|^N} \, dy =: I_1 + I_2 + I_3 \end{aligned}$$

(cp. (3.3)). Again as in the proof of Lemma 3.3, we have that $|I_1| \leq C(1 + |x|)$ and

$$|I_3| \leq C|x|^2(1 + |x|)^{\frac{(N-1)(1+\delta)}{2} + 1 - N} \leq C(1 + |x|)^{\frac{1+3\delta}{2}} \quad \text{for } N \geq 4$$

(cp. (3.4) and (3.7)). For I_2 , we observe that the first term is non-negative and we estimate the remaining two as in (3.5) and (3.6), so to get

$$\begin{aligned} I_2 &\geq - \int_{T_{\delta} \cap (B_{2|x|} \setminus B_{2\hat{r}})} \left(\frac{1}{|y|^{N-2}} + (N-2) \frac{|x|}{|y|^{N-1}} \right) \, dy \\ &\geq -C(1 + |x|)^{\frac{(N-1)(1+\delta)}{2} + 3 - N} \geq -C(1 + |x|)^{\frac{1+3\delta}{2}} \quad \text{for } N \geq 4. \end{aligned}$$

Choosing $\delta \leq 1/3$ proves that $V_{\mathcal{C}}(x) \geq -C(1 + |x|)$ for all $x \in \mathbb{R}^N$.

Now, applying Proposition 3.4 and combining this bound with (2.1), whenever $x \in \{u = 0\}$ we obtain

$$0 = u(x) = p(x) + V_{\mathcal{C}}(x) \geq c_P|x'|^2 - C(1 + |x|) \geq c_P|x'|^2 - C - C|x'| - C|x_N|.$$

From this estimate, we easily deduce that

$$|x'|^2 \leq C(1 + |x_N|) = C(1 + x_N) \quad \text{for all } x \in \{u = 0\}$$

(recall that $\{u = 0\} \subset \{x_N \geq 0\}$), so (i) follows.

• **The case $N = 3$.** This case follows from Lemmas 4.2 and 4.3 below. \square

The rest of the section is devoted to the proof of Lemmas 4.2 and 4.3.

Lemma 4.2 (Sections of \mathcal{C} are controlled by their measure). *Let $N = 3$, and let u be an x_N -monotone solution in the sense of Definition 2.8. Define $\mathcal{C}_t := \{y' \in \mathbb{R}^2 : (y', t) \in \mathcal{C}\}$ and $H(t) := \mathcal{H}^2(\mathcal{C}_t)$ for all $t \geq 0$. Then:*

- either $\{\mathcal{C}_t\}_{t \geq 0}$ is bounded, i.e., $\sup_{t \geq 0} \text{diam}(\mathcal{C}_t) < \infty$;
- or there exist $a_0 > 1$ and $C_0 < \infty$ such that, for all $x_3 \geq a_0$,

$$\mathcal{C}_{x_3} \subset \{|x'|^2 < C_0 H(x_3)\}.$$

Proof. We may assume that

$$\sup_{t \geq 0} \text{diam}(\mathcal{C}_t) = \infty. \tag{4.1}$$

Suppose, towards a contradiction, that the statement of the lemma is not true. Then, there exists a sequence $(x^n)_{n \in \mathbb{N}} \subset \mathbb{R}^2 \times (0, \infty)$, with $x^n = ((x^n)', x_3^n)$, such that $x_3^n \rightarrow \infty$ as $n \rightarrow \infty$ and, for all $n \in \mathbb{N}$,

$$x^n \in \mathcal{C} \quad \text{and} \quad |(x^n)'|^2 > nH(x_3^n). \quad (4.2)$$

Define $d_n := \text{diam}(\mathcal{C}_{x_3^n})$. From Lemma 2.10 we know that, given $\delta \in (0, 1)$,

$$d_n \leq (x_3^n)^{\frac{1+\delta}{2}} \quad \text{for all } n \text{ sufficiently large.} \quad (4.3)$$

On the other hand, (4.1) together with the monotonicity of $t \mapsto \text{diam}(\mathcal{C}_t)$ (recall that, by Definition 2.8, u is decreasing in the e^3 -direction) imply that

$$d_n \rightarrow \infty \quad \text{as } n \rightarrow \infty.$$

Let us define for each $n \in \mathbb{N}$ the rescaling

$$u_n(x) := \frac{u((0, x_3^n) + d_n x)}{d_n^2}.$$

Note that, as a consequence of (4.2), the convex sets $\{u_n = 0\} \cap \{x_3 = 0\} \subset \mathbb{R}^2$ have diameter 1, contain the origin, and their \mathcal{H}^2 -measure goes to zero. Hence, thanks to Lemma 2.4(ii), the fact that $t \mapsto \mathcal{C}(t)$ is increasing (again by the fact that u is decreasing in the e^3 -direction), and the convergence of coincidence sets (see [17, Proposition 3.17 (iv) and Proposition 3.17 (v)]), passing if necessary to a subsequence we obtain that

$$u_n \rightarrow u_0 \quad \text{in } C_{\text{loc}}^{1,\alpha}(\mathbb{R}^3) \text{ as } n \rightarrow \infty,$$

with

$$\text{diam}(\{u_0 = 0\} \cap \{x_3 = 0\}) \geq 1, \quad |\{u_0 = 0\} \cap \{-1 < x_3 < 0\}| = 0, \quad (4.4)$$

where u_0 is a global solution to the obstacle problem. Also, since $0 \in \mathcal{C}$ (cf. Definition 2.8(ii)) and $\delta \in (0, 1)$, it follows from (4.3) that

$$\frac{\text{dist}(0, (0, x_3^n))}{d_n} \geq \frac{x_3^n}{(x_3^n)^{\frac{1+\delta}{2}}} \rightarrow \infty \quad \text{as } n \rightarrow \infty.$$

Thus, by the convexity of \mathcal{C} (cf. Remark 2.2), we deduce that $\{te^3 : t \leq 0\} \subset \{u_0 = 0\}$. On the other hand, the fact that $\{te^3 : t \geq 0\} \subset \mathcal{C}$ (see Remark 2.9) implies that $\{te^3 : t \geq 0\} \subset \{u_0 = 0\}$. Hence

$$\{te^3 : t \in \mathbb{R}\} \subset \{u_0 = 0\},$$

and therefore it follows from Lemma 2.7 that u_0 is invariant the e^3 -direction, i.e.

$$u_0(x) = u_0(x', 0) \quad \text{for all } x \in \mathbb{R}^3.$$

Combining this information with (4.4), we deduce that the coincidence set of u_0 has measure zero, hence Remark 2.5 implies that u_0 coincides with a quadratic polynomial $q = q(x')$. On the other hand, [7, Lemma B.2] implies that the blow-down limit of u_0 is p (being the blow-down limit of u), and therefore the only possibility is that $u_0 = q = p$. By the nondegeneracy of p in \mathbb{R}^2 (see Definition 2.8(iv)), this implies that $\{u_0 = 0\} \cap \{x_3 = 0\}$ coincides with the origin, a contradiction to the fact that this set has diameter at least 1 (see (4.4)). This contradiction proves the lemma. \square

Lemma 4.3 (The measure of sections grows at most linearly). *Let $N = 3$, and let u be an x_N -monotone solution in the sense of Definition 2.8. Then there exists a constant C such that, for all $t \geq 0$,*

$$\mathcal{H}^2(\mathcal{C} \cap \{x_3 = t\}) \leq C(1 + t).$$

Proof. We split the proof into 8 steps.

Step 1. *Preliminary observations about sections of \mathcal{C} .*

Recall the notation $\mathcal{C}_t := \{y' \in \mathbb{R}^2 : (y', t) \in \mathcal{C}\}$ and $H(t) := \mathcal{H}^2(\mathcal{C}_t)$.

First, we claim that \sqrt{H} is a concave function, i.e., for all $\lambda \in [0, 1]$ and $t_1, t_2 \geq 0$,

$$\lambda\sqrt{H(t_1)} + (1 - \lambda)\sqrt{H(t_2)} \leq \sqrt{H(\lambda t_1 + (1 - \lambda)t_2)}.$$

Indeed, by the Brunn-Minkowski inequality in \mathbb{R}^2 ,

$$\left(\lambda\sqrt{\mathcal{H}^2(\mathcal{C}_{t_1})} + (1 - \lambda)\sqrt{\mathcal{H}^2(\mathcal{C}_{t_2})}\right)^2 \leq \mathcal{H}^2(\lambda\mathcal{C}_{t_1} + (1 - \lambda)\mathcal{C}_{t_2}).$$

Hence, since $\lambda\mathcal{C}_{t_1} + (1 - \lambda)\mathcal{C}_{t_2} \subset \mathcal{C}_{\lambda t_1 + (1 - \lambda)t_2}$ (by the convexity of \mathcal{C}), the claim follows.

Now, the concavity of \sqrt{H} together with the smoothness³ of $\partial\mathcal{C}$ implies that

$$(\sqrt{H})'(t)(t - s) \leq \sqrt{H(t)} - \sqrt{H(s)} \leq (\sqrt{H})'(s)(t - s) \quad \text{for all } 0 \leq s \leq t. \quad (4.5)$$

In particular, since $H(0) \geq 0$,

$$(\sqrt{H})'(t) \leq \frac{\sqrt{H(t)}}{t} \quad \text{for all } t > 0. \quad (4.6)$$

Furthermore, by the monotonicity of u in the x_3 -direction,

$$0 \leq H(s) \leq H(t) \quad \text{for all } 0 \leq s \leq t. \quad (4.7)$$

Finally, from Lemma 2.10 we infer that for every $\delta \in (0, 1)$ there is $a(\delta) > 0$ such that

$$H(t) \leq t^{1+\delta} \quad \text{for all } t \geq a(\delta). \quad (4.8)$$

Step 2. *The generalized Newtonian potential expansion.*

Let $V_{\mathcal{C}}$ be the generalized Newtonian potential of the coincidence set \mathcal{C} , cf. Definition 3.1.

Thanks to Lemma 4.2 and (4.8) it follows that, for $t > 0$ sufficiently large,

$$0 \leq \frac{1}{|te^3 - y|} - \frac{1}{|y|} - \frac{ty_3}{|y|^3} \quad \text{for all } y = (y', y_3) \in \mathcal{R}_t := \left\{t - \sqrt{H(t)} < y_3 < t + \sqrt{H(t)}\right\} \cap \mathcal{C}.$$

Combining this with Proposition 3.4, Remark 2.9, and Definition 2.8(iv), we find that, for t sufficiently large,

$$0 = u(te^3) = V_{\mathcal{C}}(te^3) \geq \tilde{V}_{\mathcal{C}}(te^3) := \alpha_3 \int_{\mathcal{C} \setminus \mathcal{R}_t} \left(\frac{1}{|te^3 - y|} - \frac{1}{|y|} - \frac{ty_3}{|y|^3} \right) dy. \quad (4.9)$$

Step 3. *A one-dimensionalized version of $\tilde{V}_{\mathcal{C}}$.*

The objective of this step is to replace the potential integral defining $\tilde{V}_{\mathcal{C}}$ by a one-dimensional

³Since \mathcal{C} is a convex set with non-empty interior, it follows from the regularity theory of the free boundary for the obstacle problem that $\partial\mathcal{C}$ is smooth (see for instance [3]). However, if one does not want to rely on this result, it suffices to replace $(\sqrt{H})'$ with the right or left limit of the derivative of \sqrt{H} , which always exist by the concavity of \sqrt{H} .

integral, up to a well-controlled error. To be more precise, we claim that there exists a constant C such that, for all $t > 0$ sufficiently large,

$$\left| \int_{(\mathcal{C} \cap \{y_3 \geq a\}) \setminus \mathcal{R}_t} \left(\frac{1}{|te^3 - y|} - \frac{1}{|y|} - \frac{ty_3}{|y|^3} \right) dy - W(t) \right| \leq C(H(t) + t), \quad (4.10)$$

where

$$W(t) := \int_a^{t - \sqrt{H(t)}} \left(\frac{1}{t-s} - \frac{1}{s} - \frac{t}{s^2} \right) H(s) ds + \int_{t + \sqrt{H(t)}}^{\infty} \left(\frac{1}{s-t} - \frac{1}{s} - \frac{t}{s^2} \right) H(s) ds,$$

with $a := \max\{a_0, a(\delta)\} > 0$, where a_0 is as in Lemma 4.2, $a(\delta)$ is as in (4.8), and $\delta > 0$ is sufficiently small ($\delta < 1/2$ suffices here).

For the remainder of this step, fix a point $y = (y', s) \in \mathcal{C} \setminus \mathcal{R}_t$. Note that, by the definition of \mathcal{R}_t , we have $|t - s| \geq \sqrt{H(t)}$. Combining Lemma 4.2, (4.5), (4.6), and (4.7),

$$|y'|^2 \leq CH(s) \leq CH(t) \quad \text{for all } s \in (a, t - \sqrt{H(t)}) \text{ and} \quad (4.11)$$

$$\begin{aligned} |y'|^2 &\leq CH(s) \leq C \left(\sqrt{H(t)} + \frac{\sqrt{H(t)}}{t} (s-t) \right)^2 \\ &\leq C \left(H(t) + \frac{H(t)}{t^2} (s-t)^2 \right) \quad \text{for all } s \geq t + \sqrt{H(t)}. \end{aligned} \quad (4.12)$$

Let us now note that, by the mean value theorem, there is $\xi_{s,t} \in (0, |y'|^2)$ such that

$$\begin{aligned} &\left| \frac{1}{\sqrt{(s-t)^2 + |y'|^2}} - \frac{1}{\sqrt{s^2 + |y'|^2}} - \frac{ts}{(s^2 + |y'|^2)^{\frac{3}{2}}} - \left(\frac{1}{|s-t|} - \frac{1}{s} - \frac{t}{s^2} \right) \right| \\ &\leq \frac{1}{2} \left| \frac{1}{((s-t)^2 + \xi_{s,t})^{\frac{3}{2}}} - \frac{1}{(s^2 + \xi_{s,t})^{\frac{3}{2}}} - 3 \frac{ts}{(s^2 + \xi_{s,t})^{\frac{5}{2}}} \right| |y'|^2. \end{aligned} \quad (4.13)$$

For $s \in (a, t - \sqrt{H(t)})$, we can estimate the right-hand side above as

$$\begin{aligned} &\frac{1}{2} \left| \frac{1}{((s-t)^2 + \xi_{s,t})^{\frac{3}{2}}} - \frac{1}{(s^2 + \xi_{s,t})^{\frac{3}{2}}} - 3 \frac{ts}{(s^2 + \xi_{s,t})^{\frac{5}{2}}} \right| |y'|^2 \\ &\leq \frac{1}{2} \left| \frac{1}{(t-s)^3} + \frac{1}{s^3} + \frac{3ts}{s^5} \right| |y'|^2 \leq 2 \left(\frac{1}{(t-s)^3} + \frac{t}{s^4} \right) |y'|^2, \end{aligned}$$

so (4.13) implies that

$$\begin{aligned} &\left| \frac{1}{\sqrt{(s-t)^2 + |y'|^2}} - \frac{1}{\sqrt{s^2 + |y'|^2}} - \frac{ts}{(s^2 + |y'|^2)^{\frac{3}{2}}} - \left(\frac{1}{|s-t|} - \frac{1}{s} - \frac{t}{s^2} \right) \right| \\ &\leq 2 \left(\frac{1}{(t-s)^3} + \frac{t}{s^4} \right) |y'|^2 \quad \text{for all } s \in (a, t - \sqrt{H(t)}). \end{aligned} \quad (4.14)$$

Now, combining first (4.14), (4.11), (4.8), and (4.7), we see that

$$\begin{aligned}
& \left| \int_{\mathcal{C} \cap \{a \leq y_3 \leq t - \sqrt{H(t)}\}} \left(\frac{1}{|te^3 - y|} - \frac{1}{|y|} - \frac{ty_3}{|y|^3} \right) dy - \int_a^{t - \sqrt{H(t)}} \left(\frac{1}{t-s} - \frac{1}{s} - \frac{t}{s^2} \right) H(s) ds \right| \\
& \leq C \int_a^{t - \sqrt{H(t)}} \left(\frac{1}{(t-s)^3} + \frac{t}{s^4} \right) H(s)^2 ds \\
& \leq CH(t)^2 \int_a^{t - \sqrt{H(t)}} \frac{1}{(t-s)^3} ds + Ct \int_a^{t - \sqrt{H(t)}} \frac{s^{2+2\delta}}{s^4} ds \leq C(H(t) + t). \tag{4.15}
\end{aligned}$$

Concerning the case $s \geq t + \sqrt{H(t)}$, we can either apply the Taylor formula $f(1) = f(0) + f'(0) + \int_0^1 (1-\tau)f''(\tau) d\tau$ with $f(\tau) := ((s-t\tau)^2 + \xi_{s,t})^{-\frac{3}{2}}$ to get

$$\left| \frac{1}{((s-t)^2 + \xi_{s,t})^{\frac{3}{2}}} - \frac{1}{(s^2 + \xi_{s,t})^{\frac{3}{2}}} - 3 \frac{ts}{(s^2 + \xi_{s,t})^{\frac{5}{2}}} \right| \leq C \frac{t^2}{((s-t)^2 + \xi_{s,t})^{\frac{5}{2}}} \leq C \frac{t^2}{(s-t)^5},$$

or, by a direct estimate, we can bound

$$\left| \frac{1}{((s-t)^2 + \xi_{s,t})^{\frac{3}{2}}} - \frac{1}{(s^2 + \xi_{s,t})^{\frac{3}{2}}} - 3 \frac{ts}{(s^2 + \xi_{s,t})^{\frac{5}{2}}} \right| \leq \left| \frac{1}{(s-t)^3} + \frac{1}{s^3} + \frac{3t}{s^4} \right| \leq C \frac{1}{(s-t)^3}.$$

Combining the last two inequalities with (4.13), this implies

$$\begin{aligned}
& \left| \frac{1}{\sqrt{(s-t)^2 + |y'|^2}} - \frac{1}{\sqrt{s^2 + |y'|^2}} - \frac{ts}{(s^2 + |y'|^2)^{\frac{3}{2}}} - \left(\frac{1}{|s-t|} - \frac{1}{s} - \frac{t}{s^2} \right) \right| \\
& \leq C \min \left\{ \frac{1}{(s-t)^3}, \frac{t^2}{(s-t)^5} \right\} |y'|^2 \quad \text{for all } s \geq t + \sqrt{H(t)}. \tag{4.16}
\end{aligned}$$

Using (4.7), (4.12), (4.16), and (4.8), we obtain

$$\begin{aligned}
& \left| \int_{\mathcal{C} \cap \{y_3 \geq t + \sqrt{H(t)}\}} \left(\frac{1}{|te^3 - y|} - \frac{1}{|y|} - \frac{ty_3}{|y|^3} \right) dy - \int_{t + \sqrt{H(t)}}^{\infty} \left(\frac{1}{s-t} - \frac{1}{s} - \frac{t}{s^2} \right) H(s) ds \right| \\
& \leq C \int_{t + \sqrt{H(t)}}^{\infty} \min \left\{ \frac{1}{(s-t)^3}, \frac{t^2}{(s-t)^5} \right\} H(s)^2 ds \\
& \leq C \int_{t + \sqrt{H(t)}}^{2t} \frac{1}{(s-t)^3} H(s)^2 ds + C \int_{2t}^{\infty} \frac{t^2}{(s-t)^5} H(s)^2 ds \\
& \leq CH(2t)^2 \int_{t + \sqrt{H(t)}}^{2t} \frac{1}{(s-t)^3} ds + CH(t) \int_{2t}^{\infty} \left(\frac{t^2}{(s-t)^5} + \frac{1}{(s-t)^3} \right) H(s) ds
\end{aligned}$$

$$\leq C \left(H(t) + \frac{H(t)}{t^2} (2t-t)^2 \right)^2 \int_{t+\sqrt{H(t)}}^{2t} \frac{1}{(s-t)^3} ds + CH(t) \int_{2t}^{\infty} \left(\frac{s^2}{\left(\frac{s}{2}\right)^5} + \frac{1}{\left(\frac{s}{2}\right)^3} \right) s^{1+\delta} ds \leq CH(t).$$

This bound, together with (4.15), concludes the proof of (4.10).

Step 4. *Estimating W from below.*

To simplify notation we set, for $t \geq a$,

$$W_1(t) := \int_a^{t-\sqrt{H(t)}} \left(\frac{1}{t-s} - \frac{1}{s} - \frac{t}{s^2} \right) H(s) ds, \quad W_2(t) := \int_{t+\sqrt{H(t)}}^{\infty} \left(\frac{1}{s-t} - \frac{1}{s} - \frac{t}{s^2} \right) H(s) ds,$$

so that $W(t) = W_1(t) + W_2(t)$.

To estimate W_1 from below, we split the integral so that the integrand in each part has a sign. More precisely, since

$$\frac{1}{t-s} - \frac{1}{s} - \frac{t}{s^2} \leq 0 \quad \text{for } s \in \left[a, \frac{t}{\sqrt{2}} \right] \quad \text{and} \quad \frac{1}{t-s} - \frac{1}{s} - \frac{t}{s^2} \geq 0 \quad \text{for } s \in \left[\frac{t}{\sqrt{2}}, t - \sqrt{H(t)} \right],$$

we set $W_1 = W_{1,1} + W_{1,2}$ with

$$W_{1,1}(t) := \int_a^{t/\sqrt{2}} \left(\frac{1}{t-s} - \frac{1}{s} - \frac{t}{s^2} \right) H(s) ds, \quad W_{1,2}(t) := \int_{t/\sqrt{2}}^{t-\sqrt{H(t)}} \left(\frac{1}{t-s} - \frac{1}{s} - \frac{t}{s^2} \right) H(s) ds.$$

We estimate $W_{1,1}$ by neglecting the first term, so to get

$$W_{1,1}(t) \geq - \int_a^{t/\sqrt{2}} \left(\frac{1}{s} + \frac{t}{s^2} \right) H(s) ds \geq - \int_a^t \frac{H(s)}{s} ds - t \int_a^t \frac{H(s)}{s^2} ds.$$

To estimate $W_{1,2}$, using (4.5), (4.7), and (4.6), we obtain

$$\begin{aligned} W_{1,2}(t) &= H(t) \int_{t/\sqrt{2}}^{t-\sqrt{H(t)}} \frac{1}{t-s} ds + \int_{t/\sqrt{2}}^{t-\sqrt{H(t)}} \frac{H(s) - H(t)}{t-s} ds - \int_{t/\sqrt{2}}^{t-\sqrt{H(t)}} \left(\frac{1}{s} + \frac{t}{s^2} \right) H(s) ds \\ &\geq H(t) \left(\log(t) + \log \left(1 - \frac{1}{\sqrt{2}} \right) \right) - \frac{1}{2} H(t) \log(H(t)) \\ &\quad - \int_{t/\sqrt{2}}^{t-\sqrt{H(t)}} \left(\sqrt{H(t)} + \sqrt{H(s)} \right) \frac{\sqrt{H(t)} - \sqrt{H(s)}}{t-s} ds - CH(t) \int_{t/\sqrt{2}}^{t-\sqrt{H(t)}} \left(\frac{1}{s} + \frac{t}{s^2} \right) ds \end{aligned}$$

$$\begin{aligned}
&\geq H(t) \log(t) - \frac{1}{2}H(t) \log(H(t)) - 2\sqrt{H(t)} \int_{t/\sqrt{2}}^{t-\sqrt{H(t)}} \frac{\sqrt{H(t)} - \sqrt{H(s)}}{t-s} ds - CH(t) \\
&\geq H(t) \log(t) - \frac{1}{2}H(t) \log(H(t)) - 2\sqrt{H(t)} \int_{t/\sqrt{2}}^{t-\sqrt{H(t)}} (\sqrt{H})'(s) ds - CH(t) \\
&\geq H(t) \log(t) - \frac{1}{2}H(t) \log(H(t)) - CH(t).
\end{aligned}$$

It remains to estimate $W_2(t)$. Since the integrand in W_2 is non-negative, using (4.7) we obtain that

$$W_2(t) \geq H(t) \int_{t+\sqrt{H(t)}}^{\infty} \left(\frac{1}{s-t} - \frac{1}{s} - \frac{t}{s^2} \right) ds \geq H(t) \log(t) - \frac{1}{2}H(t) \log(H(t)) - H(t).$$

Combining all estimates, we find the lower bound

$$W(t) \geq 2H(t) \log(t) - H(t) \log(H(t)) - CH(t) - \int_a^t \frac{H(s)}{s} ds - t \int_a^t \frac{H(s)}{s^2} ds. \quad (4.17)$$

Step 5. *An integral inequality.*

Combining (4.9), (4.10), and (4.17), we deduce the existence of a constant C such that, for $t > 0$ sufficiently large,

$$C(H(t) + t) \geq 2H(t) \log(t) - H(t) \log(H(t)) - \int_a^t \frac{H(s)}{s} ds - t \int_a^t \frac{H(s)}{s^2} ds$$

which implies, in particular, that

$$\begin{aligned}
C(H(t) + t) &\geq 2(H(t) + t) \log(t) - (H(t) + t) \log(H(t) + t) \\
&\quad - \int_a^t \frac{H(s) + s}{s} ds - t \int_a^t \frac{H(s) + s}{s^2} ds.
\end{aligned}$$

Hence, setting

$$\psi(t) := \frac{H(t) + t}{t},$$

for all t sufficiently large we get

$$\begin{aligned}
C\psi(t) &\geq \psi(t) \log(t) - \psi(t) \log(\psi(t)) - \frac{1}{t} \int_a^t \psi(s) ds - \int_a^t \frac{\psi(s)}{s} ds \\
&\geq -\psi(t) \log(\psi(t)) - \frac{1}{t} \int_a^t \psi(s) ds + \int_a^t \frac{\psi(t) - \psi(s)}{s} ds. \quad (4.18)
\end{aligned}$$

Our goal in the following is to show that ψ is globally bounded. To this end, let us replace ψ by the monotone function

$$\Psi(t) := \sup_{s \in [a, t]} \psi(s).$$

In order to find an integral inequality for Ψ , given $t \geq a$ let $\tau = \tau(t) \in [a, t]$ be such that $\Psi(t) = \psi(\tau)$. Then, by the definition of Ψ ,

$$\Psi(s) = \Psi(t) \quad \text{for all } s \in [\tau, t]. \quad (4.19)$$

Noticing that $\frac{1}{t} \int_a^t \Psi(s) ds \geq \frac{1}{\tau} \int_a^\tau \Psi(s) ds$ (since Ψ is increasing) and that $\Psi(t) \geq \psi(t)$ by construction, we deduce that

$$C\Psi(t) + \Psi(t) \log(\Psi(t)) + \frac{1}{t} \int_a^t \Psi(s) ds \geq C\psi(\tau) + \psi(\tau) \log(\psi(\tau)) + \frac{1}{\tau} \int_a^\tau \psi(s) ds.$$

Thus, since $\Psi(s) \geq \psi(s)$ for all $s \geq a$, it follows from (4.18) and (4.19) that

$$\begin{aligned} C\Psi(t) + \Psi(t) \log(\Psi(t)) + \frac{1}{t} \int_a^t \Psi(s) ds &\geq \int_a^\tau \frac{\psi(\tau) - \psi(s)}{s} ds = \int_a^\tau \frac{\Psi(t) - \psi(s)}{s} ds \\ &\geq \int_a^\tau \frac{\Psi(t) - \Psi(s)}{s} ds = \int_a^t \frac{\Psi(t) - \Psi(s)}{s} ds. \end{aligned}$$

Since $\frac{1}{t} \int_a^t \Psi(s) ds \leq \Psi(t)$ (by the monotonicity of Ψ), we can simplify the relation above to conclude that

$$C\Psi(t) + \Psi(t) \log(\Psi(t)) \geq \Psi(t) \log(t) - \int_a^t \frac{\Psi(s)}{s} ds. \quad (4.20)$$

Step 6. *Switching to a differential inequality and comparison.*

Define $F(t) := \int_a^t \frac{\Psi(s)}{s} ds$. Then $F'(t) = \frac{\Psi(t)}{t} > 0$ and (4.20) becomes

$$-CF'(t) - F'(t) \log(F'(t)) \leq \frac{F(t)}{t}. \quad (4.21)$$

Since $H(t) \leq t^{1+\delta}$ (see (4.8)) it follows that $\psi(t) \leq 2t^\delta$. Therefore $\Psi(t) \leq 2t^\delta$, from which it follows that

$$F'(t) \leq 2t^{\delta-1} \quad \text{and} \quad F(t) \leq \frac{2}{\delta} t^\delta. \quad (4.22)$$

In particular, this yields

$$F'(t) \rightarrow 0 \quad \text{and} \quad 0 > F'(t) \log(F'(t)) \rightarrow 0 \quad \text{as } t \rightarrow \infty.$$

Note now that, for $\tau_0 > 0$ small enough, the function

$$h : (0, \tau_0) \rightarrow (0, \infty), \quad h(\tau) := -C\tau - \tau \log(\tau) = -C\tau + \tau |\log \tau|$$

is strictly increasing, invertible, and has a locally Lipschitz-continuous inverse. Also, for $t_0 > 0$ sufficiently large, $F'(t) \leq \tau_0$ and

$$h(F'(t)) \leq \frac{F(t)}{t} \quad \text{for all } t \geq t_0.$$

This implies that the above ODE enjoys the comparison principle, i.e., if $G : (t_0, \infty) \rightarrow \mathbb{R}$, with $0 < G'(t) < \tau_0$, satisfies

$$h(G'(t)) \geq \frac{G(t)}{t} \quad \text{for all } t \geq t_0 \quad \text{and} \quad F(t_0) \leq G(t_0)$$

then $F(t) \leq G(t)$ for all $t \geq t_0$.

Step 7. *Construction of a comparison solution.*

Let $A > 1, B > 0$, and define $G_{A,B} : (0, \infty) \rightarrow \mathbb{R}$ as $G_{A,B}(t) := A \log(t) - B$. Then $G'_{A,B}(t) = \frac{A}{t} \in (0, \tau_0)$ provided $t > A\tau_0$, and with the choice $B := A(C + \log(A))$ it holds

$$\begin{aligned} h(G'_{A,B}(t)) &= -C \frac{A}{t} - \frac{A}{t} \log\left(\frac{A}{t}\right) = -\frac{A}{t}(C + \log(A)) + \frac{A \log(t)}{t} \\ &= \frac{A \log(t) - B}{t} = \frac{G_{A,B}(t)}{t} \quad \text{for all } t > 0. \end{aligned}$$

Fix now $\delta \in (0, \frac{1}{2})$ and define $A := t_0^{2\delta}$. If t_0 is chosen sufficiently large, then $G'_{A,B}(t) \in (0, \tau_0)$ for $t \geq t_0$. Also, thanks to (4.22) we get

$$G_{A,B}(t_0) = t_0^{2\delta} \log(t_0) - 2\delta t_0^{2\delta} \log(t_0) - C t_0^{2\delta} = ((1 - 2\delta) \log(t_0) - C) t_0^{2\delta} \geq F(t_0).$$

Step 8. *Conclusion.*

By the comparison principle mentioned in Step 6, choosing A and B as in Step 7 we deduce that

$$F(t) \leq G_{A,B}(t) \leq A \log(t) \quad \text{for all } t \geq t_0. \quad (4.23)$$

Also, for all $0 < x, y \ll 1$,

$$y \geq h(x) = -Cx + x |\log(x)| \implies x \leq \frac{2y}{|\log(x)|} \leq \frac{2y}{|\log(y)|} = \frac{2y}{-\log(y)}.$$

Hence, recalling (4.21), (4.23), and (4.22), for $t \geq t_0$ we obtain

$$F'(t) \leq 2 \frac{\frac{F(t)}{t}}{-\log\left(\frac{F(t)}{t}\right)} \leq 2 \frac{\frac{A \log(t)}{t}}{-\log\left(\frac{A \log(t)}{t}\right)} = \frac{2A \log(t)}{t(\log(t) - \log(\log(t)) - \log(A))} \leq \frac{4A}{t},$$

provided that t_0 has been chosen sufficiently large. Recalling that $F'(t) = \frac{\Psi(t)}{t} \geq \frac{\psi(t)}{t}$ this implies that

$$\frac{H(t) + t}{t} = \psi(t) \leq 4A \quad \text{for all } t \geq t_0$$

which concludes the proof. \square

5. LINEAR AND ALMOST-LINEAR BEHAVIOR OF V_C

In this section we prove that, for $N \geq 4$, the generalized Newtonian potential V_C can be written as the sum of a linear function and a correction with sublinear growth towards infinity. In contrast, for $N = 3$, the best one can show is the following *BMO-type property*: on every large ball B_R there exists an affine function A_C^R whose slope grows like $\log R$ and whose average distance from V_C is of order R .

We begin with the case $N \geq 4$.

Lemma 5.1 (Asymptotic growth of the Generalized Newtonian Potential in dimension $N \geq 4$). *Let $N \geq 4$, let u be an x_N -monotone solution in the sense of Definition 2.8, and let V_C be the generalized Newtonian potential (as defined in (3.1)) of the coincidence set C . Then V_C can be written as $V_C := W_C - \ell_C$ with*

$$W_C(x) := \alpha_N \int_C \left(\frac{1}{|x-y|^{N-2}} - \frac{1}{|y|^{N-2}} \right) dy, \quad \ell_C(x) := x \cdot \left(\alpha_N(N-2) \int_C \frac{y}{|y|^N} dy \right),$$

where both integrals are well-defined. Also, there exists a constant C such that

$$\int_{B_R} |W_C(x)| dx \leq CR^{1/2} \quad \text{for all } R \text{ sufficiently large.} \quad (5.1)$$

Proof. Since V_C is well-defined, it suffices to show that ℓ_C is well-defined. For this, we need to prove that $\frac{y}{|y|^N} \chi_C \in L^1(\mathbb{R}^N)$.

Let a_0 and γ_0 be as in Proposition 4.1. Then, since $N \geq 4$, thanks to Proposition 4.1 and noticing that $\frac{1}{|y|} \leq \frac{1}{y_N}$ for $y_N > 0$, we can estimate

$$\begin{aligned} \int_C \frac{1}{|y|^{N-1}} dy &\leq \int_{C \cap \{y_N \leq a_0\}} \frac{1}{|y|^{N-1}} dy + \int_{\{y_N > a_0\} \cap \{|y'|^2 < \gamma_0 y_N\}} \frac{1}{|y|^{N-1}} dy \\ &\leq C + \int_{a_0}^{\infty} \mathcal{H}^{N-1}(B'_{\sqrt{\gamma_0 t}}) \frac{1}{t^{N-1}} dt \leq C \left(1 + \int_{a_0}^{\infty} t^{(N-1)/2 + (1-N)} dt \right) \\ &\leq C \left(1 + \int_{a_0}^{\infty} t^{-3/2} dt \right) < \infty, \end{aligned}$$

which proves that $\ell_C(x)$ is a well-defined linear function.

Now, to prove (5.1), by the mean value theorem it holds

$$\left| \frac{1}{|x-y|^{N-2}} - \frac{1}{|y|^{N-2}} \right| \leq C \frac{|x|}{|y|^{N-1}} \quad \text{for } |y| > 2|x|. \quad (5.2)$$

Hence, given $x \in B_R$ with sufficiently large R , we can write

$$\begin{aligned} |W_C(x)| &\leq \alpha_N \int_{C \cap \{0 \leq y_N \leq a\}} \left| \frac{1}{|x-y|^{N-2}} - \frac{1}{|y|^{N-2}} \right| dy \\ &\quad + \alpha_N \int_{C \cap \{a \leq y_N \leq 2R\}} \left| \frac{1}{|x-y|^{N-2}} - \frac{1}{|y|^{N-2}} \right| dy \\ &\quad + \alpha_N \int_{C \cap \{y_N \geq 2R\}} \left| \frac{1}{|x-y|^{N-2}} - \frac{1}{|y|^{N-2}} \right| dy =: J_1(x) + J_2(x) + J_3(x). \end{aligned}$$

Thanks to (5.2) and Proposition 4.1(i), we can estimate

$$J_3(x) \leq C|x| \int_{2R}^{\infty} \mathcal{H}^{N-1}(\{|y'|^2 < \gamma_0 t\}) \frac{1}{t^{N-1}} dt \leq CR \int_{2R}^{\infty} t^{-3/2} dt \leq CR^{1/2}.$$

Also, thanks to Proposition 4.1(ii), it follows that $|J_1(x)| \leq C$. Finally, for $J_2(x)$ we have

$$\begin{aligned} J_2(x) &\leq \alpha_N \int_{\{|y'|^2 < \gamma_0 y_N\} \cap \{0 \leq y_N \leq 2R\}} \frac{1}{|x-y|^{N-2}} dy + C \int_{a_0}^{2R} \mathcal{H}^{N-1}(B'_{\sqrt{\gamma_0 t}}) \frac{1}{t^{N-2}} dt \\ &\leq \alpha_N \int_{\{|y'|^2 < 2\gamma_0 R\} \cap \{0 \leq y_N \leq 2R\}} \frac{1}{|x-y|^{N-2}} dy + CR^{1/2}. \end{aligned}$$

Now, for the first term on the right hand side, if we write $x = Rz$ with $z \in B_1$ and we perform the change of variables $y \mapsto Ry$, we see that

$$\begin{aligned} \int_{\{|y'|^2 < 2\gamma_0 R\} \cap \{0 \leq y_N \leq 2R\}} \frac{1}{|x - y|^{N-2}} dy &= R^N \int_{\{|y'|^2 < \frac{2\gamma_0}{R}\} \cap \{0 \leq y_N \leq 2\}} \frac{1}{|x - Ry|^{N-2}} dy \\ &= R^2 \int_{\{|y'|^2 < \frac{2\gamma_0}{R}\} \cap \{0 \leq y_N \leq 2\}} \frac{1}{|z - y|^{N-2}} dy. \end{aligned} \quad (5.3)$$

Hence, combining all these bounds, we obtain that

$$\begin{aligned} \int_{B_R} |W_C(x)| dx &\leq CR^{1/2} + \int_{B_R} J_2(x) dx \\ &\leq CR^{1/2} + \alpha_N R^2 \int_{B_1} \int_{\{|y'|^2 < \frac{2\gamma_0}{R}\} \cap \{0 \leq y_N \leq 2\}} \frac{1}{|z - y|^{N-2}} dy dz. \end{aligned}$$

Concerning the last integral we observe that

$$\int_{B_1} \frac{1}{|z - y|^{N-2}} dz \leq C \quad \text{for all } y \in \mathbb{R}^N. \quad (5.4)$$

Thus, since $\left| \left\{ |y'|^2 < \frac{2\gamma_0}{R} \right\} \cap \{0 \leq y_N \leq 2\} \right| \leq \frac{C}{R^{3/2}}$ for $N \geq 4$, it follows from Fubini's Theorem that

$$\begin{aligned} \int_{B_R} |W_C(x)| dx &\leq CR^{1/2} + \alpha_N R^2 \int_{\{|y'|^2 < \frac{2\gamma_0}{R}\} \cap \{0 \leq y_N \leq 2\}} \int_{B_1} \frac{1}{|z - y|^{N-2}} dz dy \\ &\leq CR^{1/2} + CR^2 \left| \left\{ |y'|^2 < \frac{2\gamma_0}{R} \right\} \cap \{0 \leq y_N \leq 2\} \right| \leq CR^{1/2}. \end{aligned}$$

□

We now focus on the three-dimensional case.

Lemma 5.2 (BMO-type estimate in dimension $N = 3$). *Let $N = 3$, let u be an x_N -monotone solution in the sense of Definition 2.8, and let V_C be the generalized Newtonian potential (as defined in Definition 3.1) of the coincidence set \mathcal{C} . Then, for each $R > 0$ the affine function A_C^R given by*

$$A_C^R(x) := \alpha_3 \int_{\mathcal{C}} \left(-\frac{1}{|y|} + \frac{1}{|Re^3 + y|} - \frac{x \cdot y}{|y|^3} + \frac{(Re^3 + y) \cdot (x + Re^3)}{|Re^3 + y|^3} \right) dy$$

is well-defined. Also, there exists a constant C such that

$$\int_{B_R} |V_C(x) - A_C^R(x)| dx \leq CR \quad \text{for all } R \text{ sufficiently large.} \quad (5.5)$$

Proof. We split the proof into two steps.

Step 1. A_C^R is well-defined and affine.

Set

$$a^R(x, y) := -\frac{1}{|y|} + \frac{1}{|Re^3 + y|} - \frac{x \cdot y}{|y|^3} + \frac{(Re^3 + y) \cdot (x + Re^3)}{|Re^3 + y|^3} \quad (5.6)$$

and write

$$\int_{\mathcal{C}} |a^R(x, y)| \, dy \leq \int_{\mathcal{C} \cap \{y_3 \leq 2R\}} |a^R(x, y)| \, dy + \int_{\mathcal{C} \cap \{y_3 \geq 2R\}} |a^R(x, y)| \, dy =: I_1 + I_2.$$

Since $a^R(x, \cdot) \in L^1_{\text{loc}}(\mathbb{R}^3)$ and $\mathcal{C} \cap \{y_3 \leq 2R\}$ is bounded (thanks to Lemma 2.10), it follows that $I_1 \leq C_R$ for some constant depending on R . To estimate I_2 , by a Taylor expansion and the mean value theorem, we have

$$\begin{aligned} |a^R(x, y)| &\leq \left| \frac{1}{|Re^3 + y|} - \frac{1}{|y|} + \frac{(Re^3) \cdot y}{|y|^3} \right| + \frac{|Re^3 \cdot (x + Re^3)|}{|Re^3 + y|^3} \\ &\quad + \left| (Re^3 + x) \cdot y \left(\frac{1}{|y|^3} - \frac{1}{|Re^3 + y|^3} \right) \right| \\ &\leq C \frac{R^2}{|y|^3} + C \frac{R(R + |x|)}{|y|^3} + C \frac{R(R + |x|)}{|y|^3} \quad \text{for } |y| > 2R. \end{aligned} \quad (5.7)$$

Hence, recalling Proposition 4.1(i), we can estimate (here and below, a_0 and γ_0 are the constants from Proposition 4.1)

$$\begin{aligned} I_2 &\leq CR(R + |x|) \int_{\mathcal{C} \cap \{y_3 \geq 2R\}} \frac{1}{|y|^3} \, dy \leq CR(R + |x|) \int_{2R}^{\infty} \mathcal{H}^2(\{|y'|^2 < \gamma_0 t\}) \frac{1}{t^3} \, dt \\ &\leq CR(R + |x|) \int_{2R}^{\infty} \frac{1}{t^2} \, dt \leq C(R + |x|), \end{aligned} \quad (5.8)$$

which proves that $A_{\mathcal{C}}^R$ is well-defined.

Observe now that the integrand a^R in the definition of $A_{\mathcal{C}}^R$ is integrable (by what we have just seen) and differentiable in x . Also, for each $y \in \mathcal{C}$ and $b := \max\{a_0, R\}$, it holds

$$\begin{aligned} |\nabla_x a^R(x, y)| &= \left| \frac{y}{|y|^3} - \frac{Re^3 + y}{|Re^3 + y|^3} \right| \\ &\leq \frac{2}{|y|^2} \chi_{\mathcal{C} \cap \{y_3 \leq b\}}(y) + \left| \int_0^1 \frac{d}{ds} \left(\frac{sRe^3 + y}{|sRe^3 + y|^3} \right) ds \right| \chi_{\mathcal{C} \cap \{y_3 \geq b\}}(y) \\ &\leq \frac{2}{|y|^2} \chi_{\mathcal{C} \cap \{y_3 \leq b\}}(y) + C \frac{R}{|y|^3} \chi_{\mathcal{C} \cap \{y_3 \geq b\}}(y). \end{aligned} \quad (5.9)$$

Since the right-hand side is integrable in \mathbb{R}^3 (again, thanks to Proposition 4.1), it follows from dominated convergence that

$$\nabla A_{\mathcal{C}}^R(x) = \alpha_3 \int_{\mathcal{C}} \nabla_x a^R(x, y) \, dy = \alpha_3 \int_{\mathcal{C}} \left(-\frac{y}{|y|^3} + \frac{Re^3 + y}{|Re^3 + y|^3} \right) \, dy, \quad (5.10)$$

which is constant in \mathbb{R}^3 . This proves that $A_{\mathcal{C}}^R$ is an affine function.

Step 2. *Proof of (5.5).* Recalling the definition of $G(x, y)$ in Definition 3.1, we have

$$\begin{aligned} |V_{\mathcal{C}}(x) - A_{\mathcal{C}}^R(x)| &\leq \alpha_3 \int_{\mathcal{C} \cap \{0 \leq y_3 \leq 2R\}} |G(x, y) - a^R(x, y)| \, dy \\ &\quad + \alpha_3 \int_{\mathcal{C} \cap \{y_3 \geq 2R\}} |G(x, y) - a^R(x, y)| \, dy =: J_1(x) + J_2(x). \end{aligned}$$

By (3.2) and (5.7), for $x \in B_R$ with $R \geq a_0$ we can estimate

$$J_2(x) \leq CR^2 \int_{2R}^{\infty} \mathcal{H}^2(\{|y'|^2 < \gamma_0 t\}) \frac{1}{t^3} dt \leq CR \quad (5.11)$$

(cp. (5.8)). For $J_1(x)$, thanks to Proposition 4.1, for R sufficiently large we have

$$\mathcal{C} \cap \{y_3 \leq 2R\} \subset \{|y'|^2 < 2\gamma_0 R\} \times \{0 \leq y_3 \leq 2R\}.$$

Thus, since

$$G(Rx, Ry) - a^R(Rx, Ry) = \frac{1}{R}[G(x, y) - a^1(x, y)],$$

if we write $x = Rz$ with $z \in B_1$ and perform the change of variables $y \mapsto Ry$, we see that

$$\begin{aligned} \int_{\mathcal{C} \cap \{y_3 \leq 2R\}} |G(x, y) - a^R(x, y)| dy &\leq \int_{\{|y'|^2 < 2\gamma_0 R\} \cap \{0 \leq y_3 \leq 2R\}} |G(x, y) - a^R(x, y)| dy \\ &= R^2 \int_{\{|y'|^2 < \frac{2\gamma_0}{R}\} \cap \{0 \leq y_3 \leq 2\}} |G(z, y) - a^1(z, y)| dy \end{aligned}$$

(cp. (5.3)). Altogether we proved that

$$\begin{aligned} \int_{B_R} |V_{\mathcal{C}}(x) - A_{\mathcal{C}}^R(x)| dx &\leq \int_{B_R} (J_1(x) + J_2(x)) dx \leq CR + \int_{B_R} J_1(x) dx \\ &\leq CR + \alpha_3 R^2 \int_{B_1} \int_{\{|y'|^2 < \frac{2\gamma_0}{R}\} \cap \{0 \leq y_3 \leq 2\}} |G(z, y) - a^1(z, y)| dy dz. \end{aligned} \quad (5.12)$$

Concerning the last integral, we observe that, on the domain of integration, $|e^3 + y| \geq 1$ and $|z + e^3| \leq 3$. Hence, we can estimate

$$|G(z, y) - a^1(z, y)| = \left| \frac{1}{|z - y|} - \frac{1}{|e^3 + y|} - \frac{(e^3 + y) \cdot (z + e^3)}{|e^3 + y|^3} \right| \leq \frac{1}{|z - y|} + 4. \quad (5.13)$$

Thus, since $\left| \left\{ |y'|^2 < \frac{2\gamma_0}{R} \right\} \cap \{0 \leq y_3 \leq 2\} \right| \leq \frac{C}{R}$, it follows from Fubini's Theorem and (5.4) that

$$\begin{aligned} \int_{B_R} |V_{\mathcal{C}}(x) - A_{\mathcal{C}}^R(x)| dx &\leq CR + \alpha_3 R^2 \int_{\{|y'|^2 < \frac{2\gamma_0}{R}\} \cap \{0 \leq y_3 \leq 2\}} \int_{B_1} \left(\frac{1}{|z - y|} + 4 \right) dz dy \\ &\leq CR + CR^2 \left| \left\{ |y'|^2 < \frac{2\gamma_0}{R} \right\} \cap \{0 \leq y_3 \leq 2\} \right| \leq CR, \end{aligned} \quad (5.14)$$

as desired. \square

We conclude this section by proving a sharp bound on the growth of $A_{\mathcal{C}}^R(0)$ and the different components of its gradient. This result will be used later in the proofs of Corollary 6.5, Proposition 8.1, and Proposition 8.2.

Lemma 5.3 (Growth of $A_{\mathcal{C}}^R$). *Let $N = 3$, let u be an x_N -monotone solution in the sense of Definition 2.8, and let $A_{\mathcal{C}}^R$ be as in Lemma 5.2. Then there exists a constant C such that*

$$|\nabla' A_{\mathcal{C}}^R| \leq C, \quad |\partial_3 A_{\mathcal{C}}^R| \leq C \log R, \quad |A_{\mathcal{C}}^R(0)| \leq CR \quad \text{for all } R \text{ sufficiently large.}$$

(Here, ∇' denotes the gradient with respect to the first 2 variables.)

Proof. For the first bound we note that, for $y \in \mathcal{C}$,

$$|\nabla'_x a^R(x, y)| = \left| \frac{y'}{|y|^3} - \frac{y'}{|Re^3 + y|^3} \right| \leq 2 \frac{|y'|}{|y|^3}, \quad (5.15)$$

where a^R is the function defined in (5.6). Thus, thanks to Proposition 4.1,

$$\begin{aligned} \int_{\mathcal{C}} |\nabla'_x a^R(x, y)| \, dy &\leq 2 \int_{\mathcal{C} \cap \{y_3 \leq a_0\}} \frac{1}{|y|^2} \, dy + 2 \int_{\mathcal{C} \cap \{y_3 \geq a_0\}} \frac{|y'|}{y_3^3} \, dy \\ &\leq C + 2 \int_{a_0}^{\infty} \mathcal{H}^2(\{|y'|^2 < \gamma_0 t\}) \frac{(\gamma_0 t)^{1/2}}{t^3} \, dt \leq C + C \int_{a_0}^{\infty} \frac{1}{t^{3/2}} \, dt \leq C. \end{aligned}$$

Recalling (5.10), this proves that $|\nabla' A_{\mathcal{C}}^R| \leq C$.

For the second bound, we apply (5.9) and Proposition 4.1 to obtain that, for $R \geq a_0$,

$$\begin{aligned} \int_{\mathcal{C}} |\partial_{x_3} a^R(x, y)| \, dy &\leq C \int_{\mathcal{C} \cap \{y_3 \leq a_0\}} \frac{1}{|y|^2} \, dy + C \int_{\mathcal{C} \cap \{a_0 \leq y_3 \leq R\}} \frac{1}{|y|^2} \, dy + C \int_{\mathcal{C} \cap \{y_3 \geq R\}} \frac{R}{|y|^3} \, dy \\ &\leq C + C \int_{a_0}^R \mathcal{H}^2(\{|y'|^2 < \gamma_0 t\}) \frac{1}{t^2} \, dt + CR \int_R^{\infty} \mathcal{H}^2(\{|y'|^2 < \gamma_0 t\}) \frac{1}{t^3} \, dt \\ &\leq C + C \int_{a_0}^R \frac{1}{t} \, dt + CR \int_R^{\infty} \frac{1}{t^2} \, dt \leq C \log R. \end{aligned}$$

Thus, $|\partial_3 A_{\mathcal{C}}^R| \leq C \log R$.

Finally, in order to estimate $A_{\mathcal{C}}^R(0)$, we write

$$\int_{\mathcal{C}} |a^R(0, y)| \, dy \leq \int_{\mathcal{C} \cap \{y_3 \leq 2R\}} |a^R(0, y)| \, dy + \int_{\mathcal{C} \cap \{y_3 \geq 2R\}} |a^R(0, y)| \, dy =: J_1 + J_2.$$

Using (5.8) with $x = 0$, we immediately get $J_2 \leq CR$. Concerning J_1 , for $y \in \mathcal{C} \cap \{y_3 \leq 2R\}$ we can estimate

$$|a^R(0, y)| \leq \frac{2}{|y|} + \frac{R}{R^2 + |y|^2} \leq \frac{3}{|y|}, \quad (5.16)$$

hence

$$\begin{aligned} J_1 &\leq 3 \int_{\mathcal{C} \cap \{y_3 \leq a_0\}} \frac{1}{|y|} \, dy + C \int_{\mathcal{C} \cap \{a_0 \leq y_3 \leq 2R\}} \frac{1}{|y|} \, dy \\ &\leq C + C \int_{a_0}^{2R} \mathcal{H}^2(\{|y'|^2 < \gamma_0 t\}) \frac{1}{t} \, dt \leq C + C \int_{a_0}^{2R} \frac{1}{t} \, dt \leq CR, \end{aligned} \quad (5.17)$$

concluding the proof. □

6. CONSTRUCTING MATCHING PARABOLOID SOLUTIONS

In this section, given $N \geq 3$, we construct “matching” paraboloid solutions, i.e., solutions that have paraboloids as coincidence sets and have the same second-order asymptotics at infinity as the solution u . More precisely, for $N \geq 4$ we find a fixed paraboloid solution u_P such that $u - u_P$ grows sublinearly at infinity. Instead, in the critical dimension $N = 3$, for any sufficiently large R we construct a paraboloid solution u_{P_R} such that $\sup_{B_R} |u - u_{P_R}| \leq CR$, where the constant C is independent of R .

We begin with a result about the existence of paraboloid solutions with prescribed asymptotic behavior at infinity. Since the proof is rather classical and follows from a minor modification of the one of [7, Theorem 7.1], we postpone it to Appendix A.

Lemma 6.1 (Existence of paraboloid solutions with prescribed asymptotic behavior at infinity). *Given $N \geq 3$, let $p = p(x')$ be a homogeneous quadratic polynomial as in Definition 2.8(iv). Then there exists a (unique) ellipsoid*

$$E' := \left\{ y' \in \mathbb{R}^{N-1} : \frac{y_1^2}{a_1^2} + \dots + \frac{y_{N-1}^2}{a_{N-1}^2} \leq 1 \right\} \subset \mathbb{R}^{N-1} \quad \text{with } a_i > 0 \text{ for } i = 1, \dots, N-1,$$

such that the following holds.

Define the paraboloid $P = P_{E'} := \{(y', y_N) \in \mathbb{R}^{N-1} \times [0, \infty) : y' \in \sqrt{y_N} E'\}$. Then there exists a global solution u_P with P as coincidence set and p as quadratic blow-down limit, i.e.,

$$\Delta u_P = \chi_{\{u_P > 0\}}, \quad u_P \geq 0 \quad \text{in } \mathbb{R}^N, \quad \{u_P = 0\} = P,$$

and

$$\frac{u_P(rx)}{r^2} \rightarrow p(x') \quad \text{in } C_{\text{loc}}^{1,\alpha}(\mathbb{R}^N) \text{ as } r \rightarrow \infty, \quad \alpha \in (0, 1).$$

Also, for each $\gamma > 0$, the function $u_{\gamma P}(x) := \gamma^2 u_P(\frac{x}{\gamma})$ is a global solution with γP as coincidence set and p as blow-down limit, and it satisfies the potential expansion

$$u_{\gamma P} = p + V_{\gamma P} \quad \text{in } \mathbb{R}^N. \quad (6.1)$$

In the next result, we show that, if $N \geq 4$, we can actually find paraboloid solutions with prescribed behavior up to linear order. Given $a \in \mathbb{R}^N$, we use the notation $u_{\gamma P - a}$ to denote the solution that has $\gamma P - a$ as coincidence set. Note that this solution is obtained simply by translating the solution having γP as coincidence set, namely $u_{\gamma P - a}(x) = u_{\gamma P}(x + a)$.

Lemma 6.2 (Existence of paraboloid solutions with prescribed linear behavior at infinity in dimension $N \geq 4$). *Let $N \geq 4$, let $p = p(x')$ be a homogeneous quadratic polynomial as in Definition 2.8(iv), and let P be as in Lemma 6.1. For any $b = (b', b_N) \in \mathbb{R}^{N-1} \times (-\infty, 0)$ there exist $\tau' \in \mathbb{R}^{N-1}$ and $\gamma > 0$ such that the following holds: for each $\sigma \in \mathbb{R}$,*

$$\frac{1}{R} \int_{B_R} |u_{\gamma P - (\tau', \sigma)}(x) - p(x') - b \cdot x| dx \rightarrow 0 \quad \text{as } R \rightarrow \infty.$$

Proof. As noted before, $u_{\gamma P - (\tau', \sigma)}(x) = u_{\gamma P}(x' + \tau', x_N + \sigma)$. Hence, since p is a quadratic polynomial, recalling Lemma 5.1 we have

$$\begin{aligned} u_{\gamma P - (\tau', \sigma)}(x) &= p(x' + \tau') + V_{\gamma P}(x' + \tau', x_N + \sigma) \\ &= p(x') + \nabla' p(x') \cdot \tau' + p(\tau') \\ &\quad - (x' + \tau', x_N + \sigma) \cdot \left(\alpha_N (N-2) \int_{\gamma P} \frac{y}{|y|^N} dy \right) + W_{\gamma P}(x' + \tau', x_N + \sigma), \end{aligned} \quad (6.2)$$

where $\nabla' p$ denotes the first $N-1$ components of $\nabla p = (\nabla' p, 0) \in \mathbb{R}^{N-1} \times \mathbb{R}$ and $W_{\gamma P}$ is defined analogously to W_C in Lemma 5.1, namely

$$W_{\gamma P}(x) := \alpha_N \int_{\gamma P} \left(\frac{1}{|x-y|^{N-2}} - \frac{1}{|y|^{N-2}} \right) dy.$$

Note now that, by symmetry,

$$\alpha_N(N-2) \int_{\gamma P} \frac{y}{|y|^N} dy = \lambda_\gamma e^N, \quad \text{where } \lambda_\gamma := \alpha_N(N-2) \int_{\gamma P} \frac{y_N}{|y|^N} dy > 0.$$

Thus

$$u_{\gamma P - (\tau', \sigma)}(x) = p(x') + \nabla' p(x') \cdot \tau' + p(\tau') - \lambda_\gamma(x_N + \sigma) + W_{\gamma P}(x' + \tau', x_N + \sigma).$$

Recalling that $\nabla' p(x') = Qx'$ with Q symmetric and invertible, choosing $\tau' := Q^{-1}b'$ we get $\nabla' p(x') \cdot \tau' = b' \cdot x'$.

In addition, it follows by monotone convergence that $\gamma \mapsto \lambda_\gamma$ is continuous and

$$\begin{aligned} \lambda_\gamma &\rightarrow \alpha_N(N-2) \int_{\mathbb{R}^{N-1} \times (0, \infty)} \frac{y_N}{|y|^N} dy = \infty & \text{as } \gamma \rightarrow \infty, \\ \lambda_\gamma &\rightarrow 0 & \text{as } \gamma \rightarrow 0. \end{aligned}$$

Thus, by continuity, there exists $\gamma > 0$ such that $\lambda_\gamma = -b_N$. Hence, with these choices of τ' and γ , we get

$$u_{\gamma P - (\tau', \sigma)}(x) - p(x') - b \cdot x = p(\tau') - \lambda_\gamma \sigma + W_{\gamma P}(x' + \tau', x_N + \sigma).$$

Applying Lemma 5.1 with γP in place of \mathcal{C} , this implies that for sufficiently large R (depending on τ' , γ , and σ),

$$\begin{aligned} \int_{B_R} |u_{\gamma P - (\tau', \sigma)}(x) - p(x') - b \cdot x| dx &\leq |p(\tau')| + \lambda_\gamma |\sigma| + \int_{B_R} |W_{\gamma P}(x' + \tau', x_N + \sigma)| dx \\ &\leq |p(\tau')| + \lambda_\gamma |\sigma| + \frac{1}{|B_R|} \int_{B_{2R}} |W_{\gamma P}(x)| dx \leq CR^{1/2}, \end{aligned}$$

and the result follows. □

Corollary 6.3. *Let $N \geq 4$, and let u be an x_N -monotone solution in the sense of Definition 2.8. Then there exist a paraboloid P as in Lemma 6.1, $\gamma > 0$, and $\tau' \in \mathbb{R}^{N-1}$ such that, for each $\sigma \in \mathbb{R}$,*

$$\frac{1}{R} \int_{B_R} |u - u_{\gamma P - (\tau', \sigma)}| dx \rightarrow 0 \quad \text{as } R \rightarrow \infty.$$

Proof. Let p be as in Definition 2.8, let \mathcal{C} be the coincidence set of u , and let $\ell_{\mathcal{C}}$ be as in Lemma 5.1, so that $u = p - \ell_{\mathcal{C}} + W_{\mathcal{C}}$. Define

$$b = -\nabla \ell_{\mathcal{C}} = -\alpha_N(N-2) \int_{\mathcal{C}} \frac{y}{|y|^N} dy.$$

Since $b_N = -\alpha_N(N-2) \int_{\mathcal{C}} \frac{y_N}{|y|^N} dy < 0$, we can apply Lemmas 6.2 and 5.1 to deduce the existence of P , γ , and τ' such that, for each $\sigma \in \mathbb{R}$,

$$\frac{1}{R} \int_{B_R} |u - u_{\gamma P - (\tau', \sigma)}| dx \leq \frac{1}{R} \int_{B_R} |W_{\mathcal{C}}(x)| dx + \frac{1}{R} \int_{B_R} |u_{\gamma P - (\tau', \sigma)} - p(x') - b \cdot x| dx \rightarrow 0$$

as $R \rightarrow \infty$. □

In the critical dimension $N = 3$, the statement of Lemma 6.2 does not hold. The best we can do in dimension $N = 3$ is to match, on any large ball B_R , the slope of the affine approximation of $V_{\gamma R P}$ in the e^3 -direction. This will be achieved in the proof of Corollary 6.5, using Lemma 6.4 below. As a consequence, on any large ball B_R , we can find a paraboloid $P_R = \gamma R P$ so that the average of $u - u_{P_R}$ over B_R grows at most linearly in R (see Corollary 6.5).

Lemma 6.4 (Matching on each ball B_R in dimension $N = 3$). *Let $N = 3$, $p = p(x')$ be the blow-down polynomial defined in Definition 2.8(iv), and P as in Lemma 6.1. Also, for $\gamma \geq 0$, let $A_{\gamma P}^R$ be defined as in Lemma 5.2.*

Then, given $B > 0$, there exist $\gamma_B > 0$ and $R_B \geq 1$ such that the following holds: For each $\beta \in [0, B]$ and $R \geq R_B$ there exists $\gamma = \gamma(\beta, R) \in [0, \gamma_B]$ such that

$$\partial_3 A_{\gamma P}^R = -\beta \log R. \quad (6.3)$$

Proof. Note that, since $u_{\gamma P}$ is an x_N -monotone solution, $A_{\gamma P}^R$ is well-defined thanks to Lemma 5.2. Also, as shown in Step 1 in the proof of Lemma 5.2,

$$\partial_3 A_{\gamma P}^R = \alpha_3 \int_{\gamma P} \left(-\frac{y_3}{|y|^3} + \frac{R + y_3}{|Re^3 + y|^3} \right) dy. \quad (6.4)$$

We now observe that

$$\begin{aligned} \int_{\gamma P} \left(-\frac{y_3}{|y|^3} + \frac{R + y_3}{|Re^3 + y|^3} \right) dy &= \lim_{\varepsilon \rightarrow 0} \int_{\gamma P} \left(-\frac{y_3}{|y|^{3+\varepsilon}} + \frac{R + y_3}{|Re^3 + y|^{3+\varepsilon}} \right) dy \\ &= \lim_{\varepsilon \rightarrow 0} - \int_{\gamma P} \frac{y_3}{|y|^{3+\varepsilon}} dy + \int_{\gamma P} \frac{R + y_3}{|Re^3 + y|^{3+\varepsilon}} dy \\ &= \lim_{\varepsilon \rightarrow 0} - \int_{\gamma P \setminus (\gamma P + Re^3)} \frac{y_3}{|y|^{3+\varepsilon}} dy = - \int_{\gamma P \setminus (\gamma P + Re^3)} \frac{y_3}{|y|^3} dy, \end{aligned}$$

where the first equality follows from dominated convergence since the integrand is uniformly convergent at infinity (see (5.9)), the second equality from the fact that each term in the integrand is integrable for $\varepsilon > 0$, the third equality from a change of variables, and the last equality from monotone convergence when $|y| > 1$ and dominated convergence when $|y| \leq 1$. This proves that

$$\partial_3 A_{\gamma P}^R = -\alpha_3 \int_{\gamma P \setminus (\gamma P + Re^3)} \frac{y_3}{|y|^3} dy < 0, \quad (6.5)$$

In particular, since $\gamma P + Re^3 \subset \{y_3 \geq R\}$, for $R > 1$ it follows that

$$\partial_3 A_{\gamma P}^R \leq -\alpha_3 \int_{\gamma P \cap \{R^{1/2} < y_3 < R\}} \frac{y_3}{|y|^3} dy. \quad (6.6)$$

Note now that, for $R^{1/2} \geq \gamma$ and for $y' \in \sqrt{\gamma y_3} E'$ with $y_3 \geq R^{1/2}$, we have

$$|y| \leq |y'| + y_3 \leq C_{E'} \gamma^{1/2} y_3^{1/2} + y_3 \leq C_{E'} R^{1/4} y_3^{1/2} + y_3 \leq (C_{E'} + 1) y_3,$$

for some constant $C_{E'}$ depending only on E' . Hence, thanks to (6.6),

$$\begin{aligned} \partial_3 A_{\gamma P}^R &\leq -\frac{\alpha_3}{(C_{E'} + 1)^3} \int_{\gamma P \cap \{R^{1/2} < y_3 < R\}} \frac{1}{y_3^2} dy \leq -\frac{\alpha_3}{(C_{E'} + 1)^3} \int_{R^{1/2}}^R \mathcal{H}^2(\sqrt{\gamma t} E') \frac{1}{t^2} dt \\ &\leq -c_{E'} \gamma \int_{R^{1/2}}^R \frac{1}{t} dt = -\frac{c_{E'}}{2} \gamma \log R, \end{aligned} \quad (6.7)$$

where $c_{E'} > 0$ is a constant depending only on E' .

Hence, given $B > 0$, set $\gamma_B := \frac{2}{c_{E'}} B$ and $R_B := \max\{\gamma_B^2, 1\}$. Then, with these choices,

$$\partial_3 A_{\gamma_B P}^R \leq -B \log R \quad \text{for all } R \geq R_B.$$

On the other hand, recalling (6.4) and (5.9), we can apply dominated convergence to deduce that $\gamma \mapsto \partial_3 A_{\gamma P}^R$ is continuous and

$$\partial_3 A_{\gamma P}^R \rightarrow 0 \quad \text{as } \gamma \rightarrow 0.$$

Hence, by continuity, given any $\beta \in [0, B]$ and $R \geq R_B$ there exists $\gamma = \gamma(\beta, R) \in [0, \gamma_B]$ such that (6.3) holds. \square

Corollary 6.5. *Let $N = 3$, and let u be an x_N -monotone solution in the sense of Definition 2.8. Then there exist a paraboloid P as in Lemma 6.1 and positive constants $\bar{\gamma}$, \bar{R} , and \bar{C} , such that the following holds: for any $R \geq \bar{R}$ there exists $\gamma_R \in [0, \bar{\gamma}]$ such that*

$$\frac{1}{R} \int_{B_R} |u - u_{P_R}| dx \leq \bar{C}, \quad \text{where } P_R := \gamma_R P. \quad (6.8)$$

Proof. We begin by noticing that, since \mathcal{C} is contained in some paraboloid (see Proposition 4.1), we can repeat the proof of (6.5) with \mathcal{C} in place of γP to show that

$$\partial_3 A_{\mathcal{C}}^R = -\alpha_3 \int_{\mathcal{C} \setminus (C + R e^3)} \frac{y_3}{|y|^3} dy.$$

We now observe that, as a consequence of the monotonicity of the contact set in the e^3 -direction (a direct consequence of the monotonicity of u), the right hand side above is strictly negative. Thus, thanks to Lemma 5.3, there exist $\hat{R} > 1$ and $B > 0$ such that

$$0 > \partial_3 A_{\mathcal{C}}^R \geq -B \log R \quad \text{for all } R \geq \hat{R}.$$

This allows us to apply Lemma 6.4 to deduce that, if we set $\bar{\gamma} := \gamma_B$ and $\bar{R} := \max\{\hat{R}, R_B\}$, then for any $R \geq \bar{R}$ there exists $\gamma_R \in (0, \bar{\gamma}]$ such that

$$\partial_3 A_{P_R}^R = \partial_3 A_{\mathcal{C}}^R < 0, \quad \text{where } P_R := \gamma_R P. \quad (6.9)$$

Using the potential expansion of both u and u_{P_R} (cf. Proposition 3.4 and (6.1)), thanks to (6.9) we find that for all $R \geq \bar{R}$,

$$\begin{aligned} \int_{B_R} |u - u_{P_R}| dx &= \int_{B_R} |V_{\mathcal{C}} - V_{P_R}| dx \leq \int_{B_R} |V_{\mathcal{C}} - A_{\mathcal{C}}^R| dx + \int_{B_R} |V_{P_R} - A_{P_R}^R| dx \\ &\quad + \int_{B_R} |A_{\mathcal{C}}^R(0) - A_{P_R}^R(0)| dx + \int_{B_R} |\nabla' A_{\mathcal{C}}^R - \nabla' A_{P_R}^R| |x| dx. \end{aligned}$$

Applying Lemmas 5.2 and 5.3 to u , we can estimate

$$\int_{B_R} |V_C - A_C^R| dx + \int_{B_R} |A_C^R(0)| dx + \int_{B_R} |\nabla' A_C^R| |x| dx \leq CR.$$

Also, since $\gamma_R \in (0, \bar{\gamma}]$, the very same arguments used for proving Lemmas 5.2 and 5.3 show that

$$\int_{\dot{B}_R} |V_{P_R} - A_{P_R}^R| dx + \int_{B_R} |A_{P_R}^R(0)| dx + \int_{\dot{B}_R} |\nabla' A_{P_R}^R| |x| dx \leq C_{\bar{\gamma}} R,$$

where $C_{\bar{\gamma}}$ depends only on P and $\bar{\gamma}$ (in particular, it is independent of R). Combining all these estimates, we conclude the validity of (6.8). \square

7. PROOF OF THEOREM 1.2: THE CASE $N \geq 4$

Given u an x_N -monotone solution as in Definition 2.8, using the ACF monotonicity formula from Lemma 2.12 we will show that u and the comparison solutions $u_{\gamma P - (\tau', \sigma)}$ provided by Corollary 6.3 are ordered. Thanks to this important fact, the result will follow easily.

In order to simplify notation, we set

$$P_\sigma := \gamma P - (\tau', \sigma) \quad \text{and} \quad u_\sigma := u_{\gamma P - (\tau', \sigma)}. \quad (7.1)$$

Proposition 7.1 (Ordering in dimension $N \geq 4$). *Let $N \geq 4$. Then, for all $\sigma \in \mathbb{R}$,*

$$\text{either } u \leq u_\sigma \text{ in } \mathbb{R}^N \quad \text{or} \quad u \geq u_\sigma \text{ in } \mathbb{R}^N.$$

Proof. Thanks to Lemma 2.13(i), we can apply Lemma 2.12(i)-(iii) with $v = u - u_\sigma$ to deduce that, for every $r > 0$,

$$\Phi(u - u_\sigma, r) \leq \limsup_{R \rightarrow \infty} \Phi(u - u_\sigma, R) \leq C_N \limsup_{R \rightarrow \infty} \left(\frac{1}{R} \int_{B_{4R}} |u - u_\sigma| \right)^4 = 0,$$

where the last equality follows from Corollary 6.3. Hence, thanks to Lemma 2.12(iv) we conclude that either $u - u_\sigma \geq 0$ or $u - u_\sigma \leq 0$, as desired. \square

We can now easily prove our main theorem.

Proof of Theorem 1.2: the case $N \geq 4$. Since u is an x_N -monotone solution,

$$u(0) = 0, \quad u(-e^N) > 0. \quad (7.2)$$

On the other hand, recalling (7.1), since P is a paraboloid contained in $\{x_N \geq 0\}$ with tip at the origin, by the definition of P_σ (see (7.1)) it follows that

$$0 \notin P_\sigma \quad \text{for } \sigma < 0, \quad -e^N \in P_\sigma \quad \text{for } \sigma \gg 1,$$

therefore

$$u_\sigma(0) > 0 \quad \text{for } \sigma < 0, \quad u_\sigma(-e^N) = 0 \quad \text{for } \sigma \gg 1. \quad (7.3)$$

Combining (7.2), (7.3), and Proposition 7.1, we conclude that

$$u_\sigma \geq u \quad \text{for } \sigma < 0, \quad u_\sigma \leq u \quad \text{for } \sigma \gg 1. \quad (7.4)$$

Now, let us define

$$\bar{\sigma} := \inf\{\sigma \in \mathbb{R} : u_\sigma \leq u\}.$$

Thanks to (7.4), $\bar{\sigma} \in \mathbb{R}$ is well-defined. We now claim that $u \equiv u_{\bar{\sigma}}$.

Indeed, by definition of $\bar{\sigma}$ there exists a sequence $\sigma_k \rightarrow \bar{\sigma}$ such that $u_{\sigma_k} \leq u$, therefore $u_{\bar{\sigma}} \leq u$. Assume now towards a contradiction that $u \not\equiv u_{\bar{\sigma}}$. Then there exists $\bar{x} \in \mathbb{R}^N$ such that $u_{\bar{\sigma}}(\bar{x}) < u(\bar{x})$, and by continuity we can find $\varepsilon > 0$ such that $u_{\bar{\sigma}-\varepsilon}(\bar{x}) < u(\bar{x})$. Since u and $u_{\bar{\sigma}-\varepsilon}$ must be ordered (because of Proposition 7.1), we conclude that $u_{\bar{\sigma}-\varepsilon} \leq u$, contradicting the definition of $\bar{\sigma}$.

Since $u \equiv u_{\bar{\sigma}}$ we conclude that $\{u = 0\}$ is a paraboloid⁴, as desired. \square

Remark 7.2. It is worth noticing that our argument gives a new proof of the characterization of global solutions with compact coincidence set for any dimension $N \geq 2$. Indeed, when $\mathcal{C} = \{u = 0\}$ is compact, we can write the expansion (cp. Lemma 5.1 and [9])

$$u(x) = p(x) + V_{\mathcal{C}}(x) = p(x) - x \cdot \int_{\mathcal{C}} \nabla K_N(y) dy + \int_{\mathcal{C}} \left(K_N(x - y) - K_N(y) \right) dy,$$

where K_N is the fundamental solution of the Laplacian (see (A.1)). Since \mathcal{C} is compact, all integrals converge and the remainder term (the last integral) is sublinear. Also, in this compact case, $p(x) = \frac{1}{2}x^T Qx$ where $Q \in \mathbb{R}^{N \times N}$ is symmetric and positive definite⁵.

Now, arguing as in Lemma 6.1, we find an ellipsoid $E \subset \mathbb{R}^N$ such that u_E has p as quadratic blow-down limit. In addition, since Q is invertible on \mathbb{R}^N , choosing $\tau := Q^{-1}b$ with $b := \int_{\mathcal{C}} \nabla K_N(y) dy$, for each $\gamma > 0$ the function $u - u_{\gamma E - \tau}$ has sublinear growth at infinity (cp. Lemma 6.2). Then the ACF monotonicity formula implies that either $u \leq u_{\gamma E - \tau}$ or $u \geq u_{\gamma E - \tau}$ (cp. Proposition 7.1), and finally a continuity argument implies the existence of a value $\bar{\gamma} > 0$ such that $u \equiv u_{\bar{\gamma} E - \tau}$ (cp. Proof of Theorem 1.2: the case $N \geq 4$), as desired.

8. PROOF OF THEOREM 1.2: THE CASE $N = 3$

Let u be an x_N -monotone solution as in Definition 2.8. Also, for each $R \geq \bar{R}$, let $P_R = \gamma_R P$ with $\gamma_R \in [0, \bar{\gamma}]$ be the paraboloid provided by Corollary 6.5, so that (6.8) holds.

To simplify the notation, for each $R \geq \bar{R}$ and $x \in \mathbb{R}^3$ we define

$$v_R(x) := \frac{(u - u_{P_R})(Rx)}{R},$$

so that (6.8) is equivalent to $\|v_R\|_{L^1(B_1)} \leq C$ for all $R \geq \bar{R}$.

Thanks to this uniform L^1 -bound and Lemma 2.13(i), we can apply Lemma 2.12(i)-(iii) to deduce that, for all $r \in (2\bar{R}, R)$,

$$\Phi(u - u_{P_R}, \frac{r}{2}) \leq \Phi(u - u_{P_R}, \frac{R}{2}) = \Phi(v_R, \frac{1}{2}) \leq C. \tag{8.1}$$

Since $0 \in \{u_{P_R} = 0\}$, it follows from Lemma 2.4(ii) that, passing if necessary to a subsequence,

$$u_{P_R} \rightarrow u_{\infty} \quad \text{in } C_{\text{loc}}^{1,\alpha}(\mathbb{R}^3) \quad \text{as } R \rightarrow \infty, \tag{8.2}$$

where u_{∞} is a global solution to the obstacle problem. Also, since $P_R = \gamma_R P$ with $\gamma_R \in [0, \bar{\gamma}]$, it follows that

$$\{u_{\infty} = 0\} = \gamma_{\infty} P \quad \text{for some } \gamma_{\infty} \in [0, \bar{\gamma}], \quad P = \{y' \in \sqrt{y_N} E'\} \tag{8.3}$$

(if $\gamma_{\infty} = 0$ then $u_{\infty} = p$).

⁴A posteriori, by the fact that $\{u = 0\}$ is a convex set contained in $\{x_N \geq 0\}$ with tip at the origin, the only possibility is that $\bar{\sigma} = 0$ and $\tau' = 0$. However this information is not relevant for our proof.

⁵This follows, for instance, from the proof of Theorem 1.1 in Section 9.

Thanks to (8.2) and Fatou's Lemma, (8.1) implies that $\Phi(u - u_\infty, \frac{r}{2}) \leq C$ for all $r \geq 2\bar{R}$. Hence, since Φ is non-decreasing in r (see Lemma 2.12(i)), we obtain that

$$\Phi(u - u_\infty, r) \leq C \quad \text{for all } r > 0. \quad (8.4)$$

8.1. Linear rescaling and ACF dichotomy. Let us now introduce the linear rescaling

$$w_r(x) := \frac{(u - u_\infty)(rx)}{r} \quad \text{for } x \in \mathbb{R}^3 \text{ and } r > 0. \quad (8.5)$$

We prove the following important dichotomy.

Proposition 8.1 (ACF alternative for $u - u_\infty$). *Let u_∞ , γ_∞ , and P be as in (8.2)-(8.3), and w_r as in (8.5). Then there exists a sequence $r_k \rightarrow \infty$ such that $w_{r_k} \rightarrow w$ strongly in $W^{1,2}(B_1)$ as $k \rightarrow \infty$. In addition:*

- (i) either w has constant sign inside B_1 (i.e., either $w \geq 0$ or $w \leq 0$ a.e. in B_1);
- (ii) or w is a linear function (i.e., there exists $b \in \mathbb{R}^3$ such that $w(x) = b \cdot x$ a.e. in B_1).

Proof. We will first prove that the family $(w_r)_{r>0}$ is bounded in $L^1(B_4)$, and use the boundedness of the ACF functional to deduce the desired dichotomy.

Step 1. *There exists a constant C such that $\|w_r\|_{L^1(B_4)} \leq C$ for all r sufficiently large.*

We begin by noticing that, thanks to Lemma 5.2, there exist affine linear functions A_C^r and $A_{\gamma_\infty P}^r$ such that, for all r sufficiently large,

$$\int_{B_{4r}} |u - p - A_C^r| dx \leq Cr \quad \text{and} \quad \int_{B_{4r}} |u_\infty - p - A_{\gamma_\infty P}^r| dx \leq Cr$$

(note that, in the case $u_\infty = p$, the second estimate is trivially satisfied with $A_{\gamma_\infty P}^r \equiv 0$). Setting $\mathcal{A}^r(x) := \frac{1}{r}(A_C^r(rx) - A_{\gamma_\infty P}^r(rx))$, it follows by the triangle inequality that

$$\int_{B_4} |w_r - \mathcal{A}^r| dx \leq C. \quad (8.6)$$

Also, applying Lemma 5.3 to both u and u_∞ we deduce that $|\mathcal{A}^r(0)| + |\nabla' \mathcal{A}^r(0)| \leq C$, and therefore (8.6) implies that

$$\int_{B_4} |w_r(x) - \alpha_r x_3| dx \leq C \text{ for all } r \text{ sufficiently large, where } \alpha_r := \partial_3 \mathcal{A}^r(0). \quad (8.7)$$

In particular, by Chebyshev's inequality,

$$|B_4 \cap \{|w_r - \alpha_r x_3| > |\alpha_r|/2\}| \leq \frac{2C}{|\alpha_r|} \quad \text{for all } r \text{ sufficiently large.} \quad (8.8)$$

Suppose now towards a contradiction that the claim of this step is false. Then there exists a sequence $\varrho_k \rightarrow \infty$ as $k \rightarrow \infty$ such that $\|w_{\varrho_k}\|_{L^1(B_4)} \rightarrow \infty$. In particular, (8.7) implies that $|\alpha_{\varrho_k}| \rightarrow \infty$ as $k \rightarrow \infty$.

Without loss of generality, we may assume that $\alpha_{\varrho_k} \rightarrow \infty$. Then (8.7) and (8.8) imply that, for sufficiently large k ,

$$|\{(w_{\varrho_k})_- = 0\} \cap B_4| \geq |\{w_{\varrho_k} < 0\} \cap B_4| \geq \frac{1}{2}|B_4 \cap \{x_3 < -1/2\}|, \quad \|(w_{\varrho_k})_-\|_{L^1(B_4)} \rightarrow \infty,$$

$$|\{(w_{\varrho_k})_+ = 0\} \cap B_4| \geq |\{w_{\varrho_k} > 0\} \cap B_4| \geq \frac{1}{2}|B_4 \cap \{x_3 > 1/2\}|, \quad \|(w_{\varrho_k})_+\|_{L^1(B_4)} \rightarrow \infty.$$

This allows us to apply the Poincaré’s inequality to $(w_{\varrho_k})_{\pm}$ and obtain

$$\infty \leftarrow \|(w_{\varrho_k})_{\pm}\|_{L^1(B_4)} \leq C \|\nabla(w_{\varrho_k})_{\pm}\|_{L^1(B_4)}.$$

Since $\frac{1}{|x|} \geq \frac{1}{2}$ inside B_4 , it follows by Hölder’s inequality that

$$\|\nabla(w_{\varrho_k})_{\pm}\|_{L^1(B_4)}^2 \leq C \int_{B_4} |\nabla(w_{\varrho_k})_{\pm}|^2 dx \leq C \int_{B_4} \frac{|\nabla(w_{\varrho_k})_{\pm}|^2}{|x|} dx.$$

Thus, recalling that $N = 3$ and the definition of Φ , we conclude that

$$\infty \leftarrow \Phi(w_{\varrho_k}, 4) = \Phi(u - u_{\infty}, 4\varrho_k),$$

a contradiction to (8.4).

Step 2. *Proof of the dichotomy.*

Thanks to Step 1 and Lemma 2.12(ii), there exists $\bar{r} > 0$ such that

$$\int_{B_1} \frac{|\nabla(w_r)_{\pm}|^2}{|x|} dx \leq C \quad \text{for all } r > \bar{r}.$$

This implies the following non-concentration estimate for the ACF integrands: for each $\delta \in (0, 1)$ and every $r > \frac{\bar{r}}{\delta}$,

$$0 \leq \int_{B_{\delta}} \frac{|\nabla(w_r)_{\pm}|^2}{|x|} dx = \delta^2 \int_{B_1} \frac{|\nabla(w_{r\delta})_{\pm}|^2}{|x|} dx \leq C\delta^2. \tag{8.9}$$

In addition, since $\frac{1}{|x|} \geq 1$ inside B_1 , we have that $\|\nabla w_r\|_{L^2(B_1)} \leq C$ for all $r \geq \bar{r}$. Since $\|w_r\|_{L^1(B_2)} \leq C$ (from Step 1), there exists a sequence $r_k \rightarrow \infty$ such that

$$w_{r_k} \rightharpoonup w \quad \text{weakly in } W^{1,2}(B_1) \text{ as } k \rightarrow \infty.$$

We now observe that the coincidence sets of u and u_{∞} satisfy the properties in Proposition 4.1, therefore $\frac{1}{r}(\{u = 0\} \cup \{u_{\infty} = 0\}) \rightarrow \{se^3 : s \geq 0\}$ as $r \rightarrow \infty$. Hence, since $\Delta(u - u_{\infty}) = 0$ outside $\{u = 0\} \cup \{u_{\infty} = 0\}$, we deduce that $\Delta w = 0$ outside $\{se^3 : s \geq 0\}$. Because this set has zero harmonic capacity in \mathbb{R}^3 , we conclude that $\Delta w \equiv 0$ and therefore, by Lemma 2.15,

$$w_{r_k} \rightarrow w \quad \text{strongly in } W_{\text{loc}}^{1,2}(B_1) \text{ as } k \rightarrow \infty. \tag{8.10}$$

Combining the strong convergence (8.10) with the non-concentration estimate (8.9) we conclude that, for each $\varrho \in (0, 1)$,

$$\limsup_{k \rightarrow \infty} \left| \int_{B_{\varrho}} \frac{|\nabla(w_{r_k})_{\pm}|^2}{|x|} dx - \int_{B_{\varrho}} \frac{|\nabla w_{\pm}|^2}{|x|} dx \right| \leq \limsup_{k \rightarrow \infty} \left| \int_{B_{\delta}} \frac{|\nabla(w_{r_k})_{\pm}|^2}{|x|} dx - \int_{B_{\delta}} \frac{|\nabla w_{\pm}|^2}{|x|} dx \right| \leq C\delta^2,$$

so, by the arbitrariness of δ ,

$$\int_{B_{\varrho}} \frac{|\nabla(w_{r_k})_{\pm}|^2}{|x|} dx \rightarrow \int_{B_{\varrho}} \frac{|\nabla w_{\pm}|^2}{|x|} dx \quad \text{as } k \rightarrow \infty.$$

In particular $\Phi(w_{r_k}, \varrho) \rightarrow \Phi(w, \varrho)$ as $k \rightarrow \infty$ and therefore, by the monotonicity of the ACF functional as well as (8.4),

$$\infty > \Phi(u - u_{\infty}, \infty) = \lim_{k \rightarrow \infty} \Phi(u - u_{\infty}, r_k \varrho) = \lim_{k \rightarrow \infty} \Phi(w_{r_k}, \varrho) = \Phi(w, \varrho) \quad \text{for each } \varrho \in (0, 1).$$

This proves that $\varrho \mapsto \Phi(w, \varrho)$ is constant on $(0, 1)$. Since w is harmonic in B_1 , the result follows from [17, Theorem 2.9]. \square

As we shall see later, if we are in Case (i) of Proposition 8.1, then it is easy to conclude. On the other hand, Case (ii) requires a delicate argument that is performed in the next section.

8.2. Fine adjustment of u_∞ at large scales. The goal of this section is to show that if Case (ii) of Proposition 8.1 occurs, then we can find some fine adjustments of u_∞ at large scales to cancel the linear function $b \cdot x$ appearing in the blow-down limit.

Proposition 8.2 (Fine adjustment of the matching). *Let u_∞ , γ_∞ , $\bar{\gamma}$, and P be as in (8.2)-(8.3) and Corollary 6.5, and assume that*

$$\frac{(u - u_{\gamma_\infty P})(r_k x)}{r_k} \rightarrow b \cdot x \quad \text{strongly in } W^{1,2}(B_1), \quad \text{where } r_k \rightarrow \infty \text{ as } k \rightarrow \infty. \quad (8.11)$$

Then there exist $\tau' \in \mathbb{R}^2$ and a sequence $\gamma_k \rightarrow \gamma_\infty$, with $\gamma_k \in [0, \bar{\gamma} + 1]$, such that, for each $\sigma \in \mathbb{R}$,

$$\frac{(u - u_{\gamma_k P - (\tau', \sigma)})(r_k x)}{r_k} \rightarrow 0 \quad \text{strongly in } L^1(B_1) \text{ as } k \rightarrow \infty. \quad (8.12)$$

To prove this result, we will need a series of preliminary estimates on the behaviour of paraboloid solutions under translation and scaling. We collect these in the lemmas below.

Lemma 8.3 (Translations of paraboloid solutions). *Let P be as in (8.3). Then there exists a constant $C = C(P, \bar{\gamma})$ such that, for every $\gamma \in [0, \bar{\gamma} + 1]$ and every $\tau \in \mathbb{R}^3$,*

$$\int_{B_R} |V_{\gamma P}(x + \tau) - V_{\gamma P}(x)| dx \leq C|\tau|R^{3/4} \quad \text{for all } R \geq \max\{|\tau|, 1\}.$$

Proof. Using the fundamental theorem of calculus, for any $R \geq |\tau|$ we can estimate

$$\int_{B_R} |V_{\gamma P}(x + \tau) - V_{\gamma P}(x)| dx \leq C|\tau| \int_{B_{2R}} |\nabla V_{\gamma P}(x)| dx.$$

Applying Hölder's inequality and Lemma 2.13(ii) (note that $V_{\gamma P} = u_{\gamma P} - p$, where both $u_{\gamma P}$ and p are solutions to the obstacle problem) to the right-hand side above, we obtain

$$\int_{B_R} |V_{\gamma P}(x + \tau) - V_{\gamma P}(x)| dx \leq C|\tau| \sqrt{\int_{B_{2R}} |\nabla V_{\gamma P}(x)|^2 dx} \leq C \frac{|\tau|}{R} \sqrt{\int_{B_{4R}} |V_{\gamma P}(x)|^2 dx}.$$

Recall now that, thanks to (3.8), there exists a constant $C_{\bar{\gamma}}$ such that $|V_{\gamma P}(x)| \leq C_{\bar{\gamma}}(1 + |x|)^{7/4}$ for all $\gamma \in [0, \bar{\gamma} + 1]$. Combining these facts, we conclude that

$$\int_{B_R} |V_{\gamma P}(x + \tau) - V_{\gamma P}(x)| dx \leq C|\tau|R^{3/4} \quad \text{for all } R \geq \max\{|\tau|, 1\}.$$

\square

Lemma 8.4 (The generalized Newtonian potential of scaled paraboloids). *Let u_∞ , γ_∞ , $\bar{\gamma}$, and P be as in (8.2)-(8.3) and Corollary 6.5, and for $\gamma \in [0, \bar{\gamma} + 1]$ and $R > 0$ define the affine function*

$$\begin{aligned} \mathcal{A}_{(\gamma P, \gamma_\infty P)}^R(x) &:= A_{\gamma P}^R(x) - A_{\gamma_\infty P}^R(x) \\ &= \alpha_3 \int_{\mathbb{R}^3} \left(-\frac{1}{|y|} + \frac{1}{|Re^3 + y|} - \frac{x \cdot y}{|y|^3} + \frac{(Re^3 + y) \cdot (x + Re^3)}{|Re^3 + y|^3} \right) (\chi_{\gamma P} - \chi_{\gamma_\infty P})(y) \, dy. \end{aligned}$$

Then there exists a modulus of continuity $\omega = \omega_{P, \bar{\gamma}} : [0, \infty) \rightarrow [0, \infty)$ such that $\omega(0) = 0$ and, for all $R \geq 1$,

$$\int_{B_R} |V_{\gamma P} - V_{\gamma_\infty P} - \mathcal{A}_{(\gamma P, \gamma_\infty P)}^R| \, dx \leq \omega(|\gamma - \gamma_\infty|)R, \quad (8.13)$$

$$|\nabla' \mathcal{A}_{(\gamma P, \gamma_\infty P)}^R| \leq \omega(|\gamma - \gamma_\infty|) \quad \text{and} \quad |\mathcal{A}_{(\gamma P, \gamma_\infty P)}^R(0)| \leq \omega(|\gamma - \gamma_\infty|)R. \quad (8.14)$$

Proof. As we shall see, the proof of is slight modification of the ones of Lemmas 5.2 and 5.3. We recall that $A \Delta B$ denotes the symmetric difference of two sets A and B .

Step 1. *Proof of (8.13).*

We follow the notation used in the proof of Lemma 5.2. Recalling Definition 3.1 and (5.6), we have

$$\begin{aligned} |V_{\gamma P}(x) - V_{\gamma_\infty P}(x) - \mathcal{A}_{(\gamma P, \gamma_\infty P)}^R(x)| &\leq \alpha_3 \int_{(\gamma P \Delta \gamma_\infty P) \cap \{0 \leq y_3 \leq 2R\}} |G(x, y) - a^R(x, y)| \, dy \\ &\quad + \alpha_3 \int_{(\gamma P \Delta \gamma_\infty P) \cap \{y_3 \geq 2R\}} |G(x, y) - a^R(x, y)| \, dy =: J_1(x) + J_2(x). \end{aligned}$$

Using (3.2) and (5.7), for $x \in B_R$ we estimate

$$\begin{aligned} J_2(x) &\leq CR^2 \int_{2R}^\infty \mathcal{H}^2 \left(\left\{ y' \in \sqrt{t} (\sqrt{\gamma} E' \Delta \sqrt{\gamma_\infty} E') \right\} \right) \frac{1}{t^3} \, dt \\ &= CR^2 \mathcal{H}^2(E') |\gamma - \gamma_\infty| \int_{2R}^\infty \frac{1}{t^2} \, dt \leq C |\gamma - \gamma_\infty| R \end{aligned}$$

(cp. (5.11)). For $J_1(x)$, we write $x = Rz$ with $z \in B_1$ and we perform the change of variables $y \mapsto Ry$, so that

$$J_1(x) \leq CR^2 \int_{\left\{ y' \in \sqrt{\frac{y_3}{R}} (\sqrt{\gamma} E' \Delta \sqrt{\gamma_\infty} E') \right\} \cap \{0 \leq y_3 \leq 2\}} |G(z, y) - a^1(z, y)| \, dy$$

(cp. (5.3)). Combining these two bounds, we get

$$\begin{aligned} \int_{B_R} |V_{\gamma P}(x) - V_{\gamma_\infty P}(x) - \mathcal{A}_{(\gamma P, \gamma_\infty P)}^R(x)| \, dx &\leq \int_{B_R} (J_1(x) + J_2(x)) \, dx \\ &\leq C |\gamma - \gamma_\infty| R + CR^2 \int_{B_1} \int_{\left\{ y' \in \sqrt{\frac{y_3}{R}} (\sqrt{\gamma} E' \Delta \sqrt{\gamma_\infty} E') \right\} \cap \{0 \leq y_3 \leq 2\}} |G(z, y) - a^1(z, y)| \, dy \, dz \end{aligned}$$

(cp. (5.12)). Concerning the last integral we note that (5.13) holds on the domain of integration. Hence, since $\left| \left\{ y' \in \sqrt{\frac{y_3}{R}} (\sqrt{\gamma} E' \Delta \sqrt{\gamma_\infty} E') \right\} \cap \{0 \leq y_3 \leq 2\} \right| \leq \frac{C|\gamma - \gamma_\infty|}{R}$, using (5.4) and Fubini's Theorem we obtain

$$\int_{B_R} |V_{\gamma P} - V_{\gamma_\infty P} - \mathcal{A}_{(\gamma P, \gamma_\infty P)}^R| \, dx \leq C |\gamma - \gamma_\infty| R,$$

(cp. (5.14)). This proves (8.13).

Step 2. *Proof of (8.14).*

For the first bound we note that, for $y \in \gamma P \cup \gamma_\infty P$, (5.15) holds. Hence

$$\begin{aligned} \int_{(\gamma P \Delta \gamma_\infty P)} |\nabla_{x'} a^R(x, y)| dy &\leq 2 \int_{(\gamma P \Delta \gamma_\infty P) \cap \{y_3 \leq 1\}} \frac{1}{|y|^2} dy + 2 \int_{(\gamma P \Delta \gamma_\infty P) \cap \{y_3 \geq 1\}} \frac{|y'|}{|y|^3} dy \\ &\leq 2 \int_{(\gamma P \Delta \gamma_\infty P) \cap \{y_3 \leq 1\}} \frac{1}{|y|^2} dy \\ &\quad + C \int_1^\infty \mathcal{H}^2 \left(\left\{ y' \in \sqrt{t} (\sqrt{\gamma} E' \Delta \sqrt{\gamma_\infty} E') \right\} \right) \frac{t^{1/2}}{t^3} dt \\ &\leq 2 \int_{(\gamma P \Delta \gamma_\infty P) \cap \{y_3 \leq 1\}} \frac{1}{|y|^2} dy + C |\gamma - \gamma_\infty| \int_1^\infty \frac{1}{t^{3/2}} dt. \end{aligned}$$

By dominated convergence, this proves the existence of a modulus of continuity ω such that $|\nabla' \mathcal{A}_{(\gamma P, \gamma_\infty P)}^R| \leq \omega(|\gamma - \gamma_\infty|)$.

For the second bound we write

$$\begin{aligned} \int_{(\gamma P \Delta \gamma_\infty P)} |a^R(0, y)| dy &\leq \int_{(\gamma P \Delta \gamma_\infty P) \cap \{y_3 \leq 2R\}} |a^R(0, y)| dy \\ &\quad + \int_{(\gamma P \Delta \gamma_\infty P) \cap \{y_3 \geq 2R\}} |a^R(0, y)| dy =: I_1 + I_2. \end{aligned}$$

Using (5.7), we immediately get

$$\begin{aligned} I_2 &\leq CR^2 \int_{2R}^\infty \mathcal{H}^2 \left(\left\{ y' \in \sqrt{t} (\sqrt{\gamma} E' \Delta \sqrt{\gamma_\infty} E') \right\} \right) \frac{1}{t^3} dt \\ &\leq CR^2 |\gamma - \gamma_\infty| \int_{2R}^\infty \frac{1}{t^2} dt \leq C |\gamma - \gamma_\infty| R, \end{aligned}$$

(cp. (5.8)). Concerning I_1 , using (5.16) we have

$$I_1 \leq 3 \int_0^R \mathcal{H}^2 \left(\left\{ y' \in \sqrt{t} (\sqrt{\gamma} E' \Delta \sqrt{\gamma_\infty} E') \right\} \right) \frac{1}{t} dt \leq C |\gamma - \gamma_\infty| \int_0^R dt \leq C |\gamma - \gamma_\infty| R$$

(cp. (5.17)). This implies that $|\mathcal{A}_{(\gamma P, \gamma_\infty P)}^R(0)| \leq C |\gamma - \gamma_\infty| R$, concluding the proof. \square

Lemma 8.5. *Let P be as in (8.3), and fix $\gamma \in [0, \bar{\gamma} + 1]$. Then there exists a constant $C = C(P, \bar{\gamma})$ such that for all $R \geq 1$,*

$$\left| \frac{\partial}{\partial R} \partial_3 \mathcal{A}_{\gamma P}^R \right| \leq \frac{C}{R}.$$

Proof. Recalling (5.10), it follows that

$$\begin{aligned} \left| \frac{\partial}{\partial R} \partial_3 \mathcal{A}_{\gamma P}^R \right| &\leq \alpha_3 \int_{\gamma P} \left| \frac{\partial}{\partial R} \left(-\frac{y_3}{|y|^3} + \frac{R + y_3}{|Re^3 + y|^3} \right) \right| dy = \alpha_3 \int_{\gamma P} \left| \frac{\partial}{\partial R} \left(\frac{R + y_3}{|Re^3 + y|^3} \right) \right| dy \\ &\leq C \int_{\gamma P} \frac{1}{|Re^3 + y|^3} dy \leq C \int_0^\infty \mathcal{H}^2 \left(\left\{ y' \in \sqrt{\gamma t} E' \right\} \right) \frac{1}{(R + t)^3} dt \end{aligned}$$

$$\leq C \int_0^\infty \frac{t}{(R+t)^3} dt \leq \frac{C}{R^3} \int_0^R t dt + C \int_R^\infty \frac{1}{t^2} dt \leq \frac{C}{R}.$$

□

Proof of Proposition 8.2. We note that, given $\gamma \in [0, \bar{\gamma} + 1]$, we have (cp. (6.2))

$$\begin{aligned} u_{\gamma P - (\tau', \sigma)}(x) &= u_{\gamma P}(x' + \tau', x_3 + \sigma) = p(x' + \tau') + V_{\gamma P}(x' + \tau', x_3 + \sigma) \\ &= p(x') + \nabla' p(x') \cdot \tau' + p(\tau') + \left(V_{\gamma P}(x' + \tau', x_3 + \sigma) - V_{\gamma P}(x) \right) \\ &\quad + \left(V_{\gamma P}(x) - V_{\gamma_\infty P}(x) \right) + V_{\gamma_\infty P}(x) \\ &= u_{\gamma_\infty P}(x) + \nabla' p(x') \cdot \tau' + \left(V_{\gamma P}(x) - V_{\gamma_\infty P}(x) \right) \\ &\quad + \left[p(\tau') + \left(V_{\gamma P}(x' + \tau', x_3 + \sigma) - V_{\gamma P}(x) \right) \right]. \end{aligned}$$

Let $b = (b', b_3) \in \mathbb{R}^2 \times \mathbb{R}$. Recalling that $\nabla' p(x') = Qx'$ with Q symmetric and invertible, we choose $\tau' := Q^{-1}b'$ so that $\nabla' p(x') \cdot \tau' = b' \cdot x'$. Also, for each $R > 1$,

$$\begin{aligned} V_{\gamma P}(x) - V_{\gamma_\infty P}(x) &= \left(V_{\gamma P}(x) - V_{\gamma_\infty P}(x) - \mathcal{A}_{(\gamma P, \gamma_\infty P)}^R(x) \right) \\ &\quad + \mathcal{A}_{(\gamma P, \gamma_\infty P)}^R(0) + \nabla' \mathcal{A}_{(\gamma P, \gamma_\infty P)}^R \cdot x' + \partial_3 \mathcal{A}_{(\gamma P, \gamma_\infty P)}^R x_3. \end{aligned}$$

Combining all these identities and applying Lemmas 8.3 and 8.4, we obtain the existence of a modulus of continuity $\omega : [0, \infty) \rightarrow [0, \infty)$ such that, for $x \in B_R$,

$$\begin{aligned} \int_{B_R} \left| u_{\gamma P - (\tau', \sigma)}(x) - u_{\gamma_\infty P}(x) - b' \cdot x' - \partial_3 \mathcal{A}_{(\gamma P, \gamma_\infty P)}^R x_3 \right| dx \\ \leq C \left(\omega(|\gamma - \gamma_\infty|)R + R^{3/4} \right). \end{aligned} \quad (8.15)$$

Note that, without loss of generality, we can assume that $\omega(t) \geq t$ (otherwise, it suffices to replace $\omega(t)$ with $\omega(t) + t$).

We now distinguish two cases, depending on whether $\gamma_\infty > 0$ or not.

• **Case 1:** $\gamma_\infty > 0$.

We note that, by a change of variables, for each $\lambda > 0$ it holds

$$\begin{aligned} \partial_3 A_{\lambda \gamma_\infty P}^R &= \int_{\lambda \gamma_\infty P} \left(-\frac{y_3}{|y|^3} + \frac{R + y_3}{|Re^3 + y|^3} \right) dy = \lambda^3 \int_{\gamma_\infty P} \left(-\frac{\lambda z_3}{\lambda^3 |z|^3} + \frac{R + \lambda z_3}{|Re^3 + \lambda z|^3} \right) dz \\ &= \lambda \int_{\gamma_\infty P} \left(-\frac{z_3}{|z|^3} + \frac{\frac{R}{\lambda} + z_3}{\left| \frac{R}{\lambda} e^3 + z \right|^3} \right) dz = \lambda \partial_3 A_{\gamma_\infty P}^{R/\lambda}. \end{aligned} \quad (8.16)$$

Thus, setting $\lambda := \frac{\gamma}{\gamma_\infty}$,

$$\partial_3 A_{(\gamma P, \gamma_\infty P)}^R = \partial_3 A_{\gamma P}^R - \partial_3 A_{\gamma_\infty P}^R = \lambda \left[\partial_3 A_{\gamma_\infty P}^{R/\lambda} - \partial_3 A_{\gamma_\infty P}^R \right] + (\lambda - 1) \partial_3 A_{\gamma_\infty P}^R. \quad (8.17)$$

Assuming now that γ is sufficiently close to γ_∞ so that $\lambda \in [1/2, 2]$, it follows from Lemma 8.5 that

$$\left| \partial_3 A_{\gamma_\infty P}^{R/\lambda} - \partial_3 A_{\gamma_\infty P}^R \right| \leq \int_{R/\lambda}^R \left| \frac{\partial}{\partial r} \partial_3 A_{\gamma P}^r \right| dr \leq C |\lambda - 1| = C \frac{|\gamma - \gamma_\infty|}{|\gamma_\infty|}.$$

Thus, combining (8.15) and (8.17), we obtain

$$\int_{B_R} \left| u_{\gamma P - (\tau', \sigma)}(x) - u_{\gamma_\infty P}(x) - b' \cdot x' - \frac{\gamma - \gamma_\infty}{\gamma_\infty} \partial_3 A_{\gamma_\infty P}^R x_3 \right| dx \leq C \left(\omega(|\gamma - \gamma_\infty|)R + R^{3/4} \right).$$

We now observe that, as a consequence of Lemma 5.3 and (6.7), there exist constants $0 < c_\infty < C_\infty$ such that

$$-c_\infty \log R \geq \partial_3 A_{\gamma_\infty P}^R \geq -C_\infty \log R \quad \text{for all } R \text{ sufficiently large.}$$

In particular, for each R sufficiently large we can choose $\gamma = \gamma_R \in \left[\gamma_\infty - \frac{|b_3| \gamma_\infty}{c_\infty \log R}, \gamma_\infty + \frac{|b_3| \gamma_\infty}{c_\infty \log R} \right]$ such that $\frac{\gamma_R - \gamma_\infty}{\gamma_\infty} \partial_3 A_{\gamma_\infty P}^R = b_3$, and with such a choice we have

$$\int_{B_R} \left| u_{\gamma_R P - (\tau', \sigma)}(x) - u_{\gamma_\infty P}(x) - b' \cdot x' - b_3 x_3 \right| dx \leq C \left(\omega \left(\frac{C}{\log R} \right) R + R^{3/4} \right) \quad (8.18)$$

for all R sufficiently large. Choosing $R = r_k$ and $\gamma_k = \gamma_{r_k}$, and combining (8.11) and (8.18), we obtain (8.12).

• **Case 2:** $\gamma_\infty = 0$.

Note that in this case $V_{\gamma_\infty P} \equiv 0$. We first claim that $b_3 \leq 0$. Indeed, from Definition 2.8(ii) we know that $u - p$ is harmonic in $\{y_3 \leq 0\}$, which combined with (8.11) implies that

$$\frac{(u - p)(r_k x)}{r_k} \rightarrow b \cdot x \quad \text{uniformly in } B_{1/2}(-e^3) \text{ as } k \rightarrow \infty.$$

On the other hand, Definition 2.8(ii)-(iv) implies that $(u - p)(-te^3) \geq 0$ for all $t \geq 0$. Thus

$$0 \leq \frac{(u - p)(-r_k e^3)}{r_k} \rightarrow -b_3 \quad \text{as } k \rightarrow \infty,$$

proving the claim.

Now, if $b_3 = 0$ then the result follows by choosing $\gamma = 0$. Otherwise, applying (8.16) with $\gamma > 0$ in place of λ and P in place of $\gamma_\infty P$, we get

$$\partial_3 \mathcal{A}_{(\gamma P, \gamma_\infty P)}^R = \partial_3 A_{\gamma P}^R = \gamma \partial_3 A_P^{R/\gamma}.$$

Moreover, for each $\gamma \in (0, 1]$, we know from Lemma 5.3 and (6.7) that

$$-c_\infty \log(R/\gamma) \geq \partial_3 A_P^{R/\gamma} \geq -C_\infty \log(R/\gamma) \quad \text{for all } R \text{ sufficiently large.}$$

Thus, recalling that $b_3 < 0$, for each R sufficiently large we can find $\gamma = \gamma_R \in \left(0, \frac{2|b_3|}{c_\infty \log R} \right]$ such that $\gamma_R \partial_3 A_P^{R/\gamma_R} = b_3$. Choosing $R = r_k$ and $\gamma_k = \gamma_{r_k}$, we conclude as in Case 1. \square

8.3. Ordering of solutions and conclusion.

Proposition 8.6 (Ordering in dimension $N = 3$). *Let u_∞ , γ_∞ , and P be as in (8.2)-(8.3). Then there exists $\tau' \in \mathbb{R}^2$ such that for each $\sigma \in \mathbb{R}$,*

$$\text{either } u \leq u_{\gamma_\infty P - (\tau', \sigma)} \text{ in } \mathbb{R}^N \quad \text{or} \quad u \geq u_{\gamma_\infty P - (\tau', \sigma)} \text{ in } \mathbb{R}^N.$$

Proof. Let w_{r_k} and w be as in Proposition 8.1. We distinguish between the two cases in the dichotomy.

• **Case 1:** Proposition 8.1(i) holds.

In this case we have that either $\|(w_{r_k})_+\|_{L^1(B_1)} \rightarrow 0$ or $\|(w_{r_k})_-\|_{L^1(B_1)} \rightarrow 0$ as $k \rightarrow \infty$. Also, since

$$u_{\gamma_\infty P}(x) - u_{\gamma_\infty P-(0,\sigma)}(x) = u_{\gamma_\infty P}(x) - u_{\gamma_\infty P}(x + \sigma e^3) = V_{\gamma_\infty P}(x) - V_{\gamma_\infty P}(x + \sigma e^3),$$

defining $w_{k,\sigma}(x) := \frac{(u - u_{\gamma_\infty P-(0,\sigma)})(r_k x)}{r_k}$, Lemma 8.3 yields

$$\|w_{r_k} - w_{k,\sigma}\|_{L^1(B_1)} = \frac{|B_1|}{r_k} \int_{B_{r_k}} |V_{\gamma_\infty P}(x) - V_{\gamma_\infty P}(x + \sigma e^3)| dx \leq C \frac{|\sigma|}{r_k^{1/4}} \rightarrow 0 \quad \text{as } k \rightarrow \infty.$$

Hence, for each $\sigma \in \mathbb{R}$, either $\|(w_{k,\sigma})_+\|_{L^1(B_1)} \rightarrow 0$ or $\|(w_{k,\sigma})_-\|_{L^1(B_1)} \rightarrow 0$ as $k \rightarrow \infty$. Therefore, thanks to Lemma 2.12(i)-(iii), for each $\varrho \in (0, \infty)$ it holds

$$\begin{aligned} 0 \leq \Phi(u - u_\infty(\cdot + \sigma e^3), \varrho) &\leq \lim_{k \rightarrow \infty} \Phi(u - u_\infty(\cdot + \sigma e^3), \frac{r_k}{4}) \\ &= \lim_{k \rightarrow \infty} \Phi(w_{k,\sigma}, \frac{1}{4}) \leq C \limsup_{k \rightarrow \infty} \|(w_{k,\sigma})_+\|_{L^1(B_1)}^2 \|(w_{k,\sigma})_-\|_{L^1(B_1)}^2 = 0. \end{aligned}$$

Applying Lemma 2.12(iv) proves the result with $\tau' = 0$.

• **Case 2:** Proposition 8.1(ii) holds.

Let γ_k and τ' be as in Proposition 8.2, and define $w'_{k,\sigma}(x) := \frac{(u - u_{\gamma_k P-(\tau',\sigma)})(r_k x)}{r_k}$. Note that, since $\gamma_k \rightarrow \gamma_\infty$ and solutions to the obstacle problems are locally bounded in $C^{1,1}$ (cf. Lemma 2.4(i)),

$$u_{\gamma_k P-(\tau',\sigma)} \rightarrow u_{\gamma_\infty P-(\tau',\sigma)} = u_\infty(\cdot + (\tau', \sigma)) \quad \text{in } C_{\text{loc}}^{1,\alpha}(\mathbb{R}^3) \quad \text{as } k \rightarrow \infty,$$

which implies in particular that, given $\varrho \in (0, \infty)$,

$$\Phi(u - u_{\gamma_k P-(\tau',\sigma)}, \varrho) \rightarrow \Phi(u - u_\infty(\cdot + (\tau', \sigma)), \varrho) \quad \text{as } k \rightarrow \infty.$$

Since $\|w'_{k,\sigma}\|_{L^1(B_1)} \rightarrow 0$, it follows from Lemma 2.12(i)-(iii) that, for each $\varrho \in (0, \infty)$,

$$\begin{aligned} 0 \leq \Phi(u - u_\infty(\cdot + (\tau', \sigma)), \varrho) &= \lim_{k \rightarrow \infty} \Phi(u - u_{\gamma_k P-(\tau',\sigma)}, \varrho) \leq \limsup_{k \rightarrow \infty} \Phi(u - u_{\gamma_k P-(\tau',\sigma)}, \frac{r_k}{4}) \\ &= \limsup_{k \rightarrow \infty} \Phi(w'_{k,\sigma}, \frac{1}{4}) \leq C \limsup_{k \rightarrow \infty} \|(w'_{k,\sigma})_+\|_{L^1(B_1)}^2 \|(w'_{k,\sigma})_-\|_{L^1(B_1)}^2 = 0. \end{aligned}$$

Hence, the result follows again from Lemma 2.12(iv). \square

We can now prove our main result.

Proof of Theorem 1.2: the case $N = 3$. The proof is almost identical to the one of Theorem 1.2 for $N \geq 4$ given in Section 7, the only difference being the application of Proposition 8.6 instead of Proposition 7.1. \square

9. PROOF OF THEOREM 1.1

As explained after the statement of [7, Main Theorem**], every non-cylindrical solution with unbounded coincidence set is x_N -monotone. Hence, to prove Theorem 1.1 for $N \geq 3$, it suffices to characterize x_N -monotone solutions for $N \geq 3$, which is exactly the result of Theorem 1.2. For completeness and convenience of the interested reader, we present here an original alternative argument to show how Theorem 1.1 follows from Theorem 1.2.

Proof of Theorem 1.1. Let u be a global solution in \mathbb{R}^N with non-empty coincidence set.

We can assume that u is non-cylindrical (see Definition 2.6). Indeed, otherwise, up to a change of coordinates $u(y, z) = v(y)$ for $(z, y) \in \mathbb{R}^{N-k} \times \mathbb{R}^k$, where v is a non-cylindrical global solution in \mathbb{R}^k , and then it suffices to prove Theorem 1.1 for v . Also, as mentioned in the introduction, Theorem 1.1 has already been proved for $N = 2$ in [18].

Hence, we assume that u is a non-cylindrical global solution in dimension $N \geq 3$, whose coincidence set $\mathcal{C} := \{u = 0\}$ has non-empty interior (recall that \mathcal{C} is convex, see Remark 2.2). Then, we set $u_r(x) := \frac{u(rx)}{r^2}$ and define $g(x) := \lim_{r \rightarrow \infty} u_r(x)$ (cf. Lemma 2.3). We distinguish several cases.

• **Case 1:** g is a half-space solution.

By a translation and a rotation we can assume that $\{u = 0\} \subset \{x_N \leq 0\}$ and that $0 \in \partial\{u = 0\}$. Then, since $\{u = 0\}$ is convex we deduce that $\{u_r = 0\} = \frac{1}{r}\{u = 0\} \subset \{u = 0\}$ for every $r \geq 1$, and letting $r \rightarrow \infty$ we conclude that $\{g = 0\} \subset \{u = 0\}$.

On the other hand, since g is a half-space solution, $\{g = 0\}$ is a half-space passing through the origin. So the only option is that $\{g = 0\} = \{u = 0\} = \{x_N \leq 0\}$.

Thanks to this fact, we deduce that $\Delta(u - g) \equiv 0$ and $(u - g)|_{\{x_N \leq 0\}} = 0$. By unique continuation this implies that $u \equiv g$, and since g is constant in the directions orthogonal to e^N we obtain that u is cylindrical, a contradiction.

• **Case 2:** $g(x) = \frac{1}{2}x^T Q x$ is a quadratic polynomial solution with Q positive definite.

Since Q is positive definite, there exists a constant $c_0 > 0$ such that $g \geq c_0$ on ∂B_1 . Hence, it follows from the local uniform convergence of u_r to g that $u(x) \geq \frac{c_0}{2}|x|^2$ for sufficiently large $|x|$. This implies that the coincidence set $\{u = 0\}$ is compact, so the result follows from [10] (see also [9] or Remark 7.2).

• **Case 3:** $g(x) = \frac{1}{2}x^T Q x$ is a quadratic polynomial solution with $\ker(Q) \neq \{0\}$.

Step 1. u is monotone in the directions of $\ker(Q)$.

Let $e \in \partial B_1 \cap \ker(Q)$. Then $\partial_e u = \partial_e(u - g)$. Also, if we define $v_r(x) := \frac{(u-g)(rx)}{r^2} = u_r(x) - g(x)$, Hölder's inequality and Lemma 2.13(ii) imply that

$$\left(\int_{B_1} |\nabla v_r| dx \right)^2 \leq \int_{B_1} |\nabla v_r|^2 dx \leq C \int_{B_2} v_r^2 dx \rightarrow 0 \text{ as } r \rightarrow \infty.$$

It follows that

$$\frac{1}{r} \int_{B_r} |\partial_e u| dx = \frac{1}{r} \int_{B_r} |\partial_e(u - g)| dx = \int_{B_1} |\partial_e v_r| dx \rightarrow 0 \quad \text{as } r \rightarrow \infty. \quad (9.1)$$

Thus, thanks to Remark 2.14, (9.1), and Lemma 2.12(iii)-(iv), we deduce that either $\partial_e u \geq 0$ or $\partial_e u \leq 0$. Since $\partial_e u \not\equiv 0$ (as u is non-cylindrical) and $\partial_e u$ is harmonic outside the coincidence set of u , it follows by the strong maximum principle that

$$\text{either } \partial_e u > 0 \text{ or } \partial_e u < 0 \quad \text{inside } \{u > 0\}. \quad (9.2)$$

Step 2. $\ker(Q)$ is one-dimensional.

Indeed, assume by contradiction that there exists a two-dimensional plane $\Pi \subset \ker(Q)$. Then, by Step 1,

$$\text{for any } e \in \partial B_1 \cap \Pi, \quad (9.2) \text{ holds.}$$

Since $\partial B_1 \cap \Pi$ is a circle, given a point $\bar{x} \in \{u > 0\}$ we can consider a curve $[0, 1] \ni s \mapsto e(s) \subset \partial B_1 \cap \Pi$ such that $e(0) = -e(1)$. Then, noticing that $\partial_{e(0)} u(\bar{x}) = -\partial_{e(1)} u(\bar{x})$, it follows by continuity that there exists $s \in (0, 1)$ such that $\partial_{e(s)} u(\bar{x}) = 0$. This contradicts (9.2) and

proves that $\ker(Q)$ is one-dimensional.

Step 3. u is x_N -monotone.

Thanks to Steps 1 and 2, we may assume that $\ker(Q) = \mathbb{R}e^N$ and that $\partial_N u \geq 0$. Thus, since by assumption $\mathcal{C} = \{u = 0\}$ has non-empty interior, u satisfies Definition 2.8(i)-(iii)-(iv). Also, up to a translation, we can assume that $\mathcal{C} \subset \{x_N \geq 0\}$ and that $0 \in \partial\mathcal{C}$.

To conclude that u is x_N -monotone, we only need to show that $\mathcal{C} \cap \{x_N = 0\} = \{0\}$. This follows from [2, 15]. Indeed, since \mathcal{C} is convex with non-empty interior, [2, Section 1.8] (see also [3, Theorem 7]) implies $\partial\mathcal{C}$ is of class C^1 , and therefore it follows from [15] that $\partial\mathcal{C}$ is an analytic hypersurface. In particular $\partial\mathcal{C}$ cannot contain a segment, and therefore $\mathcal{C} \cap \{x_N = 0\}$ is a singleton. \square

APPENDIX A. PROOF OF LEMMA 6.1

We split the proof into three steps.

Step 1. *Construction of a suitable sequence of ellipsoids.*

Let K_d denote the fundamental solution of the Laplace operator in \mathbb{R}^d , namely,

$$K_d(z) := \begin{cases} -\frac{1}{2\pi} \log |z| & \text{if } d = 2, \\ \frac{1}{d(d-2)|B_1|} \frac{1}{|z|^{d-2}} & \text{if } d \geq 3. \end{cases} \quad (\text{A.1})$$

Given p as in the statement, it follows from [5, Equation (5.4)] that there exists a unique ellipsoid $E' := \{y' \in \mathbb{R}^{N-1} : \frac{y_1^2}{a_1^2} + \dots + \frac{y_{N-1}^2}{a_{N-1}^2} \leq 1\} \subset \mathbb{R}^{N-1}$, with $a_i > 0$, such that

$$V_{E'}^{NP}(x') = V_{E'}^{NP}(0) - p(x') \quad \text{for all } x' \in E',$$

where $V_{E'}^{NP}$ denotes the $(N-1)$ -dimensional Newtonian potential of E' , i.e.

$$V_{E'}^{NP}(x') := \int_{E'} K_{N-1}(x' - y') dy' \quad \text{for all } x' \in \mathbb{R}^{N-1}.$$

Set

$$u'_{E'}(x') := p(x') - V_{E'}^{NP}(0) + V_{E'}^{NP}(x') \quad \text{for all } x' \in \mathbb{R}^{N-1}. \quad (\text{A.2})$$

Then, it follows from [4, Theorem II] that $u'_{E'}$ is a non-negative global solution to the obstacle problem in \mathbb{R}^{N-1} satisfying $\{u'_{E'} = 0\} = E'$.

We now complete E' to an N -dimensional ellipsoid approximating a paraboloid in the following way: for each $n \in \mathbb{N}$, set

$$\tilde{E}^n := \left\{ x \in \mathbb{R}^N : x' \in \sqrt{\frac{n}{2} - \frac{x_N^2}{2n}} E' \right\} = \left\{ x \in \mathbb{R}^N : \sum_{j=1}^{N-1} \frac{2x_j^2}{a_j^2 n} + \frac{x_N^2}{n^2} \leq 1 \right\}.$$

From [5, Equation (5.3)] we infer that, for each $n \in \mathbb{N}$, there is a homogeneous quadratic polynomial q^n such that $\Delta q^n = 1$ and

$$V_{\tilde{E}^n}^{NP}(x) = V_{\tilde{E}^n}^{NP}(0) - q^n(x) \quad \text{for all } x \in \tilde{E}^n, \quad (\text{A.3})$$

where

$$V_{\tilde{E}^n}^{NP}(x) := \int_{\tilde{E}^n} K_N(x - y) dy \quad \text{for all } x \in \mathbb{R}^N.$$

Let us now translate the ellipsoids \tilde{E}^n so that they all touch the origin:

$$\begin{aligned} E^n &:= \tilde{E}^n + ne^N = \left\{ x \in \mathbb{R}^N : \sum_{j=1}^{N-1} \frac{2x_j^2}{a_j^2 n} + \frac{(x_N - n)^2}{n^2} \leq 1 \right\} \\ &= \left\{ x \in \mathbb{R}^N : \sum_{j=1}^{N-1} \frac{2x_j^2}{a_j^2} + \frac{x_N^2}{n} \leq 2x_N \right\}. \end{aligned} \quad (\text{A.4})$$

Then, recalling (A.3), for all $x \in E^n$ we have

$$\begin{aligned} V_{E^n}^{N,P}(x) &= V_{\tilde{E}^n + ne^N}^{N,P}(x) = V_{\tilde{E}^n}^{N,P}(x - ne^N) \\ &= V_{\tilde{E}^n}^{N,P}(0) - q^n(x - ne^N) = V_{E^n}^{N,P}(ne^N) - q^n(x - ne^N). \end{aligned}$$

Step 2. *Switching to the obstacle problem and passing to the limit.*

For all $n \in \mathbb{N}$, we define

$$u_{E^n}(x) := q^n(x - ne^N) - V_{E^n}^{N,P}(ne^N) + V_{E^n}^{N,P}(x) \quad \text{for all } x \in \mathbb{R}^N.$$

As before, [4, Theorem II] guarantees that u_{E^n} is a non-negative global solution to the obstacle problem satisfying $\{u_{E^n} = 0\} = E^n$.

Since $0 \in E^n = \{u_{E^n} = 0\}$ we deduce from Lemma 2.4(ii) that, passing if necessary to a subsequence,

$$u_{E^n} \rightarrow u^* \quad \text{in } C_{\text{loc}}^{1,\alpha}(\mathbb{R}^N) \quad \text{and} \quad \chi_{\{u_{E^n}=0\}} \rightarrow \chi_{\{u^*=0\}} \quad \text{a.e.} \quad \text{as } n \rightarrow \infty.$$

On the other hand, by construction (cf. (A.4)),

$$\chi_{E^n} \rightarrow \chi_P \quad \text{a.e. in } \mathbb{R}^N \text{ as } n \rightarrow \infty, \quad \text{where } P := \left\{ \sum_{j=1}^{N-1} \frac{x_j^2}{a_j^2} \leq x_N \right\} = \{x' \in \sqrt{x_N} E'\},$$

and therefore $\{u^* = 0\} = P$.

Step 3. *Identification of the blow-down limit of u^* and conclusion.*

Let us define the following sequence of rescalings $(u_k^*)_{k \in \mathbb{N}}$:

$$u_k^*(x) := \frac{u^*(r_k x + x^k)}{r_k^2} \quad \text{for } x \in \mathbb{R}^N, \quad \text{where } x^k := (0, k) \text{ and } r_k := \sqrt{k}.$$

Since $0 \in \{u_k^* = 0\}$, using Lemma 2.4(ii) once more we deduce that, passing if necessary to a subsequence,

$$u_k^* \rightarrow u_0^* \quad \text{in } C_{\text{loc}}^{1,\alpha}(\mathbb{R}^N) \quad \text{as } k \rightarrow \infty,$$

where u_0^* is a non-negative global solution to the obstacle problem. Also, arguing as in Step 2, we see that the coincidence sets of u_k^* converge to $E' \times \mathbb{R}$, hence $\{u_0^* = 0\} = E' \times \mathbb{R}$. This implies that $\{u_0^* = 0\}$ contains the ray $\{te^N : t \in \mathbb{R}\}$, so it follows from Lemma 2.7 that u_0^* is independent of x_N , i.e.

$$u_0^*(x) = u_0^*(x', 0) =: u_0^{\star'}(x') \quad \text{for all } x \in \mathbb{R}^N. \quad (\text{A.5})$$

Since $u'_{E'}$ constructed in Step 1 (cf. (A.2)) is the unique global solution to the obstacle problem in \mathbb{R}^{N-1} with E' as coincidence set, we deduce that $u_0^{\star'} \equiv u'_{E'}$. Also, since the

classical Newton-potential $V_{E'}^{NP}(x')$ has subquadratic growth as $|x'| \rightarrow \infty$, we have that $p(x')$ is the blow-down limit of $u'_{E'}$. Thus, recalling (A.5),

$$\frac{u_0^{*\prime}(\varrho x)}{\varrho^2} = \frac{u'_{E'}(\varrho x')}{\varrho^2} \rightarrow p(x') \quad \text{in } C_{\text{loc}}^{1,\alpha}(\mathbb{R}^{N-1}) \text{ as } \varrho \rightarrow \infty.$$

On the other hand, since in the limit as $\varrho \rightarrow \infty$ the contact set of $\frac{u^*(\varrho x)}{\varrho^2}$ has measure zero (as a consequence of Lemma 2.4(ii)), the blow-down limit of u^* is a homogeneous quadratic polynomial \tilde{p} as well, i.e.

$$\frac{u^*(\varrho x)}{\varrho^2} \rightarrow \tilde{p}(x) \quad \text{in } C_{\text{loc}}^{1,\alpha}(\mathbb{R}^N) \text{ as } \varrho \rightarrow \infty.$$

Hence, we are in the position to apply [7, Lemma B.2] to u^* and the sequence of rescalings $(u_k^*)_{k \in \mathbb{N}}$ to deduce that $\tilde{p} = p$, which proves that p is the blow-down limit of u^* . In conclusion, u^* is the desired paraboloid solution.

Finally, the remaining statements follow from Proposition 3.4 and Lemma 3.2. □

REFERENCES

- [1] H. W. Alt, L. A. Caffarelli, and A. Friedman. Variational problems with two phases and their free boundaries. *Trans. Amer. Math. Soc.*, 282(2):431–461, 1984.
- [2] L. A. Caffarelli. The regularity of free boundaries in higher dimensions. *Acta Math.*, 139(3-4):155–184, 1977.
- [3] L. A. Caffarelli. The obstacle problem revisited. *J. Fourier Anal. Appl.*, 4(4-5):383–402, 1998.
- [4] L. A. Caffarelli, L. Karp, and H. Shahgholian. Regularity of a free boundary with application to the Pompeiu problem. *Ann. of Math. (2)*, 151(1):269–292, 2000.
- [5] E. DiBenedetto and A. Friedman. Bubble growth in porous media. *Indiana Univ. Math. J.*, 35(3):573–606, 1986.
- [6] P. Dive. Attraction des ellipsoïdes homogènes et réciproques d’un théorème de Newton. *Bull. Soc. Math. France*, 59:128–140, 1931.
- [7] S. Eberle, H. Shahgholian, and G. S. Weiss. On global solutions of the obstacle problem. *Duke Math. J.*, 172(11):2149–2193, 2023.
- [8] S. Eberle, H. Shahgholian, and G. S. Weiss. The structure of the regular part of the free boundary close to singularities in the obstacle problem. *J. Differential Equations*, 377:873–887, 2023.
- [9] S. Eberle and G. S. Weiss. Characterizing compact coincidence sets in the obstacle problem – a short proof. *Algebra i Analiz, St. Petersburg mathematical journal*, 32:137–145, 2020.
- [10] A. Friedman and M. Sakai. A characterization of null quadrature domains in \mathbf{R}^N . *Indiana Univ. Math. J.*, 35(3):607–610, 1986.
- [11] W. K. Hayman and E. F. Lingham. *Research problems in function theory*. Problem Books in Mathematics. Springer, Cham, 2019. Fiftieth anniversary edition of [MR0217268].
- [12] L. Karp. Generalized Newton potential and its applications. *J. Math. Anal. Appl.*, 174(2):480–497, 1993.
- [13] L. Karp and A. S. Margulis. Newtonian potential theory for unbounded sources and applications to free boundary problems. *J. Anal. Math.*, 70:1–63, 1996.
- [14] L. Karp and A. S. Margulis. Null quadrature domains and a free boundary problem for the Laplacian. *Indiana Univ. Math. J.*, 61(2):859–882, 2012.
- [15] D. Kinderlehrer and L. Nirenberg. Regularity in free boundary problems. *Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4)*, 4(2):373–391, 1977.
- [16] H. Lewy. An inversion of the obstacle problem and its explicit solution. *Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4)*, 6(4):561–571, 1979.
- [17] A. Petrosyan, H. Shahgholian, and N. Uraltseva. *Regularity of free boundaries in obstacle-type problems*, volume 136 of *Graduate Studies in Mathematics*. American Mathematical Society, Providence, RI, 2012.
- [18] M. Sakai. Null quadrature domains. *J. Analyse Math.*, 40:144–154 (1982), 1981.
- [19] H. Shahgholian. On quadrature domains and the Schwarz potential. *J. Math. Anal. Appl.*, 171(1):61–78, 1992.