

NON-TRAPPING ESTIMATES NEAR NORMALLY HYPERBOLIC TRAPPING

PETER HINTZ AND ANDRAS VASY

ABSTRACT. In this paper we prove semiclassical resolvent estimates for operators with normally hyperbolic trapping which are lossless relative to non-trapping estimates but take place in weaker function spaces. In particular, we obtain non-trapping estimates in standard L^2 spaces for the resolvent sandwiched between operators which localize away from the trapped set Γ in a rather weak sense, namely whose principal symbols vanish on Γ .

1. INTRODUCTION

The purpose of this paper is to obtain semiclassical estimates for pseudodifferential operators $P_h(z)$ with normally hyperbolic trapping for z real which are lossless relative to non-trapping estimates, but take place in weaker function spaces which are defined in a manner related to the Hamiltonian dynamics. Thus, the main result is an estimate of the form

$$\|u\|_{\mathcal{H}_{h,\Gamma}} \leq Ch^{-1} \|P_h(z)u\|_{\mathcal{H}_{h,\Gamma}^*},$$

with certain function spaces $\mathcal{H}_{h,\Gamma}$ and $\mathcal{H}_{h,\Gamma}^*$, described below; away from the trapped set these are just standard L^2 spaces. As the main application of such estimates is in so-called b-spaces, e.g. Kerr-de Sitter spaces, for which the estimates follow from the semiclassical ones immediately in the presence of dilation invariance, we also prove their counterpart in the general, non-dilation-invariant, b-setting. Extensions of special cases of these estimates play an important role in the recent global analysis of nonlinear wave equations on asymptotically Kerr-de Sitter spaces by the authors [7].

So at first we consider a family $P_h(z)$ of semiclassical pseudodifferential operators $P_h(z) \in \Psi_h(X)$ on a closed manifold X , depending smoothly on the parameter $z \in \mathbb{C}$, with normally hyperbolic trapping at the trapped set Γ , and assume that $P_h(z)$ is formally self-adjoint near Γ for $z \in \mathbb{R}$; moreover, we add complex absorption W in such a way that all forward and backward bicharacteristics outside Γ either enter the elliptic set of W in finite time or tend to Γ , and in at least one of the two directions they tend to the elliptic set of W . The bicharacteristics tending to Γ in the forward/backward directions are forward/backward trapped; denote by Γ_- , resp. Γ_+ the forward, resp. backward trapped set,¹ and assume that these are

Date: May 12, 2014.

The authors were supported in part by A.V.'s National Science Foundation grants DMS-0801226 and DMS-1068742 and P. H. was supported in part by a Gerhard Casper Stanford Graduate Fellowship and the German National Academic Foundation. They are grateful to the referee for comments that significantly improved the exposition in the manuscript.

¹In the notation of Wunsch and Zworski [13], which we recall below in Section 2.1, Γ_{\pm} are the backward/forward trapped sets for all (not necessarily null) bicharacteristics near the, say, zero

smooth codimension one submanifolds of T^*X which intersect transversally in Γ , which we moreover assume to be symplectic.

In this normally hyperbolic setting, under additional hypotheses, Wunsch and Zworski [13] have shown polynomial semiclassical resolvent estimates

$$\|u\| \leq Ch^{-N} \|P_h(z)u\|, \quad 0 < h < h_0, \quad (1.1)$$

in small strips $|\operatorname{Im} z| \leq ch$, $c > 0$ sufficiently small, $N > 1$, and indeed for z real, the loss (as compared to non-trapping estimates, which hold in many cases where there is no trapping, and which lose a power of h^{-1}) is merely logarithmic, i.e. one has

$$\|u\| \leq Ch^{-1}(\log h^{-1}) \|P_h(z)u\|, \quad 0 < h < h_0, \quad (1.2)$$

where $\|\cdot\|$ is the L^2 -norm; Bony, Burq and Ramond [1] showed that (1.2) is indeed sharp. Dyatlov [6] improved these estimates in $\operatorname{Im} z < 0$ by making c and N explicit; in a more general setting, Nonnenmacher and Zworski [10] obtained the optimal value for c .

We are concerned with improved estimates (for z almost real) if one localizes u and $P_h(z)u$ away from the trapping Γ in a rather weak sense, such as by applying pseudodifferential operators with symbols vanishing at Γ . To place this into context, recall that Datchev and Vasy [3, 4] have shown that under our assumptions, with $\operatorname{Im} z = \mathcal{O}(h^\infty)$, if $A, B \in \Psi_h(X)$ with $\operatorname{WF}'_h(A) \cap \Gamma = \operatorname{WF}'_h(B) \cap \Gamma = \emptyset$, B elliptic on $\operatorname{WF}'_h(A)$, then for all M there is N such that

$$\|Au\| \leq Ch^{-1} \|BP_h(z)u\| + C'h^M \|u\| + C''h^{-N} \|(\operatorname{Id} - B)P_h(z)u\|. \quad (1.3)$$

Thus, if $P_h(z)u$ is $\mathcal{O}(h^{N-1})$ at Γ (corresponding to the $\operatorname{Id} - B$ term in the estimate), then on the elliptic set of A , hence off Γ by appropriate choice of A , u satisfies non-trapping semiclassical estimates:

$$\|AP_h(z)^{-1}Av\| \leq Ch^{-1} \|v\|,$$

with A as above (take B as above with $\operatorname{WF}'_h(\operatorname{Id} - B) \cap \operatorname{WF}'_h(A) = \emptyset$). Here the $\mathcal{O}(h^\infty)$ bound on $\operatorname{Im} z$ arises from the a priori estimate, (1.1), and if $1 < N < 3/2$, e.g. as is on, and sufficiently near, the real axis,² then one can take $\operatorname{Im} z = \mathcal{O}(h^{-1+2N})$. The purpose of this paper is to improve this result by relaxing the conditions on $\operatorname{WF}'_h(A)$ and $\operatorname{WF}'_h(B)$ in (1.3).

The main point of the theorem below is thus that its estimate degenerates only at, as opposed to near, Γ . The proof given here is closely related to the proof of Wunsch and Zworski [13, §4] but can take place in a significantly simpler, standard semiclassical pseudodifferential algebra, at the cost of being suboptimal in terms of the L^2 -estimate, even though it is optimal (i.e. non-trapping) when a pseudodifferential operator with vanishing principal symbol at Γ is applied from both sides. To set this up, let $Q_\pm \in \Psi_h^{-\infty}(X)$ be self-adjoint and have symbols which are defining functions of Γ_\pm near Γ , say on a neighborhood O of Γ . Let $Q_0 \in \Psi_h^0(X)$ be a semiclassical operator with $\operatorname{WF}'_h(Q_0) \cap \Gamma = \emptyset$ which is elliptic on O^c (and thus on a neighborhood of O^c), with real principal symbol for convenience. One considers *normally isotropic* spaces at Γ , denoted $\mathcal{H}_{h,\Gamma}$, with squared norms given by

$$\|u\|_{\mathcal{H}_{h,\Gamma}}^2 = \|Q_0u\|^2 + \|Q_+u\|^2 + \|Q_-u\|^2 + h\|u\|^2;$$

level set of the semiclassical principal symbol $p_{h,z}$, and Γ_\pm^λ are the corresponding sets within the λ -level set of $p_{h,z}$.

²In the latter case by the Phragmén-Lindelöf theorem.

this is just the standard L^2 -space microlocally away from Γ as one of Q_+ , Q_- or Q_0 is elliptic there, and it does not depend on the choice of Q_0 as on $O \setminus \Gamma$ one of Q_+ and Q_- is elliptic at every point. The dual space relative to L^2 is then³

$$\mathcal{H}_{h,\Gamma}^* = h^{1/2}L^2 + Q_+L^2 + Q_-L^2 + Q_0L^2$$

(which is L^2 as a space, but with this norm); $P_h(z)u$ will then be measured in $\mathcal{H}_{h,\Gamma}^*$.

Theorem 1.1. *Let $P = P_h(z)$, Q_\pm be as above, $\text{Im } z = \mathcal{O}(h^2)$. Then*

$$\|Q_+u\| + \|Q_-u\| \leq Ch^{-1}\|Pu\|_{\mathcal{H}_{h,\Gamma}^*} + C'h^{1/2}\|u\|, \quad (1.4)$$

and thus by (1.2),

$$\|u\|_{\mathcal{H}_{h,\Gamma}} \leq Ch^{-1}\|Pu\|_{\mathcal{H}_{h,\Gamma}^*}. \quad (1.5)$$

In fact, we also obtain a direct proof of (1.5) without using (1.2) at the end of Section 2. Note that this theorem in particular implies the main result of [3] in this setting, in that the estimates are of the same kind, except that in [3] Pu is assumed to be microlocalized away from Γ , and u is estimated microlocally away from Γ .

The aforementioned b-estimates will be proved in Section 3, see Theorem 3.2.

2. SEMICLASSICAL RESOLVENT ESTIMATES ON THE REAL LINE

2.1. Notation and definitions. We will review some definitions of semiclassical analysis, partially in order to fix our notation. For a general reference, see Zworski [14].

Let X be a compact n -dimensional manifold without boundary, and fix a smooth density on X .

- For $u \in L^2(X)$, denote by $\|u\|$ its $L^2(X)$ norm; moreover, denote by $\langle \cdot, \cdot \rangle$ the (sesquilinear) inner product on $L^2(X)$.
- A family of functions $u = (u_h)_{h \in (0,1)}$ on X is *polynomially bounded* if $\|u\| \leq Ch^{-N}$ for some N . If $k \in \mathbb{R}$, we say that $u \in \mathcal{O}(h^k)$ if $\|u\| \leq C_k h^k$, and $u \in \mathcal{O}(h^\infty)$ if $\|u\| \leq C_N h^N$ for every N .
- For $a = (a_h)_{h \in (0,1)} \in \mathcal{C}^\infty(T^*X)$, we say $a \in h^k S^m(T^*X)$ if a satisfies

$$|\partial_z^\alpha \partial_\zeta^\beta a_h(z, \zeta)| \leq C_{\alpha\beta} h^k \langle \zeta \rangle^{m-|\beta|}$$

for all multiindices α, β and all $N \in \mathbb{N}$ in any coordinate chart, where the z are coordinates in the base and ζ coordinates in the fiber. We define the *semiclassical quantization* $\text{Op}_h(a)$ of a by

$$\text{Op}_h(a)u(z) = (2\pi h)^{-n} \int e^{iz\zeta/h} a(z, \zeta) \hat{u}(\zeta/h) d\zeta$$

for $u \in \mathcal{C}_c^\infty(X)$ supported in a chart and for general $u \in \mathcal{C}^\infty(X)$ by using a partition of unity. We write $\text{Op}_h(a) \in h^k \Psi_h^m(X)$. The quantization depends on the choice of partition of unity, but the resulting class of operators does not, modulo operators that have Schwartz kernel in $h^\infty \mathcal{C}^\infty(X^2)$. We say that a is a *symbol* of $\text{Op}_h(a)$. The equivalence class of a in $h^k S^m(T^*X)/h^{k-1} S^{m-1}(T^*X)$ is invariantly defined and is called the *principal symbol* of $\text{Op}_h(a)$. All operators below except $Q_0 \in \Psi_h^0(X)$ will in fact have compact microsupport in

³One really has Q_\pm^* and Q_0^* in this formula, but the reality of the principal symbols assures that one may replace them by Q_\pm and Q_0 modulo hL^2 . See [9, Appendix A] for a general discussion of the underlying functional analysis; also see Footnote 11.

the sense that they are quantizations of symbols $a \in h^k S^m(T^*X)$ satisfying in addition for all N

$$|\partial_z^\alpha \partial_\zeta^\beta a_h(z, \zeta)| \leq C_N h^N \langle \zeta \rangle^{-N} \text{ for all multiindices } \alpha, \beta$$

for ζ outside of a compact subset of T^*X . We denote the class of such symbols by $h^k S(T^*X)$ and the corresponding class of operators by $h^k \Psi_h(X)$.

- If $A, B \in \Psi_h(X)$, then $[A, B] \in h\Psi_h(X)$, and its principal symbol is $\frac{h}{i} \mathbf{H}_a b$, where we define the *Hamilton vector field* in a coordinate chart by

$$\mathbf{H}_a = (\partial_\zeta a) \partial_z - (\partial_z a) \partial_\zeta.$$

- By a *bicharacteristic* of A we mean an integral curve of the Hamilton vector field of the principal symbol of A . We denote the integral curve passing through the point $\rho \in T^*X$ by γ_ρ , i.e. $\gamma_\rho(0) = \rho$ and $\gamma'_\rho(s) = \mathbf{H}_a(\gamma_\rho(s))$. We shall also write $\phi^s(\rho) := \gamma_\rho(s)$ for the bicharacteristic flow.
- For a polynomially bounded family $(u_h)_{h \in (0,1)}$ and $k \in \mathbb{R} \cup \{\infty\}$, we say that $u = \mathcal{O}(h^k)$ at a point $\rho \in T^*X$ if there exists $a \in S(T^*X)$ with $a(\rho) \neq 0$ such that $\|\text{Op}_h(a)u\| = \mathcal{O}(h^k)$. We define the *semiclassical wave front set* $\text{WF}_h(u)$ of u as the complement of the set of all $\rho \in T^*X$ at which $u = \mathcal{O}(h^\infty)$.
- The *microsupport* of $A = \text{Op}_h(a) \in h^k \Psi_h(X)$, denoted $\text{WF}'_h(A)$, is the complement of the set of all $\rho \in T^*X$ so that $|\partial^\alpha a| = \mathcal{O}(h^\infty)$ near ρ for every multiindex α , in any (and therefore in every) coordinate chart.
- For $A \in h^k \Psi_h(X)$ with principal symbol $a \in h^k S(T^*X)$, we say that A is *elliptic* at $\rho \in T^*X$ if there is a constant $C > 0$ such that $|a(\rho')| \geq Ch^k$ for ρ' near ρ and h sufficiently small. For a subset $E \Subset T^*X$, we say that A is elliptic on E if A is elliptic at each point of E . If $A \in h^k \Psi_h(X)$ is elliptic on $E \Subset T^*X$ and $Au = f$ with u, f polynomially bounded and f is $\mathcal{O}(1)$ on E , then *microlocal elliptic regularity* states that u is $\mathcal{O}(h^{-k})$ on E .
- The *semiclassical characteristic set* of the semiclassical operator $A \in \Psi_h(X)$ with principal symbol a is defined by $\Sigma_h = \{\rho \in T^*X : a(\rho) = 0\}$.
- If $A \in \Psi_h(X)$ has a principal symbol with non-positive imaginary part, u, f are polynomially bounded, $Au = f$, and $u = \mathcal{O}(h^k)$ at ρ , $f = \mathcal{O}(h^{k+1})$ on $\gamma_\rho([0, T])$ for some $T > 0$, then the *propagation of singularities* states that $u = \mathcal{O}(h^k)$ at $\gamma_\rho(T)$.
- Let $P \in \Psi_h(X)$ be a semiclassical operator. Let $U \subset X$ denote an open subset so that the cotangent bundle over U contains what will be the trapped set, and place complex absorbing potentials in a neighborhood of U^c .⁴ We recall the notion of *normal hyperbolicity* from [13]: Define the *backward*, resp. *forward, trapped set* Γ_+ , resp. Γ_- , by

$$\Gamma_\pm = \{\rho \in T^*X : \gamma_\rho(s) \notin T_{U^c}^*X \text{ for all } \mp s \geq 0\}.$$

Let $\Gamma_\pm^\lambda = \Gamma_\pm \cap p^{-1}(\lambda)$ be the backward/forward trapped set within the energy surface $p^{-1}(\lambda)$, and define the *trapped set* $\Gamma_\lambda := \Gamma_+^\lambda \cap \Gamma_-^\lambda$. We say that P is *normally hyperbolically trapping* if:

- (1) There exists $\delta > 0$ such that $dp \neq 0$ on $p^{-1}(\lambda)$ for $|\lambda| < \delta$;
- (2) $\Gamma_\pm \cap p^{-1}(-\delta, \delta)$ are smooth codimension one submanifolds intersecting transversally at $\Gamma \cap p^{-1}(-\delta, \delta)$, and $\Gamma \cap p^{-1}(-\delta, \delta)$ is symplectic;

⁴See [13, 11] for details; the point here is that the relevant part of our analysis takes place microlocally near the trapped set, and the complex absorbing potentials allow us to ‘cut off’ the bicharacteristic flow in a neighborhood of the trapped set.

- (3) the flow is hyperbolic in the normal directions to Γ_λ within the energy surface: There exist subbundles E_λ^\pm of $T_{\Gamma_\lambda}(\Gamma_\pm^\lambda)$ such that $T_{\Gamma_\lambda}\Gamma_\pm^\lambda = T\Gamma_\lambda \oplus E_\lambda^\pm$, where $d\phi^s: E_\lambda^\pm \rightarrow E_\lambda^\pm$, and there exists $\theta > 0$ such that for all $|\lambda| < \delta$

$$\|d\phi^s(v)\| \leq Ce^{-\theta|t|}\|v\| \text{ for all } v \in E_\lambda^\mp, \pm t \geq 0.$$

2.2. Details on the setup and proof of the main result. Let $p = p_{h,z}$ be the semiclassical principal symbol of $P = P_h(z)$. Recall from the work of Wunsch and Zworski [13, Lemma 4.1], with a corrected argument in [12], that for defining functions ϕ_\pm of Γ_\pm (near Γ , namely in a neighborhood O of Γ) one can take ϕ_\pm with

$$H_p\phi_\pm = \mp c_\pm^2\phi_\pm$$

with $c_\pm > 0$ near Γ , and with⁵

$$\{\phi_+, \phi_-\} > 0$$

near Γ . *This is the only relevant feature of normal hyperbolicity for this paper; thus these identities and estimates could be taken as its definition for our purposes.* By shrinking O if necessary we may assume that this Poisson bracket as well as c_\pm have positive lower bounds on O . Then notice that

$$H_p\phi_+^2 = -2c_+^2\phi_+^2, \quad H_p\phi_-^2 = 2c_-^2\phi_-^2.$$

As indicated in the introduction, we consider *normally isotropic* spaces at Γ , denoted $\mathcal{H}_{h,\Gamma}$, with squared norms given by

$$\|u\|_{\mathcal{H}_{h,\Gamma}}^2 = \|Q_0u\|^2 + \|Q_+u\|^2 + \|Q_-u\|^2 + h\|u\|^2;$$

we can take Q_\pm with principal symbol ϕ_\pm , while Q_0 is elliptic on O^c with real principal symbol. This is just the standard L^2 -space microlocally away from Γ as one of Q_+ , Q_- and Q_0 is elliptic there, and it does not depend on the choice of Q_0 as on $O \setminus \Gamma$ one of Q_+ and Q_- is elliptic at every point. Notice that in fact

$$(Q_+ - iQ_-)^*(Q_+ - iQ_-) = Q_+^*Q_+ + Q_-^*Q_- - i[Q_+, Q_-]$$

and if $B \in \Psi_h(X)$ with $\text{WF}'_h(B) \subset O$ then

$$h\|Bv\|^2 \leq C \text{Re}\langle i[Q_+, Q_-]Bv, Bv \rangle + Ch^{N'}\|v\|^2,$$

$C > 0$, in view of $\{\phi_+, \phi_-\} > 0$ on O , so

$$Q_+^*Q_+ + Q_-^*Q_- = \frac{1}{2}(Q_+^*Q_+ + Q_-^*Q_- + (Q_+ - iQ_-)^*(Q_+ - iQ_-) + i[Q_+, Q_-])$$

shows that, for $h > 0$ small, the norm on $\mathcal{H}_{h,\Gamma}$ is equivalent to just the norm

$$\|u\|_{\mathcal{H}_{h,\Gamma,2}}^2 = \|Q_0u\|^2 + \|Q_+u\|^2 + \|Q_-u\|^2.$$

As mentioned in the introduction, the dual space relative to L^2 is then

$$\mathcal{H}_{h,\Gamma}^* = h^{1/2}L^2 + Q_+L^2 + Q_-L^2 + Q_0L^2.$$

Then $\Psi_h(X)$ acts on $\mathcal{H}_{h,\Gamma}$, and thus on $\mathcal{H}_{h,\Gamma}^*$, for $B \in \Psi_h(X)$ preserves $h^{-1/2}L^2$ and gives

$$\|Q_+Bu\| \leq \|BQ_+u\| + \|[Q_+, B]u\| \leq C\|Q_+u\| + h\|u\|_{L^2},$$

⁵These defining functions exist globally when Γ_\pm is orientable; but even if Γ_\pm is not such, the square is globally defined. There is only a minor change required below if ϕ_\pm are not well defined; see Footnote 6.

with a similar result for Q_- and Q_0 . We remark that the notation $\mathcal{H}_{h,\Gamma}$ is justified as the space depends only on Γ , not on the particular defining functions ϕ_\pm as any other defining functions would change Q_\pm by an elliptic factor modulo an element of $h\Psi_h(X)$, whose contribution to the squared norm can be absorbed into $Ch^2\|u\|_{L^2}^2$, and thus dropped altogether (for h small) in view of the equivalence of the two norms discussed above.

We are now ready to prove Theorem 1.1. We remark that the microlocal version of the two estimates of the theorem is that given any neighborhood O' of Γ with closure in O , there exist $B_0 \in \Psi_h(X)$ elliptic at Γ , $B_1, B_2 \in \Psi_h(X)$ with $\text{WF}'_h(B_2) \cap \Gamma_+ = \emptyset$, $\text{WF}'_h(B_j) \subset O'$ for $j = 0, 1, 2$ such that

$$\|B_0Q_+u\| + \|B_0Q_-u\| \leq h^{-1}\|B_1Pu\|_{\mathcal{H}_{h,\Gamma}^*} + \|B_2u\|_{L^2} + C'h^{1/2}\|u\|_{L^2}, \quad (2.1)$$

respectively

$$\|B_0u\|_{\mathcal{H}_{h,\Gamma}} \leq h^{-1}\|B_1Pu\|_{\mathcal{H}_{h,\Gamma}^*} + \|B_2u\|_{L^2} + C'h\|u\|_{L^2}; \quad (2.2)$$

see (2.9). The theorem is then proved by controlling the B_2u term using the backward non-trapped nature of $\Gamma_- \setminus \Gamma$.

Proof of Theorem 1.1. We first prove (1.4), which proves (1.5) by (1.2). In fact, one can also give a direct proof of (1.5) without using (1.2); see the discussion following this proof.

Let $\chi_0(t) = e^{-1/t}$ for $t > 0$, $\chi_0(t) = 0$ for $t \leq 0$, $\chi \in C_c^\infty([0, \infty))$ be identically 1 near 0 with $\chi' \leq 0$, and indeed with $\chi'\chi = -\chi_1^2$, $\chi_1 \geq 0$, $\chi_1 \in C_c^\infty([0, \infty))$, and let $\psi \in C_c^\infty(\mathbb{R})$ be identically 1 near 0. Let

$$a = \chi_0(\phi_+^2 - \phi_-^2 + \kappa)\chi(\phi_+^2)\psi(p),$$

$\kappa > 0$ small. Notice that on $\text{supp } a$, if χ is supported in $[0, R]$,

$$\phi_+^2 \leq R, \quad \phi_-^2 \leq \phi_+^2 + \kappa = R + \kappa,$$

so a is localized near Γ if R and κ are taken sufficiently small. Then

$$\begin{aligned} \frac{1}{4}\mathbb{H}_p(a^2) &= -(c_+^2\phi_+^2 + c_-^2\phi_-^2)(\chi_0\chi_0')(\phi_+^2 - \phi_-^2 + \kappa)\chi(\phi_+^2)^2\psi(p)^2 \\ &\quad - c_+^2\phi_+^2(\chi'\chi)(\phi_+^2)\chi_0(\phi_+^2 - \phi_-^2 + \kappa)^2\psi(p)^2. \end{aligned}$$

Now $\chi_0' \geq 0$, so the two terms have opposite signs. Let⁶

$$a_\pm = \phi_\pm \sqrt{(\chi_0\chi_0')(\phi_+^2 - \phi_-^2 + \kappa)\chi(\phi_+^2)\psi(p)},$$

and

$$e_- = c_+\phi_+\chi_1(\phi_+^2)\chi_0(\phi_+^2 - \phi_-^2 + \kappa)\psi(p);$$

then

$$\frac{1}{4}\mathbb{H}_p(a^2) = -c_+^2a_+^2 - c_-^2a_-^2 + e_-^2. \quad (2.3)$$

Here

$$\begin{aligned} \text{supp } e_- &\subset \text{supp } a, \\ \text{supp } e_- \cap \Gamma_+ &= \emptyset, \end{aligned}$$

⁶If ϕ_\pm is not defined globally, a_\pm are not defined as stated. (The term e_-^2 need not have a sign, so this issue does not arise for it; see the Weyl quantization argument below.) However, a_\pm need not be real below, so as long as one can choose ψ_\pm complex valued with $|\psi_\pm|^2 = \phi_\pm^2$, replacing the first factor of ϕ_\pm with ψ_\pm in the definition of a_\pm allows one to complete the argument in general.

with the last statement following from ϕ_+^2 taking values away from 0 on $\text{supp } \chi_1$; see Figure 1.




FIGURE 1. Supports of the commutant a and the error term e_- in the positive commutator argument of the non-trapping estimate near the trapped set Γ , Theorem 1.1. The support of a is indicated in light gray; on $\text{supp } a \setminus \text{supp } e_-$, darker colors correspond to larger values of a . Also shown are the forward, resp. backward, trapped set Γ_- , resp. Γ_+ , and the bicharacteristic flow nearby. The figure already suggests that $H_p(a^2)$ is non-positive away from $\text{supp } e_-$, and actually negative away from $\text{supp } e_- \cup \Gamma$; see equation (2.3).

One then takes $A \in \Psi_h(X)$ with principal symbol a , and with $\text{WF}'_h(A) \subset \text{supp } a$, $A_\pm \in \Psi_h(X)$ with principal symbols of a_\pm , and with $\text{WF}'_h(A_\pm) \subset \text{supp } a_\pm$, C_\pm have symbol c_\pm and with $\text{WF}'_h(C_\pm) \subset \text{supp } c_\pm$; one similarly lets $E_- \in \Psi_h(X)$ have principal symbol e_- , and wave front set in the support of e_- . This gives that

$$\frac{i}{4h}[P, A^*A] = -(C_+A_+)^*(C_+A_+) - (C_-A_-)^*(C_-A_-) + E_-^*E_- + hF, \quad (2.4)$$

for some $F \in \Psi_h(X)$ with

$$\text{WF}'_h(F) \subset \text{supp } a.$$

Thus

$$\frac{i}{4h}\langle [P, A^*A]u, u \rangle = -\|C_+A_+u\|^2 - \|C_-A_-u\|^2 + \|E_-u\|^2 + h\langle Fu, u \rangle.$$

Expanding the left hand side gives

$$\begin{aligned} & \langle PA^*Au, u \rangle - \langle A^*APu, u \rangle \\ &= \langle Au, APu \rangle - \langle APu, Au \rangle + \langle (P - P^*)A^*Au, u \rangle. \end{aligned}$$

As we are assuming that $P - P^*$ is $\mathcal{O}(h^2)$ near Γ , we may also assume that this holds on $\text{supp } a$, thus the last term is $\mathcal{O}(h^2)\|u\|^2$. Thus,

$$\|C_+A_+u\|^2 + \|C_-A_-u\|^2 \leq \|E_-u\|^2 + h^{-1}|\langle APu, Au \rangle| + C_1h\|u\|^2. \quad (2.5)$$

Now, by the duality of $\mathcal{H}_{h,\Gamma}$ and $\mathcal{H}_{h,\Gamma}^*$ relative to the L^2 inner product,

$$|\langle APu, Au \rangle| \leq \|APu\|_{\mathcal{H}_{h,\Gamma}^*} \|Au\|_{\mathcal{H}_{h,\Gamma}} \leq \frac{h\epsilon}{2} \|Au\|_{\mathcal{H}_{h,\Gamma}}^2 + \frac{1}{2h\epsilon} \|APu\|_{\mathcal{H}_{h,\Gamma}^*}^2. \quad (2.6)$$

Further, for $\epsilon > 0$ small, $\epsilon \|Q_+ Au\|^2$ can be estimated in terms of $\|C_+ A_+ u\|^2 + \mathcal{O}(h)\|u\|^2$, as can be seen by comparing the principal symbols, in particular using the ellipticity of C_+ on $\text{supp } a$. One can thus absorb $\frac{\epsilon}{2} \|Au\|_{\mathcal{H}_{h,\Gamma}}^2$ into the left hand side of (2.5). This shows

$$\|C_+ A_+ u\|^2 + \|C_- A_- u\|^2 \leq C \|E_- u\|^2 + Ch^{-2} \|APu\|_{\mathcal{H}_{h,\Gamma}^*}^2 + Ch \|u\|^2.$$

Now, since the region $\text{supp } e_-$ is disjoint from Γ_+ , it is backward non-trapped, and thus the standard propagation of singularities with complex absorption (see e.g. [4, Lemma 5.1]) implies that $E_- u$ is controlled by Pu microlocalized off Γ_+ , hence by $Q_+ Pu$, modulo higher order (in h) terms in Pu . This proves the first part of Theorem 1.1 since A_\pm is an elliptic multiple of Q_\pm microlocally near Γ . Thus, if we have a bound $\|u\| \leq C'h^{-1-s} \|Pu\|_{L^2}$, $0 < s < 1/2$, and thus $h\|u\|^2 \leq C'h^{-1-2s} \|Pu\|_{L^2}^2 \leq C''h^{-1-2s} \|Pu\|_{\mathcal{H}_{h,\Gamma}^*}^2$, this implies a non-trapping estimate:

$$\|u\|_{\mathcal{H}_{h,\Gamma}} \leq Ch^{-1} \|Pu\|_{\mathcal{H}_{h,\Gamma}^*}.$$

This completes the proof of Theorem 1.1. \square

In fact, as mentioned earlier, a slight change of point of view proves Theorem 1.1 directly. To see this, we use the Weyl quantization⁷ when choosing a, a_\pm, c_\pm, e_- ; since we are on a manifold, this requires identifying functions with half-densities via trivialization of the half-density bundle by the Riemannian metric; this identification preserves self-adjointness. We also write $P_{h,z}$ as the Weyl quantization of $p_0 + hp_1$ with p_0, p_1 real modulo $\mathcal{O}(h^2)$. Then the principal symbol calculation above holds with p_0 in place of p , and with p_1 included it yields additional terms

$$\begin{aligned} \frac{1}{4} \mathbf{H}_p(a^2) &= - (c_+^2 \phi_+^2 + c_-^2 \phi_-^2 - h\phi_+ \mathbf{H}_{p_1} \phi_+ + h\phi_- \mathbf{H}_{p_1} \phi_-) \\ &\quad \times (\chi_0 \chi'_0) (\phi_+^2 - \phi_-^2 + \kappa) \chi(\phi_+^2) \psi(p)^2 \\ &\quad - (c_+^2 \phi_+^2 - h\phi_+ \mathbf{H}_{p_1} \phi_+) (\chi' \chi) (\phi_+^2) \chi_0 (\phi_+^2 - \phi_-^2 + \kappa) \psi(p)^2. \end{aligned}$$

Now, (2.4) becomes

$$\begin{aligned} \frac{i}{4h} [P, A^* A] &= - (C_+ A_+)^* (C_+ A_+) - (C_- A_-)^* (C_- A_-) \\ &\quad + h(A_+^* G_+ + G_+^* A_+ + A_-^* G_- + G_-^* A_-) + E + h^2 F, \end{aligned} \quad (2.7)$$

with G_\pm being the Weyl quantization of

$$g_\pm = \pm \frac{1}{2} (\mathbf{H}_{p_1} \phi_\pm) \sqrt{(\chi_0 \chi'_0) (\phi_+^2 - \phi_-^2 + \kappa) \chi(\phi_+^2) \psi(p)},$$

and with $F \in \Psi_h(X)$ with

$$\text{WF}'_h(F) \subset \text{supp } a.$$

⁷The Weyl quantization is actually irrelevant. It is straightforward to see that if $A \in \Psi_h(X)$ and if the principal symbol of A is real then the real part of the subprincipal symbol is defined independently of choices. This is all that is needed for the argument below.

Correspondingly, (2.5) becomes

$$\begin{aligned} \|C_+A_+u\|^2 + \|C_-A_-u\|^2 &\leq |\langle Eu, u \rangle| + h^{-1}|\langle APu, Au \rangle| \\ &\quad + 2h\|A_+u\|\|G_+u\| + 2h\|A_-u\|\|G_-u\| + C_1h^2\|u\|^2. \end{aligned} \quad (2.8)$$

The terms with G_\pm on the right hand side can be estimated by

$$\epsilon\|A_+u\|^2 + \epsilon^{-1}h^2\|G_+u\|^2 + \epsilon\|A_-u\|^2 + \epsilon^{-1}h^2\|G_-u\|^2,$$

and for $\epsilon > 0$ sufficiently small, the $\|A_\pm u\|^2$ terms can now be absorbed into the left hand side of (2.8). Proceeding as above yields

$$\|C_+A_+u\|^2 + \|C_-A_-u\|^2 \leq C|\langle Eu, u \rangle| + Ch^{-2}\|APu\|_{\mathcal{H}_{h,\Gamma}^*}^2 + Ch^2\|u\|^2. \quad (2.9)$$

Together with the non-trapping for the E term this gives the global estimate

$$\|u\|_{\mathcal{H}_{h,\Gamma}}^2 \leq Ch^{-2}\|Pu\|_{\mathcal{H}_{h,\Gamma}^*}^2 + Ch^2\|u\|^2,$$

and now the last term on the right hand side can be absorbed into the left hand side for sufficiently small h , giving the estimate (1.5).

Notice that this also directly gives a weaker version of the Wunsch-Zworski estimate (1.2), namely

$$\|u\|_{L^2} \leq Ch^{-2}\|Pu\|_{L^2},$$

in view of the continuity of the inclusions $\mathcal{H}_{h,\Gamma} \hookrightarrow h^{-1/2}L^2$ and $h^{1/2}L^2 \hookrightarrow \mathcal{H}_{h,\Gamma}^*$.

Remark 2.1. If one is interested in a fixed operator, rather than in a parameter-dependent family of operators, one can naturally strengthen the estimates (1.4)–(1.5) by adding $h^{-1}\|\hat{P}_0u\|$ to the left hand sides, where \hat{P}_0 is any elliptic multiple of P . A more natural way of phrasing such an improvement is to use ‘coisotropic, normally isotropic’ spaces $\tilde{\mathcal{H}}_{h,\Gamma}$ and $\tilde{\mathcal{H}}_{h,\Gamma}^*$, where the squared norm on $\tilde{\mathcal{H}}_{h,\Gamma}$ is defined by

$$\|u\|_{\tilde{\mathcal{H}}_{h,\Gamma}}^2 = \|Q_0u\|^2 + \|Q_+u\|^2 + \|Q_-u\|^2 + h^{-1}\|\hat{P}_0u\|^2 + h\|u\|^2,$$

which strengthens the space and therefore weakens its dual. Using these spaces instead in (2.6), one obtains an additional term from $h\|Au\|_{\tilde{\mathcal{H}}_{h,\Gamma}}^2$, namely $\|\hat{P}_0Au\|^2$, which is bounded by $C(h^{-1}\|APu\|_{\tilde{\mathcal{H}}_{h,\Gamma}^*}^2 + h\|u\|^2)$, and thus the remainder of the second proof goes through.

3. NON-TRAPPING ESTIMATES IN NON-DILATION INVARIANT SETTINGS

We now transfer Theorem 1.1 into the b-setting; the discussion in the previous section is essentially the dilation invariant special case of this, as we will explain below, though in the b-setting there is additional localization near the boundary. One main application of the b-estimate is in the analysis of linear and non-linear waves on asymptotically Kerr-de Sitter spaces; see [11, 7] for details.

3.1. Notation and definitions. For a general reference for b-analysis, see Melrose [8].

Let M be an n -dimensional compact manifold with boundary X .

- Let $\mathcal{V}_b(M)$ be the Lie algebra of *b-vector fields* on M , i.e. of vector fields on M which are tangent to X . Elements of $\mathcal{V}_b(M)$ are sections of a natural vector bundle on M , namely the *b-tangent bundle* ${}^bT^*M$; in local coordinates (τ, x) near X , the fibers of ${}^bT^*M$ are spanned by $\tau\partial_\tau$ and ∂_x . The fibers of the dual bundle ${}^bT^*M$, called *b-cotangent bundle*, are spanned by $\frac{d\tau}{\tau}$ and dx .

It is often convenient to consider the fiber compactification $\overline{{}^bT^*M}$ of ${}^bT^*M$, where the fibers are replaced by their radial compactification. The new boundary of $\overline{{}^bT^*M}$ at fiber infinity is the *b-cosphere bundle* ${}^bS^*M$; it still possesses the compactification of the ‘old’ boundary $\overline{{}^bT^*}_X M$, see Figure 2. ${}^bS^*M$ is naturally the quotient of ${}^bT^*M \setminus o$ by the \mathbb{R}^+ -action of dilation in the fibers of the cotangent bundle. Many sets that we will consider below are conic subsets of ${}^bT^*M \setminus o$, and we will often view them as subsets of ${}^bS^*M$.




FIGURE 2. The radially compactified cotangent bundle $\overline{{}^bT^*M}$ near $\overline{{}^bT^*}_X M$; the cosphere bundle ${}^bS^*M$, which is the boundary at fiber infinity of $\overline{{}^bT^*M}$, is also shown, as well as the zero section $o_M \subset \overline{{}^bT^*M}$ and the zero section over the boundary $o_X \subset \overline{{}^bT^*}_X M$.

- For $a \in \mathcal{C}^\infty({}^bT^*M)$, we say $a \in S^m({}^bT^*M)$ if a satisfies

$$|\partial_z^\alpha \partial_\zeta^\beta a(z, \zeta)| \leq C_{\alpha\beta} \langle \zeta \rangle^{m-|\beta|} \text{ for all multiindices } \alpha, \beta$$

in any coordinate chart, where z are coordinates in the base and ζ coordinates in the fiber; more precisely, in local coordinates (τ, x) near X , we take $\zeta = (\sigma, \xi)$, where we write b-covectors as

$$\sigma \frac{d\tau}{\tau} + \sum_j \xi_j dx_j.$$

We define the quantization $\text{Op}(a)$ of a , acting on smooth functions u supported in a coordinate chart, by

$$\begin{aligned} \text{Op}(a)u(\tau, x) &= (2\pi)^{-n} \int e^{i(\tau-\tau')\tilde{\sigma}+i(x-x')\xi} \phi\left(\frac{\tau-\tau'}{\tau}\right) \\ &\quad \times a(\tau, x, \tau\tilde{\sigma}, \xi) u(\tau', x') d\tau' dx' d\tilde{\sigma} d\xi, \end{aligned}$$

where the τ' -integral is over $[0, \infty)$, and $\phi \in \mathcal{C}_c^\infty((-1/2, 1/2))$ is identically 1 near 0.⁸ For general u , define $\text{Op}(a)u$ using a partition of unity. We write $\text{Op}(a) \in \Psi_b^m(M)$. We say that a is a *symbol* of $\text{Op}(a)$. The equivalence class of a in $S^m({}^bT^*M)/S^{m-1}({}^bT^*M)$ is invariantly defined on ${}^bT^*M$ and is called

⁸The cutoff ϕ ensures that these operators lie in the ‘small b-calculus’ of Melrose, in particular that such quantizations act on weighted b-Sobolev spaces, defined below.

the *principal symbol* of $\text{Op}(a)$. We will tacitly assume that all our operators have homogeneous principal symbols.

- If $A \in \Psi_b^{m_1}(M)$ and $B \in \Psi_b^{m_2}(M)$, then $[A, B] \in \Psi_b^{m_1+m_2-1}(M)$, and its principal symbol is $\frac{1}{i}H_a b \equiv \frac{1}{i}\{a, b\}$, where the Hamilton vector field H_a of the principal symbol a of A is the extension of the Hamilton vector field from $T^*M^\circ \setminus o$ to ${}^bT^*M \setminus o$, which is a homogeneous degree $m-1$ vector field on ${}^bT^*M \setminus o$ tangent to the boundary ${}^bT_X^*M$. In local coordinates (τ, x, σ, ξ) on ${}^bT^*M$ as above, this has the form

$$H_a = (\partial_\sigma a)(\tau \partial_\tau) - (\tau \partial_\tau a) \partial_\sigma + \sum_j ((\partial_{\xi_j} a) \partial_{x_j} - (\partial_{x_j} a) \partial_{\xi_j}). \quad (3.1)$$

- We define bicharacteristics completely analogously to the semiclassical setting.
- The *microsupport* $\text{WF}_b^l(A) \subset {}^bT^*M \setminus o$ of $A = \text{Op}(a) \in \Psi_b^m(M)$ is the complement of the set of all $\rho \in {}^bT^*M \setminus o$ such that a is rapidly decaying in a conic neighborhood around ρ . Note that $\text{WF}_b^l(A)$ is conic, hence we will also view it as a subset of ${}^bS^*M$.
- Fix a *b-density* on M , which is locally of the form $a \left| \frac{d\tau}{\tau} dz \right|$, $a > 0$.
- Define the *b-Sobolev space* $H_b^k(M)$ for $k \in \mathbb{Z}_{\geq 0}$ by

$$H_b^k(M) = \{u \in L^2(M) : X_1 \cdots X_k u \in L^2(M), X_1, \dots, X_j \in \mathcal{V}_b(M)\},$$

and for general $k \in \mathbb{R}$ by duality and interpolation. Moreover, define the weighted b-Sobolev spaces $H_b^{s,\alpha}(M) := \tau^\alpha H_b^s(M)$ for $s, \alpha \in \mathbb{R}$, where τ is a boundary defining function, i.e. $\tau = 0$ at X and $d\tau \neq 0$ there. Every b-pseudodifferential operator $A \in \Psi_b^m(M)$ is a bounded operator $A: H_b^{s,\alpha}(M) \rightarrow H_b^{s-m,\alpha}(M)$, $s, \alpha \in \mathbb{R}$.

- For $A \in \Psi_b^m(M)$ with principal symbol $a \in S^m({}^bT^*M)$, we say that A is *elliptic* at $\rho \in {}^bT^*M \setminus o$ if there is a constant $C > 0$ such that $|a(z, \zeta)| \geq C|\zeta|^m$ for (z, ζ) in a conic neighborhood of ρ . The *characteristic set* of A is the complement (in ${}^bT^*M \setminus o$) of the set of all ρ at which A is elliptic.
- For $u \in H_b^{-\infty,\alpha}(M)$, define its $H_b^{s,\alpha}$ *wave front set* $\text{WF}_b^{s,\alpha}(u) \subset {}^bT^*M \setminus o$ as the complement of the set of all $\rho \in {}^bT^*M \setminus o$ for which there exists $a \in S^0({}^bT^*M)$ elliptic at ρ such that $\text{Op}(a)u \in H_b^{s,\alpha}(M)$. In particular, $\text{WF}_b^{s,\alpha}(u) = \emptyset$ if and only if $u \in H_b^{s,\alpha}(M)$.
- *Microlocal elliptic regularity* states that if $Au = f$ with $A \in \Psi_b^m(M)$, $u, f \in H_b^{-\infty,\alpha}(M)$, $\rho \notin \text{WF}_b^{s-m,\alpha}(f)$ and A is elliptic at ρ , then $\rho \notin \text{WF}_b^{s,\alpha}(u)$.
- If $A \in \Psi_b^m(M)$ has a principal symbol with non-positive imaginary part, $u, f \in H_b^{-\infty,\alpha}(M)$, $Au = f$, moreover $\rho \notin \text{WF}_b^{s,\alpha}(u)$ and $\gamma_\rho([0, T]) \cap \text{WF}_b^{s-m+1,\alpha}(f) = \emptyset$ for some $T > 0$, then the *propagation of singularities* states that $\gamma_\rho(T) \notin \text{WF}_b^{s,\alpha}(u)$.

3.2. Setup, statement and proof of the result. Suppose $\mathcal{P} \in \Psi_b^m(M)$, $\mathcal{P} - \mathcal{P}^* \in \Psi_b^{m-2}(M)$. Let p be the principal symbol of \mathcal{P} , which is thus a homogeneous degree m function on ${}^bT^*M \setminus o$, which we assume to be *real-valued*. Let $\tilde{\rho}$ denote a homogeneous degree -1 defining function of ${}^bS^*M$. Then the rescaled Hamilton vector field

$$V = \tilde{\rho}^{m-1} H_p$$

is a \mathcal{C}^∞ vector field on $\overline{{}^bT^*M}$ away from the 0-section, and it is tangent to all boundary faces. The characteristic set Σ is the zero-set of the smooth function $\tilde{\rho}^m p$ in ${}^bS^*M$. We will, somewhat imprecisely, refer to the flow of V in $\Sigma \subset {}^bS^*M$ as the Hamilton, or (null)bicharacteristic flow; its integral curves, the (null)bicharacteristics, are reparameterizations of those of the Hamilton vector field H_p , projected by the quotient map ${}^bT^*M \setminus o \rightarrow {}^bS^*M$.

We first work microlocally near the trapped set, namely assume that

- (1) $\Gamma \subset \Sigma \cap {}^bS_X^*M$ is a smooth submanifold disjoint from the image of $T^*X \setminus o$ (so τD_τ is elliptic near Γ),
- (2) Γ_+ is a smooth submanifold of $\Sigma \cap {}^bS_X^*M$ in a neighborhood U_1 of Γ ,
- (3) Γ_- is a smooth submanifold of Σ transversal to $\Sigma \cap {}^bS_X^*M$ in U_1 ,
- (4) Γ_+ has codimension 2 in Σ , Γ_- has codimension 1,
- (5) Γ_+ and Γ_- intersect transversally in Σ with $\Gamma_+ \cap \Gamma_- = \Gamma$,
- (6) the rescaled Hamilton vector field $V = \tilde{\rho}^{m-1} H_p$ is tangent to both Γ_+ and Γ_- , and thus to Γ .




FIGURE 3. An exemplary situation with trapping: Shown are the (projection from ${}^bS^*M$ to the base M of the) trapped set Γ , the b-cosphere bundle over X as well as a forward bicharacteristic starting at a point $\rho \in \Gamma_-$.

We assume that Γ_+ is backward trapped for the Hamilton flow (i.e. bicharacteristics in Γ_+ near Γ tend to Γ as the parameter goes to $-\infty$), i.e. is the unstable manifold of Γ , while Γ_- is forward trapped, i.e. is the stable manifold of Γ , see Figure 3; indeed, we assume a quantitative version of this. (There is a completely analogous statement if Γ_+ is forward trapped and Γ_- is backward trapped: replacing \mathcal{P} by $-\mathcal{P}$ preserves all assumptions, but reverses the Hamilton flow.) To state this, let ϕ_- be a defining function of Γ_- , and let $\phi_+ \in \mathcal{C}^\infty({}^bS^*M)$ be a defining function of Γ_+ in ${}^bS_X^*M$; thus Γ_+ is defined within ${}^bS^*M$ by $\tau = 0, \phi_+ = 0$. Notice that V being tangent to ${}^bS_X^*M$ (due to (3.1)) implies that $V\tau$ is a multiple of τ ; we assume that, near Γ ,

$$V\tau = -c_\partial^2 \tau, \quad c_\partial > 0; \quad (3.2)$$

this is consistent with the stability of Γ_- . By the tangency requirement, with

$$\hat{p}_0 = \tilde{\rho}^m p,$$

$V\phi_- = \alpha_- \phi_- + \nu_- \hat{p}_0$, α_- smooth; notice that changing ϕ_- by a smooth non-zero multiple f gives $V(f\phi_-) = \alpha_- f\phi_- + \nu_- f\hat{p}_0 + (Vf)\phi_-$, so α_- depends on the choice of ϕ_- . On the other hand, the tangency requirement gives $V\phi_+ = \alpha_+ \phi_+ + \beta_+ \tau + \nu_+ \hat{p}_0$. For the sake of conciseness, rather than stating the assumptions on the Hamilton flow as in [13], we assume directly that ϕ_{\pm} satisfy

$$V\phi_- = c_-^2 \phi_- + \nu_- \hat{p}_0, \quad V\phi_+ = -c_+^2 \phi_+ + \beta_+ \tau + \nu_+ \hat{p}_0, \quad (3.3)$$

with $c_{\pm} > 0$ smooth near Γ , β_+, ν_{\pm} smooth near Γ and

$$\{\phi_+, \phi_-\} > 0 \quad (3.4)$$

near Γ . Let $U_0 \subset \overline{U_0} \subset U_1$ be a neighborhood of Γ such that the Poisson bracket in (3.4) as well as c_{\pm} have positive lower bounds.

Now, given a boundary defining function τ , ${}^bS_X^*M \setminus S^*X$ (where S^*X is the image of T^*X under the \mathbb{R}^+ -dilation quotient) can be identified with $(\frac{d\tau}{\tau} + T^*X) \cup (-\frac{d\tau}{\tau} + T^*X)$ (which in turn can be identified with two copies of T^*X) since ${}^bT_X^*M = \text{Span}\{\frac{d\tau}{\tau}\} \oplus T^*X$, and each \mathbb{R}^+ -orbit outside T^*X intersects $(\frac{d\tau}{\tau} + T^*X) \cup (-\frac{d\tau}{\tau} + T^*X)$ in a unique point. This provides the connection between the b- and the semiclassical perspectives, i.e. analysis on ${}^bT_X^*M$ and that of T^*X . In fact, if p is a homogeneous degree m function, then $\tilde{\rho}^m p$, where $\tilde{\rho}$ can be taken as the reciprocal of the absolute value of the symbol of τD_{τ} in this region (which is well-defined, independent of choices), gives a function on $\{\pm 1\} \times T^*X$; this is exactly the semiclassical rescaling, with $\tilde{\rho}^m p$ the semiclassical principal symbol (depending on the parameter ± 1) of the rescaled operator family (cf. [11, §2.1]).

Notice that if we merely assume the normal hyperbolicity within ${}^bS_X^*M$, in the sense of this identification with T^*X , as in [13, §1.2], then [13, Lemma 4.1], as corrected in [12], actually gives such defining functions ϕ_{\pm}^0 *within* ${}^bS_X^*M$ (i.e. letting $\tau = 0$); taking an arbitrary extension in case of ϕ_+ , and an extension which is a defining function in case of Γ_- , all the requirements above are satisfied. In particular, Kerr and Kerr-de Sitter spaces satisfy these assumptions, as do their perturbations when the angular momentum $|a|$ is small; see [13, Proposition 2.1] for the Kerr setting and [11, §6] for the Kerr-de Sitter one. Indeed, in the Kerr case the full range of $|a| < M$, M the black hole mass, satisfies the hypotheses, as shown by Dyatlov [5].

There is an asymmetry between the roles of ϕ_{\pm} and τ , and thus we consider the parabolic defining function

$$\rho_+ = \phi_+^2 + M\tau$$

for Γ_+ , $M > 0$, to be chosen. Then, near Γ ,

$$\begin{aligned} \hat{\rho}_+ &= V\rho_+ = -2c_+^2 \phi_+^2 + 2\beta_+ \phi_+ \tau + 2\nu_+ \phi_+ \hat{p}_0 - M c_{\partial}^2 \tau \\ &= -2c_+^2 \phi_+^2 - (M c_{\partial}^2 - 2\beta_+ \phi_+) \tau + 2\nu_+ \phi_+ \hat{p}_0 \\ &\leq -\tilde{c}_+^2 \rho_+ + 2\nu_+ \phi_+ \hat{p}_0, \quad \tilde{c}_+ > 0, \end{aligned} \quad (3.5)$$

if $M > 0$ is chosen sufficiently large, consistently with the forward trapped nature of Γ_- . (Here the term with \hat{p}_0 is considered harmless as one essentially restricts to the characteristic set, $\hat{p}_0 = 0$.) Also, note that one can use⁹ the reciprocal

⁹Indeed, in the semiclassical setting, after Mellin transforming this problem, $|\sigma|^{-1}$ plays the role of the semiclassical parameter h , which in that case *commutes* with the operator.

$\tilde{\rho} = |\sigma|^{-1}$ of the principal symbol σ of τD_τ as the local defining function of ${}^bS^*M$ as fiber-infinity in ${}^bT^*M$ near Γ ; then

$$V\tilde{\rho} = \tilde{\alpha}\tilde{\rho}\tau \quad (3.6)$$

for some $\tilde{\alpha}$ smooth in view of (3.1).

Similar to the normally isotropic spaces in the semiclassical setting, we introduce spaces which are *normally isotropic at Γ* .¹⁰ Concretely, let $Q_\pm \in \Psi_b^0(M)$ have principal symbol ϕ_\pm as before, $\hat{P}_0 \in \Psi_b^0(M)$ have principal symbol \hat{p}_0 and let $Q_0 \in \Psi_b^0(M)$ be elliptic, with real principal symbol for convenience, on U_0^c (and thus nearby). Define the (global) b-normally isotropic spaces at Γ of order s , $\mathcal{H}_{b,\Gamma}^s$, by the norm

$$\|u\|_{\mathcal{H}_{b,\Gamma}^s}^2 = \|Q_0u\|_{H_b^s}^2 + \|Q_+u\|_{H_b^s}^2 + \|Q_-u\|_{H_b^s}^2 + \|\tau^{1/2}u\|_{H_b^s}^2 + \|\hat{P}_0u\|_{H_b^s}^2 + \|u\|_{H_b^{s-1/2}}^2, \quad (3.7)$$

and let $\mathcal{H}_{b,\Gamma}^{*, -s}$ be the dual space relative to L^2 , which is thus¹¹

$$Q_0H_b^{-s} + Q_+H_b^{-s} + Q_-H_b^{-s} + \tau^{1/2}H_b^{-s} + \hat{P}_0H_b^{-s} + H_b^{-s+1/2}.$$

Note that microlocally away from Γ , $\mathcal{H}_{b,\Gamma}^s$ is just the standard H_b^s space while $\mathcal{H}_{b,\Gamma}^{*, -s}$ is H_b^{-s} since at least one of Q_0, Q_\pm and τ is elliptic. Moreover, $\Psi_b^k(M) \ni A : \mathcal{H}_{b,\Gamma}^s \rightarrow \mathcal{H}_{b,\Gamma}^{s-k}$ is continuous since $[Q_+, A] \in \Psi_b^{k-1}(M)$ etc.; the analogous statement also holds for the dual spaces. Further, the last term in (3.7) can be replaced by $\|u\|_{H_b^{s-1}}^2$ as $i[Q_+, Q_-] = B^*B + R$, $B \in \Psi_b^{-1/2}(M)$, $R \in \Psi_b^{-2}(M)$, using the same argument as in the semiclassical setting (however, it cannot be dropped altogether unlike in the semiclassical setting!).

Remark 3.1. The notation $\mathcal{H}_{b,\Gamma}^s(M)$ is justified for the space is independent of the particular defining functions ϕ_\pm chosen; near Γ any other choice would replace ϕ_\pm by smooth non-degenerate linear combinations plus a multiple of τ and of \hat{p}_0 , denote these by $\check{\phi}_\pm$, and thus the corresponding \check{Q}_\pm can be expressed as

$$B_+Q_+ + B_-Q_- + B_\partial\tau + \hat{B}\hat{P}_0 + B_0Q_0 + R, \quad B_\pm, B_0, B_\partial, \hat{B} \in \Psi_b^0(M), \quad R \in \Psi_b^{-1}(M),$$

so the new norm can be controlled by the old norm, and conversely in view of the non-degeneracy.

Our result is then:

Theorem 3.2. *With $\mathcal{P}, \mathcal{H}_{b,\Gamma}^s, \mathcal{H}_{b,\Gamma}^{*,s}$ as above, for any neighborhood U of Γ and for any N there exist $B_0 \in \Psi_b^0(M)$ elliptic at Γ and $B_1, B_2 \in \Psi_b^0(M)$ with $\text{WF}'_b(B_j) \subset U$, $j = 0, 1, 2$, $\text{WF}'_b(B_2) \cap \Gamma_+ = \emptyset$ and $C > 0$ such that*

$$\|B_0u\|_{\mathcal{H}_{b,\Gamma}^s} \leq \|B_1\mathcal{P}u\|_{\mathcal{H}_{b,\Gamma}^{*,s-m+1}} + \|B_2u\|_{H_b^s} + C\|u\|_{H_b^{-N}}, \quad (3.8)$$

¹⁰Note that ${}^bT^*M$ is *not* a symplectic manifold (in a natural way) since the symplectic form on ${}^bT_{M^o}^*M$ does not extend smoothly to ${}^bT^*M$. Thus, the word ‘normally isotropic’ is not completely justified; we use it since it reflects that in the analogous semiclassical setting, see [13], the set Γ is symplectic, and the origin in the symplectic orthocomplement $(T_\alpha\Gamma)^\perp$ of $T_\alpha\Gamma$, which is also symplectic, is isotropic within $(T_\alpha\Gamma)^\perp$.

¹¹We refer to [9, Appendix A] for a general discussion of the underlying functional analysis. In particular, Lemma A.3 there essentially gives the density of $\dot{C}^\infty(M)$ in $\mathcal{H}_{b,\Gamma}^s(M)$: one can simply drop the subscript ‘e’ in the statement of that lemma to conclude that $H_b^\infty(M)$ (so in particular $H_b^s(M)$) is dense in $\mathcal{H}_{b,\Gamma}^s(M)$, and then the density of $\dot{C}^\infty(M)$ in $H_b^{s'}(M)$ for any s' completes the argument. The completeness of $\mathcal{H}_{b,\Gamma}^s(M)$ follows from the continuity of $\Psi_b^0(M)$ on $H_b^{s-1/2}(M)$.

i.e. if all the functions on the right hand side are in the indicated spaces: $B_1\mathcal{P}u \in \mathcal{H}_{b,\Gamma}^{*,s-m+1}$, etc., then $B_0u \in \mathcal{H}_{b,\Gamma}^s$, and the inequality holds.

The same conclusion also holds if we assume $\text{WF}'_b(B_2) \cap \Gamma_- = \emptyset$ instead of $\text{WF}'_b(B_2) \cap \Gamma_+ = \emptyset$.

Finally, if $r < 0$, then, with $\text{WF}'_b(B_2) \cap \Gamma_+ = \emptyset$, (3.8) becomes

$$\|B_0u\|_{H_b^{s,r}} \leq \|B_1\mathcal{P}u\|_{H_b^{s-m+1,r}} + \|B_2u\|_{H_b^{s,r}} + C\|u\|_{H_b^{-N,r}}, \quad (3.9)$$

while if $r > 0$, then, with $\text{WF}'_b(B_2) \cap \Gamma_- = \emptyset$,

$$\|B_0u\|_{H_b^{s,r}} \leq \|B_1\mathcal{P}u\|_{H_b^{s-m+1,r}} + \|B_2u\|_{H_b^{s,r}} + C\|u\|_{H_b^{-N,r}}, \quad (3.10)$$

Remark 3.3. Note that the weighted versions (3.9)-(3.10) use *standard* weighted b-Sobolev spaces; this corresponds to non-trapping semiclassical estimates if the subprincipal symbol has the correct, definite, sign at Γ .

Proof. We may assume that $U \subset U_0$ is disjoint from a neighborhood of $\text{WF}'_b(Q_0)$, and thus ignore Q_0 in the definition of $\mathcal{H}_{b,\Gamma}^s$ below.

We first prove that there exist B_0, B_1, B_2 as above and $B_3 \in \Psi_b^0(M)$ with $\text{WF}'_b(B_3) \subset U$ such that

$$\|B_0u\|_{\mathcal{H}_{b,\Gamma}^s} \leq \|B_1\mathcal{P}u\|_{\mathcal{H}_{b,\Gamma}^{*,s-m+1}} + \|B_2u\|_{H_b^s} + \|B_3u\|_{H_b^{s-1}} + C\|u\|_{H_b^{-N}}. \quad (3.11)$$

An iterative argument will then prove the theorem.

The proof is a straightforward modification of the construction in the semiclassical setting above, replacing ϕ_+^2 by $\phi_+^2 + M\tau$, $M > 0$ large, in accordance with (3.5).

We start by pointing out that for any $\tilde{B}_0 \in \Psi_b^0(M)$ and any $\tilde{B}_3 \in \Psi_b^0(M)$ elliptic on $\text{WF}'_b(\tilde{B}_0)$, we have

$$\|\hat{P}_0\tilde{B}_0u\|_{H_b^s} \leq C\|\tilde{B}_0\mathcal{P}u\|_{H_b^{s-m}} + C'\|\tilde{B}_3u\|_{H_b^{s-1}}, \quad (3.12)$$

by simply using that \hat{P}_0 is an elliptic multiple of \mathcal{P} modulo $\Psi_b^{-1}(M)$. Since $\|\tilde{B}_0\mathcal{P}u\|_{H_b^{s-m}} \leq C\|\tilde{B}_0\mathcal{P}u\|_{\mathcal{H}_{b,\Gamma}^{*,s-m}}$, the \hat{P}_0 contribution to $\|\tilde{B}_0u\|_{\mathcal{H}_{b,\Gamma}^s}$ in (3.11) is thus automatically controlled.

So let $\chi_0(t) = e^{-F/t}$ for $t > 0$, $\chi_0(t) = 0$ for $t \leq 0$, with $F > 0$ (large) to be specified, $\chi \in \mathcal{C}_c^\infty([0, \infty))$ be identically 1 near 0 with $\chi' \leq 0$, and indeed with $\chi'\chi = -\chi_1^2$, $\chi_1 \geq 0$, $\chi_1 \in \mathcal{C}_c^\infty([0, \infty))$, and let $\psi \in \mathcal{C}_c^\infty(\mathbb{R})$ be identically 1 near 0. As we use the Weyl quantization,¹² we write \mathcal{P} as the Weyl quantization of $p = p_0 + \tilde{\rho}p_1$, with $\tilde{\rho}p_1$ of order $m-1$. Let

$$a = \tilde{\rho}^{-s+(m-1)/2}\chi_0(\rho_+ - \phi_-^2 + \kappa)\chi(\rho_+)\psi(\tilde{\rho}^m p), \quad (3.13)$$

$\kappa > 0$ small. Notice that on $\text{supp } a$, if χ is supported in $[0, R]$,

$$\rho_+ \leq R, \quad \phi_-^2 \leq \rho_+ + \kappa = R + \kappa,$$

so a is localized near Γ if R and κ are taken sufficiently small. In particular, the argument of χ_0 is bounded above by $R + \kappa$, so given any $M_0 > 0$ one can take $F > 0$ large so that

$$\chi'_0\chi_0 - M_0\chi_0^2 = b^2\chi'_0\chi_0,$$

¹²Again, the Weyl quantization is irrelevant: If $A \in \Psi_b^m(X)$ and the principal symbol of A is real then the real part of the subprincipal symbol is defined independently of choices, which suffices below.

with $b \geq 1/2$, \mathcal{C}^∞ , on the range of the argument of χ_0 .

In fact, we also need to regularize, namely introduce

$$a_\epsilon = (1 + \epsilon \tilde{\rho}^{-1})^{-2} a, \quad \epsilon \in [0, 1], \quad (3.14)$$

which is a symbol of order $s - (m - 1)/2 - 2$ for $\epsilon > 0$, and is uniformly bounded in symbols of order $s - (m - 1)/2$ as ϵ varies in $[0, 1]$. In order to avoid more cumbersome notation below, we ignore the regularizer and work directly with a ; since the regularizer gives the same kind of contributions to the commutator as the weight $\tilde{\rho}^{-s+(m-1)/2}$, these contributions can be dominated in exactly the same way.

Then, with $p = p_0 + \tilde{\rho}p_1$ as above, $W = \tilde{\rho}^{m-2}\mathbf{H}_{\tilde{\rho}p_1}$, which is a smooth vector field near ${}^bS^*M$ as $\tilde{\rho}p_1$ is order $m - 1$, noting $W\tilde{\rho} = \tilde{\alpha}_1\tau\tilde{\rho}$ similarly to (3.6), and $W\tau = \alpha_{\partial,1}\tau$ by the tangency of W to $\tau = 0$,

$$\begin{aligned} \frac{1}{4}\mathbf{H}_p(a^2) &= -(-\hat{\rho}_+/2 + c_-^2\phi_-^2 + \nu_- \phi_- \hat{p}_0 - \tilde{\rho}\phi_+(W\phi_+) - \tilde{\rho}\mathbf{M}\alpha_{\partial,1}\tau + \tilde{\rho}\phi_-(W\phi_-)) \\ &\quad \times \tilde{\rho}^{-2s}(\chi_0\chi'_0)(\rho_+ - \phi_-^2 + \kappa)\chi(\rho_+)^2\psi(\tilde{\rho}^m p)^2 \\ &\quad + \frac{1}{4}(-2s + m - 1)\tilde{\rho}^{-2s}(\tilde{\alpha} + \tilde{\rho}\tilde{\alpha}_1)\tau\chi_0(\rho_+ - \phi_-^2 + \kappa)^2\chi(\rho_+)^2\psi(\tilde{\rho}^m p)^2 \\ &\quad + \frac{1}{2}\tilde{\rho}^{-2s}(\hat{\rho}_+ + \tilde{\rho}W\rho_+)(\chi'\chi)(\rho_+)\chi_0(\rho_+ - \phi_-^2 + \kappa)^2\psi(\tilde{\rho}^m p)^2 \\ &\quad + \frac{m}{2}(\tilde{\alpha} + \tilde{\rho}\tilde{\alpha}_1)\tilde{\rho}^{-2s}(\tilde{\rho}^m p)\tau\chi_0(\rho_+ - \phi_-^2 + \kappa)^2\chi(\rho_+)^2(\psi\psi')(\tilde{\rho}^m p). \end{aligned} \quad (3.15)$$

A key point is that the second term on the right hand side, given by the weight $\tilde{\rho}^{-2s+m-1}$ being differentiated, can be absorbed into the first by making $F > 0$ large so that $\hat{\rho}_+\chi'_0(\rho_+ - \phi_-^2 + \kappa)$ dominates

$$|-2s + m - 1||\tilde{\alpha}|\tau\chi_0(\rho_+ - \phi_-^2 + \kappa)$$

on $\text{supp } a$, which can be arranged as $|-2s + m - 1||\tilde{\alpha}|\tau$ is bounded by a sufficiently large multiple of $\hat{\rho}_+$ there. Thus,

$$\frac{1}{4}\mathbf{H}_p(a^2) = -c_+^2 a_+^2 - c_-^2 a_-^2 - a_\partial^2 + 2g_+ a_+ + 2g_- a_- + e + \tilde{e} + 2a_+ j_+ p + 2a_- j_- p \quad (3.16)$$

with

$$\begin{aligned} a_\pm &= \tilde{\rho}^{-s}\phi_\pm\sqrt{(\chi_0\chi'_0)(\rho_+ - \phi_-^2 + \kappa)\chi(\rho_+)\psi(\tilde{\rho}^m p)}, \\ a_\partial &= \tilde{\rho}^{-s}\tau^{1/2}\left(\mathbf{M}(c_\partial^2/2) - \beta_+\phi_+ - \tilde{\rho}\mathbf{M}a_{\partial,1}\right)(\chi_0\chi'_0)(\rho_+ - \phi_-^2 + \kappa) \\ &\quad - \frac{1}{4}(-2s + m - 1)(\tilde{\alpha} + \tilde{\rho}\tilde{\alpha}_1)\chi_0(\rho_+ - \phi_-^2 + \kappa)^2)^{1/2}\chi(\rho_+)\psi(\tilde{\rho}^m p), \\ g_\pm &= \pm\frac{1}{2}\tilde{\rho}^{-s+1}((W\phi_\pm) - \nu_\pm\tilde{\rho}^{m-1}p_1)\sqrt{(\chi_0\chi'_0)(\rho_+ - \phi_-^2 + \kappa)\chi(\rho_+)\psi(\tilde{\rho}^m p)}, \\ e &= -\frac{1}{2}\tilde{\rho}^{-2s}(\hat{\rho}_+ + \tilde{\rho}W\rho_+)\chi_1(\rho_+)^2\chi_0(\rho_+ - \phi_-^2 + \kappa)^2\psi(\tilde{\rho}^m p)^2, \\ \tilde{e} &= \frac{m}{2}\tilde{\rho}^{-2s}(\tilde{\rho}^m p)(\tilde{\alpha} + \tilde{\rho}\tilde{\alpha}_1)\tau\chi_0(\rho_+ - \phi_-^2 + \kappa)^2\chi(\rho_+)^2(\psi\psi')(\tilde{\rho}^m p), \\ j_\pm &= \pm\frac{1}{2}\nu_\pm\tilde{\rho}^{-s+m}\sqrt{(\chi_0\chi'_0)(\rho_+ - \phi_-^2 + \kappa)\chi(\rho_+)\psi(\tilde{\rho}^m p)}; \end{aligned}$$

the square root in a_∂ is that of a non-negative quantity and is \mathcal{C}^∞ for \mathbf{M} large (so that $\beta_+\phi_+$ can be absorbed into $\mathbf{M}(c_\partial^2/2)$) and F large (so that a small multiple of

χ'_0 can be used to dominate χ_0), as discussed earlier. Moreover,

$$\begin{aligned} \text{supp } e &\subset \text{supp } a, \quad \text{supp } e \cap \Gamma_+ = \emptyset, \\ \text{supp } \tilde{e} &\subset \text{supp } a, \quad \text{supp } \tilde{e} \cap \Sigma = \emptyset. \end{aligned}$$

This gives, with the various operators being Weyl quantizations of the corresponding lower case symbols,

$$\begin{aligned} \frac{i}{4}[\mathcal{P}, A^*A] &= -(C_+A_+)^*(C_+A_+) - (C_-A_-)^*(C_-A_-) - A_\partial^*A_\partial \\ &\quad + G_+^*A_+ + A_+^*G_+ + G_-^*A_- + A_-^*G_- \\ &\quad + E + \tilde{E} + A_+^*J_+\mathcal{P} + \mathcal{P}^*J_+^*A_+ + A_-^*J_-\mathcal{P} + \mathcal{P}^*J_-^*A_- + F \end{aligned} \quad (3.17)$$

where now $A \in \Psi_b^{s-(m-1)/2}(M)$, $A_\pm, A_\partial \in \Psi_b^s(M)$, $G_\pm \in \Psi_b^{s-1}(M)$, $E \in \Psi_b^{2s}(M)$, $\tilde{E} \in \Psi_b^{2s}(M)$, $J_\pm \in \Psi_b^{s-m}(M)$, $F \in \Psi_b^{2s-2}(M)$ with $\text{WF}'_b(F) \subset \text{supp } a$.

After this point the calculations repeat the semiclassical argument: First using $\mathcal{P} - \mathcal{P}^* \in \Psi_b^{m-2}(M)$,

$$\begin{aligned} &\|C_+A_+u\|^2 + \|C_-A_-u\|^2 + \|A_\partial u\|^2 \\ &\leq |\langle Eu, u \rangle| + |\langle \tilde{E}u, u \rangle| + |\langle A\mathcal{P}u, Au \rangle| + 2\|A_+u\|\|G_+u\| + 2\|A_-u\|\|G_-u\| \\ &\quad + 2|\langle J_+\mathcal{P}u, A_+u \rangle| + 2|\langle J_-\mathcal{P}u, A_-u \rangle| + C_1\|\tilde{F}_1u\|_{H_b^{s-1}}^2 + C_1\|u\|_{H_b^{-N}}^2, \end{aligned} \quad (3.18)$$

where we took $\tilde{F}_1 \in \Psi_b^0(M)$ elliptic on $\text{WF}'_b(F)$ and with $\text{WF}'_b(\tilde{F}_1)$ near Γ . Noting that $\text{WF}'_b(\tilde{E}) \cap \Sigma = \emptyset$, the elliptic estimates give

$$|\langle \tilde{E}u, u \rangle| \leq C\|B_1\mathcal{P}u\|_{H_b^{s-m}}^2 + C\|u\|_{H_b^{-N}}^2$$

if $B_1 \in \Psi_b^0(M)$ is elliptic on $\text{supp } \tilde{e}$. Let $\Lambda \in \Psi_b^{(m-1)/2}(M)$ be elliptic with real principal symbol λ , and let $\Lambda^- \in \Psi_b^{-(m-1)/2}(M)$ be a parametrrix for it so that $\Lambda\Lambda^- - \text{Id} = R_0 \in \Psi_b^{-\infty}(M)$. Then

$$\begin{aligned} |\langle A\mathcal{P}u, Au \rangle| &\leq |\langle \Lambda^-A\mathcal{P}u, \Lambda^*Au \rangle| + |\langle R_0A\mathcal{P}u, Au \rangle| \\ &\leq \frac{1}{2\epsilon}\|\Lambda^-A\mathcal{P}u\|_{\mathcal{H}_{b,\Gamma}^{s,0}}^2 + \frac{\epsilon}{2}\|\Lambda^*Au\|_{\mathcal{H}_{b,\Gamma}^0}^2 + C'\|u\|_{H_b^{-N}}^2 \end{aligned}$$

As $\Lambda^*A \in \Psi_b^s(M)$, for sufficiently small $\epsilon > 0$, $\frac{\epsilon}{2}\|\Lambda^*Au\|_{\mathcal{H}_{b,\Gamma}^0}^2$ can be absorbed into¹³ $\|C_+A_+u\|^2 + \|C_-A_-u\|^2 + \|A_\partial u\|^2$ plus $\|\tilde{B}_0\hat{\mathcal{P}}_0u\|_{H_b^s}^2$, and as discussed above, the latter already has the control required for (3.11). On the other hand, taking $B_1 \in \Psi_b^0(M)$ elliptic on $\text{WF}'_b(A)$, as $\Lambda^-A \in \Psi_b^{s-m+1}(M)$,

$$\|\Lambda^-A\mathcal{P}u\|_{\mathcal{H}_{b,\Gamma}^{s,0}}^2 \leq C''\|B_1\mathcal{P}u\|_{\mathcal{H}_{b,\Gamma}^{s-m+1}}^2 + C''\|u\|_{H_b^{-N}}^2.$$

¹³The point being that $A_+^*C_+^*C_+A_+ - \epsilon A^*\Lambda Q_+^*Q_+\Lambda^*A$ has principal symbol $c_+^2a_+^2 - \epsilon a^2\phi_+^2\lambda^2$ which can be written as the square of a real symbol for $\epsilon > 0$ small in view of the main difference in vanishing factors in the two terms being that χ'_0 in a_+^2 is replaced by χ_0 in a , and thus the corresponding operator can be expressed as $\tilde{C}^*\tilde{C}$ for suitable \tilde{C} , modulo an element of $\Psi_b^{2s-2}(M)$, with the latter contributing to the H_b^{s-1} error term on the right hand side of (3.11).

Similarly, to deal with the J_{\pm} terms on the right hand side of (3.18), one writes

$$\begin{aligned} |\langle J_{\pm} \mathcal{P}u, A_{\pm} u \rangle| &\leq \frac{1}{2\epsilon} \left(\|B_1 \mathcal{P}u\|_{H_b^{s-m}}^2 + C'' \|u\|_{H_b^{-N}}^2 \right) + \frac{\epsilon}{2} \|A_{\pm} u\|_{L^2}^2 \\ &\leq \frac{1}{2\epsilon} \left(\|B_1 \mathcal{P}u\|_{\mathcal{H}_{b,\Gamma}^{*,s-m}}^2 + C'' \|u\|_{H_b^{-N}}^2 \right) + \frac{\epsilon}{2} \|A_{\pm} u\|_{L^2}^2, \end{aligned}$$

while the G_{\pm} terms can be estimated by

$$\epsilon \|A_+ u\|^2 + \epsilon^{-1} \|G_+ u\|^2 + \epsilon \|A_- u\|^2 + \epsilon^{-1} \|G_- u\|^2,$$

and for $\epsilon > 0$ sufficiently small, the $\|A_{\pm} u\|^2$ terms in both cases can be absorbed into the left hand side of (3.18) while the G_{\pm} into the error term. This gives, with \tilde{F}_2 having properties as \tilde{F}_1 ,

$$\begin{aligned} &\|C_+ A_+ u\|^2 + \|C_- A_- u\|^2 + \|A_{\partial} u\|^2 \\ &\leq |\langle Eu, u \rangle| + C \|B_1 \mathcal{P}u\|_{\mathcal{H}_{b,\Gamma}^{*,s-m+1}}^2 + C_2 \|\tilde{F}_2 u\|_{H_b^{s-1}}^2 + C_2 \|u\|_{H_b^{-N}}^2. \end{aligned}$$

By the remark before the statement of the theorem, if $B_0 \in \Psi_b^0(M)$ is such that $\chi_0(\rho_+ - \phi_+^2 + \kappa)\chi(\rho_+)\psi(p) > 0$ on $\text{WF}'_b(B_0)$, $\|B_0 u\|_{H_b^{s-1/2}}^2$ can be added to the left hand side at the cost of changing the constant in front of $\|\tilde{F}_2 u\|_{H_b^{s-1}}^2 + \|u\|_{H_b^{-N}}^2$ on the right hand side. Taking such $B_0 \in \Psi_b^0(M)$, and B_1 elliptic on $\text{WF}'_b(A)$ as before, $B_2 \in \Psi_b^0(M)$ elliptic on $\text{WF}'_b(E)$ but with $\text{WF}'_b(B_2)$ disjoint from Γ_+ , we conclude that

$$\|B_0 u\|_{\mathcal{H}_{b,\Gamma}^s}^2 \leq C \|B_1 \mathcal{P}u\|_{\mathcal{H}_{b,\Gamma}^{*,s-m+1}}^2 + C \|B_2 u\|_{H_b^s}^2 + C \|\tilde{F}_2 u\|_{H_b^{s-1}}^2 + C \|u\|_{H_b^{-N}}^2,$$

proving (3.11), up to redefining B_j by multiplication by a positive constant. Recall that unless one makes sufficient a priori assumptions on the regularity of u , one actually needs to regularize, but as mentioned after (3.14), the regularizer is handled in exactly the same manner as the weight.

Now in general, with χ as before, but supported in $[0, 1]$ instead of $[0, R]$, let $\chi_R = \chi(\cdot/R)$ and write $a = a_{R,\kappa}$ to emphasize its dependence on these quantities. When R and κ are decreased, $\text{supp } a_{R,\kappa}$ also decreases in Σ in the strong sense that $0 < R < R'$ and $0 < \kappa < \kappa'$ imply that $a_{R',\kappa'}$ is elliptic on $\text{supp } a_{R,\kappa}$ within Σ , and indeed globally if the cutoff ψ is suitably adjusted as well. Thus, if $u \in H_b^{-N}$, say, one uses first (3.11) with $s = -N + 1$, and with B_j given by the proof above, so the $B_3 u$ term is a priori bounded, to conclude that $B_0 u \in \mathcal{H}_{b,\Gamma}^s$ and the estimate holds, so in particular, u is in $H_b^{-N+1/2}$ microlocally near Γ (concretely, on the elliptic set of B_0). Now one decreases κ and R by an arbitrarily small amount and applies (3.11) with $s = -N + 3/2$; the $B_3 u$ term is now a priori bounded by the microlocal membership of u in $H_b^{-N+1/2}$, and one concludes that $B_0 u \in \mathcal{H}_{b,\Gamma}^{-N+3/2}$, so in particular u is microlocally in H_b^{-N+1} . Proceeding inductively, one deduces the first statement of the theorem, (3.8).

If one reverses the role of Γ_+ and Γ_- in the statement of the theorem, one simply reverses the roles of $\rho_+ = \phi_+^2 + M\tau$ and ϕ_-^2 in the definition of a in (3.13). This reverses the signs of all terms on the right hand side of (3.15) whose sign mattered below, and thus the signs of the first three terms on the right hand side of (3.17), which then does not affect the rest of the argument.

In order to prove (3.9), one simply adds a factor τ^{-2r} to the definition of a in (3.13). This adds a factor τ^{-2r} to every term on the right hand side of (3.17), as

well as an additional term

$$\frac{r}{2}\tau^{-2r}\tilde{\rho}^{-2s}c_{\partial}^2\chi_0(\rho_+ - \phi_-^2 + \kappa)^2\chi(\rho_+)^2\psi(p)^2,$$

which for $r < 0$ has the same sign as the terms whose sign was used above, and indeed can be written as the negative of a square. Thus (3.16) becomes

$$\begin{aligned} \frac{1}{4}\mathbf{H}_p(a^2) &= -c_+^2a_+^2 - c_-^2a_-^2 - a_{\partial}^2 - a_r^2 \\ &\quad + 2g_+a_+ + 2g_-a_- + e + \tilde{e} + 2j_+a_+p + 2j_-a_-p \end{aligned} \quad (3.19)$$

with

$$a_r = \sqrt{\frac{-r}{2}}\tau^{-r}\tilde{\rho}^{-s}c_{\partial}\chi_0(\rho_+ - \phi_-^2 + \kappa)\chi(\rho_+)\psi(p),$$

and all other terms as above apart from the additional factor of τ^{-r} in the definition of a_{\pm} , etc. Since a_r is actually elliptic at Γ when $r \neq 0$, this proves the desired estimate (and one does not need to use the improved properties given by the Weyl calculus!).

When the role of Γ_+ and Γ_- is reversed, there is an overall sign change, and thus $r > 0$ gives the advantageous sign; the rest of the argument is unchanged. \square

Remark 3.4. As in the semiclassical setting, see Remark 2.1, the estimate (3.8) can be strengthened by adding the term $\|B_0\hat{P}_0u\|_{H_b^{s+1}}$ to the left hand side, which is controlled by elliptic regularity, likewise for (3.9)–(3.10). A more natural way of phrasing such an improvement is to use ‘coisotropic, normally isotropic’ spaces $\tilde{\mathcal{H}}_{b,\Gamma}^s$ and $\tilde{\mathcal{H}}_{b,\Gamma}^{*,s}$ in the estimate (3.8), where the squared norm on $\tilde{\mathcal{H}}_{b,\Gamma}^s$ is defined by

$$\|u\|_{\tilde{\mathcal{H}}_{b,\Gamma}^s}^2 = \|Q_0u\|_{H_b^s}^2 + \|Q_+u\|_{H_b^s}^2 + \|Q_-u\|_{H_b^s}^2 + \|\tau^{1/2}u\|_{H_b^s}^2 + \|\hat{P}_0u\|_{H_b^{s+1/2}}^2 + \|u\|_{H_b^{s-1/2}}^2,$$

i.e. strengthening the norm of \hat{P}_0u by a half, which strengthens the space and weakens its dual. To obtain the necessary elliptic estimate (3.12) with the strengthened norms on the terms involving \tilde{B}_0 , but keeping the norm on \tilde{B}_3u (which is required for the iterative argument at the end of the proof), one can choose \tilde{B}_0 with $\text{WF}'_b(I - \tilde{B}_0) \cap \Gamma = \emptyset$ so that \tilde{B}_3 can be chosen to be microsupported away from Γ , and thus $\|\tilde{B}_3u\|_{H_b^{s-1/2}} \leq C\|\tilde{B}_3u\|_{\tilde{\mathcal{H}}_{b,\Gamma}^{s-1/2}}$ is controlled using (3.8), with the norm on $B_1\mathcal{P}u$ being $\|B_1\mathcal{P}u\|_{\tilde{\mathcal{H}}_{b,\Gamma}^{*,s-m+1/2}} \leq C\|B_1\mathcal{P}u\|_{\tilde{\mathcal{H}}_{b,\Gamma}^{*,s-m+1}}$, and the error term being measured in $H_b^{s-3/2} \supset H_b^{s-1}$, as required.

REFERENCES

- [1] Jean-François Bony, Nicolas Burq, and Thierry Ramond. Minoration de la résolvante dans le cas captif. *Comptes Rendus Mathématique*, 348(23–24):1279–1282, 2010.
- [2] Kiril Datchev and Andrés Vasy. Gluing semiclassical resolvent estimates via propagation of singularities. *Int. Math. Res. Notices*, 2012(23):5409–5443, 2012.
- [3] Kiril Datchev and Andrés Vasy. Propagation through trapped sets and semiclassical resolvent estimates. *Annales de l’Institut Fourier*, 62(6): 2347–2377, 2012.
- [4] Kiril Datchev and Andrés Vasy. Semiclassical resolvent estimates at trapped sets. *Annales de l’Institut Fourier*, 62(6):2379–2384, 2012.
- [5] Semyon Dyatlov. Asymptotics of linear waves and resonances with applications to black holes. *Preprint, arXiv:1305.1723*, 2013.
- [6] Semyon Dyatlov. Spectral gaps for normally hyperbolic trapping. *Preprint, arXiv:1403.6401*, 2014.
- [7] Peter Hintz and Andrés Vasy. Global analysis of nonlinear wave equations on asymptotically Kerr-de Sitter spaces. *Preprint, arXiv:1404.1348*, 2014.

- [8] Richard B. Melrose. *The Atiyah-Patodi-Singer index theorem*, volume 4 of *Research Notes in Mathematics*. A K Peters Ltd., Wellesley, MA, 1993.
- [9] Richard B. Melrose, András Vasy, and Jared Wunsch. Diffraction of singularities for the wave equation on manifolds with corners. *Astérisque*, 351, vi+136pp, 2013.
- [10] Stéphane Nonnenmacher and Maciej Zworski. Decay of correlations for normally hyperbolic trapping. *Preprint*, *arXiv:1302.4483*, 2013.
- [11] András Vasy. Microlocal analysis of asymptotically hyperbolic and Kerr-de Sitter spaces. With an appendix by S. Dyatlov. *Inventiones Math.*, 194:381-513, 2013.
- [12] Jared Wunsch and Maciej Zworski. Erratum to ‘Resolvent estimates for normally hyperbolic trapped sets’. *Posted on http://www.math.northwestern.edu/~jwunsch/erratum_wz.pdf*.
- [13] Jared Wunsch and Maciej Zworski. Resolvent estimates for normally hyperbolic trapped sets. *Ann. Henri Poincaré*, 12(7):1349–1385, 2011.
- [14] Maciej Zworski. *Semiclassical Analysis*. Graduate studies in mathematics. American Mathematical Society, 2012.

DEPARTMENT OF MATHEMATICS, STANFORD UNIVERSITY, CA 94305-2125, USA

E-mail address: `phintz@math.stanford.edu`

E-mail address: `andras@math.stanford.edu`