

CYCLE INTEGRALS OF THE J-FUNCTION AND MOCK MODULAR FORMS

W. DUKE, Ö. IMAMOĞLU, AND Á. TÓTH

ABSTRACT. In this paper we construct certain mock modular forms of weight $1/2$ whose Fourier coefficients are given in terms of cycle integrals of the modular j -function. Their shadows are weakly holomorphic forms of weight $3/2$. These new mock modular forms occur as holomorphic parts of weakly harmonic Maass forms. We also construct a generalized mock modular form of weight $1/2$ having a real quadratic class number times a regulator as a Fourier coefficient. As an application of these forms we study holomorphic modular integrals of weight 2 whose rational period functions have poles at certain real quadratic integers. The Fourier coefficients of these modular integrals are given in terms of cycle integrals of modular functions. Such a modular integral can be interpreted in terms of a Shimura-type lift of a mock modular form of weight $1/2$ and yields a real quadratic analogue of a Borcherds product.

1. INTRODUCTION

Mock modular forms, especially those of weight $1/2$, have attracted much attention recently. This is mostly due to the discovery of Zwegers [51, 52] that Ramanujan's mock theta functions can be completed to become modular by the addition of a certain non-holomorphic function on the upper half plane \mathcal{H} . This complement is associated to a modular form of weight $3/2$, the *shadow* of the mock theta function. Consider, for example, the q -series

$$f(\tau) = q^{-1/24} \sum_{n \geq 0} \frac{q^{n^2}}{(1+q)^2 \cdots (1+q^n)^2} \quad (q = e(\tau) = e^{2\pi i \tau}, \tau \in \mathcal{H}).$$

Up to the factor $q^{-1/24}$ this is one of Ramanujan's original mock theta functions. The shadow of f is the weight $3/2$ cusp form (a unary theta series)

$$g(\tau) = \sum_{n \in 1+6\mathbb{Z}} n q^{n^2/24};$$

it is proved in [52] that the completion

$$\hat{f}(\tau) = f(\tau) + g^*(\tau)$$

transforms like a modular form of weight $1/2$ for $\Gamma(2)$, the well known congruence subgroup of $\Gamma = \mathrm{PSL}(2, \mathbb{Z})$, when

$$g^*(\tau) = \sum_{n \in 1+6\mathbb{Z}} \mathrm{sgn}(n) \beta\left(\frac{n^2 y}{6}\right) q^{-n^2/24} \quad (y = \mathrm{Im} \tau).$$

Here $\beta(x)$ is defined for $x > 0$ in terms of the complementary error function and the standard incomplete gamma function by

$$(1.1) \quad \beta(x) = \mathrm{erfc}(\sqrt{\pi x}) = \frac{1}{\sqrt{\pi}} \Gamma\left(\frac{1}{2}, \pi x\right), \quad \text{where} \quad \Gamma(s, x) = \int_x^\infty t^{s-1} e^{-t} dt.$$

W. Duke is supported by NSF grant DMS-0355564.

Á. Tóth is supported by OTKA grant 49693.

Observe that the Fourier expansion of the non-holomorphic “Eichler integral” $g^*(\tau)$ mirrors that of $g(\tau)$. In addition to leading to a number of new results about mock theta functions, the work of Zwegers has stimulated the study of other kinds of mock modular forms as well (see [38] and [50] for surveys on some of these developments).

In this paper we will consider mock modular forms of weight $1/2$ for $\Gamma_0(4)$. In some sense this is the simplest case, but has not been treated before because the associated shadows, if not zero, cannot be cusp forms. We will show that they are nevertheless quite interesting, and have remarkable connections with cycle integrals of the modular j -function and modular integrals having rational period functions. First let us define mock modular forms precisely in this context. Let

$$\theta(\tau) = 1 + 2q + 2q^4 + 2q^9 + 2q^{16} + \dots$$

be the Jacobi theta series, which is a modular form of weight $1/2$ for $\Gamma_0(4)$. Set

$$(1.2) \quad j(\gamma, \tau) = \theta(\gamma\tau)/\theta(\tau) \quad \text{for } \gamma \in \Gamma_0(4).$$

For $k \in 1/2 + \mathbb{Z}$ say that f defined on \mathcal{H} has weight k for $\Gamma_0(4)$ (or simply has weight k) if

$$f(\gamma\tau) = j(\gamma, \tau)^{2k} f(\tau) \quad \text{for all } \gamma \in \Gamma_0(4).$$

Let $M_k^!$ be the space comprising functions holomorphic on \mathcal{H} of weight k for $\Gamma_0(4)$ whose Fourier coefficients $a(n)$ in the expansion $f(\tau) = \sum_n a(n)q^n$ are supported on integers $n \geq -\infty$ with $(-1)^{k-1/2}n \equiv 0, 1 \pmod{4}$.

Specializing now to the case of weight $1/2$, let $\mathcal{E}(z)$ be the entire function given by any of the following formulas

$$(1.3) \quad \mathcal{E}(z) = \int_0^1 e^{-\pi zu^2} du = \frac{\operatorname{erf}(\sqrt{\pi z})}{2\sqrt{z}} = \sum_{n=0}^{\infty} \frac{(-\pi z)^n}{(2n+1)n!}.$$

For any $g(\tau) = \sum_n b_n q^n \in M_{3/2}^!$ we define the non-holomorphic Eichler integral of g by

$$(1.4) \quad g^*(\tau) = -4\sqrt{y} \sum_{n \leq 0} b_n \mathcal{E}(4ny) q^{-n} + \sum_{n > 0} \frac{b_n}{\sqrt{n}} \beta(4ny) q^{-n}.$$

Let $f(\tau) = \sum_n a_n q^n$ be holomorphic on \mathcal{H} and such that its coefficients a_n are supported on integers $-\infty < n \equiv 0, 1 \pmod{4}$. We will say that $f(\tau)$ is a mock modular form of weight $1/2$ for $\Gamma_0(4)$ if there exists a $g \in M_{3/2}^!$, its shadow, so that

$$\hat{f}(\tau) = f(\tau) + g^*(\tau)$$

has weight $1/2$ for $\Gamma_0(4)$. Denote by $\mathbb{M}_{1/2}$ the space of all mock modular forms of weight $1/2$ for $\Gamma_0(4)$. Obviously $M_{1/2}^! \subset \mathbb{M}_{1/2}$ but it is not at all clear that there are any non-modular mock modular forms.

We will show that they do exist and that they are related to the work of Borcherds and Zagier on traces of singular moduli of the classical j -function

$$j(\tau) = q^{-1} + 744 + 196884q + \dots$$

It is well-known and easily shown that $\mathbb{C}[j]$, has a unique basis $\{j_m\}_{m \geq 0}$ whose members are of the form $j_m(\tau) = q^{-m} + O(q)$. For example

$$(1.5) \quad j_0 = 1, \quad j_1 = j - 744, \quad j_2 = j^2 - 1488j + 159768, \dots$$

Here $j_1(\tau)$ is the normalized Hauptmodule for Γ . In this paper, unless otherwise specified, d is assumed to be an integer $d \equiv 0, 1 \pmod{4}$ and is called a discriminant if $d \neq 0$. For each discriminant d let \mathcal{Q}_d be the set of integral binary quadratic forms of discriminant d that are

positive definite if $d < 0$. The forms are acted on as usual by Γ , resulting in finitely many classes $\Gamma \backslash \mathcal{Q}_d$. Let Γ_Q be the group of automorphs of Q (see section 3 for more details).

Suppose that $d < 0$. For $Q \in \mathcal{Q}_d$ and τ_Q a root of Q in \mathcal{H} , the numbers $j_1(\tau_Q)$ are known by the classical theory of complex multiplication to form a $\text{Gal}(\bar{\mathbb{Q}}/\mathbb{Q})$ -invariant set of algebraic integers, so that their weighted sum

$$(1.6) \quad \text{Tr}_d(j_1) = \sum_{Q \in \Gamma \backslash \mathcal{Q}_d} |\Gamma_Q|^{-1} j_1(\tau_Q)$$

is an integer. A beautiful theorem of Zagier [49] asserts that these integers give the Fourier coefficients of a weight $3/2$ weakly holomorphic form in $\mathbf{T}_-(\tau) \in M_{3/2}^1$:

$$(1.7) \quad \begin{aligned} \mathbf{T}_-(\tau) &= -q^{-1} + 2 + \sum_{d \leq 0} \text{Tr}_d(j_1)(d) q^{|d|} \\ &= -q^{-1} + 2 - 248 q^3 + 492 q^4 - 4119 q^7 + 7256 q^8 + \cdots \end{aligned}$$

A natural question is whether one can give a similar statement for the numbers $\text{Tr}_d(j_1)$ defined for non-square $d > 0$ by

$$(1.8) \quad \text{Tr}_d(j_1) = \frac{1}{2\pi} \sum_{Q \in \Gamma \backslash \mathcal{Q}_d} \int_{C_Q} j_1(\tau) \frac{d\tau}{Q(\tau, 1)}.$$

Here C_Q is any smooth curve from any $z \in \mathcal{H}$ to $g_Q z$, where g_Q is a certain distinguished generator (see (3.1)) of the infinite cyclic group Γ_Q of automorphs of Q . Note: $\text{Tr}_d(j_1)$ is well-defined. We will see that the generating function

$$(1.9) \quad \mathbf{T}_+(\tau) = \sum_{d > 0} \text{Tr}_d(j_1) q^d$$

(with a suitable definition of $\text{Tr}_d(j_1)$ when d is a perfect square) defines a mock modular form of weight $1/2$ for $\Gamma_0(4)$ with shadow $\mathbf{T}_-(\tau)$ from (1.7).

Theorem 1. *The function $\widehat{\mathbf{T}}_+(\tau)$ on \mathcal{H} defined by*

$$\begin{aligned} \widehat{\mathbf{T}}_+(\tau) &= \mathbf{T}_+(\tau) + \mathbf{T}_-(\tau) \\ &= \sum_{d > 0} \text{Tr}_d(j_1) q^d + 4\sqrt{y} \mathcal{E}(-4y) q - 8\sqrt{y} + \sum_{d < 0} \frac{\text{Tr}_d(j_1)}{\sqrt{|d|}} \beta(4|d| y) q^d \end{aligned}$$

has weight $1/2$ for $\Gamma_0(4)$.

Zagier [49] showed that $g_1(\tau) = \mathbf{T}_-(\tau)$ from (1.7) is the first member of a basis $\{g_d\}_{0 < d \equiv 0, 1 \pmod{4}}$ for $M_{3/2}^1$, where for each $d > 0$ the function $g_d(\tau)$ is uniquely determined by having a q -expansion of the form¹

$$(1.10) \quad g_d(\tau) = -q^{-d} + \sum_{\substack{n \leq 0 \\ n \equiv 0, 1 \pmod{4}}} a(d, n) q^{|n|}.$$

We define $a(d, n) = 0$ unless $d, n \equiv 0, 1 \pmod{4}$. For $d \leq 0$ consider the “dual” form

$$(1.11) \quad f_d(\tau) = q^d + \sum_{n > 0} a(n, d) q^n.$$

¹This is the negative of the $g_d(\tau)$ defined in [49].

As was shown in [49], the set $\{f_d\}_{d \leq 0}$ coincides with the basis given by Borcherds [2] for $M_{1/2}^!$. Thus $f_0(\tau) = \theta(\tau)$ and the first few terms of the next function are

$$f_{-3}(\tau) = q^{-3} - 248q + 26752q^4 - 85995q^5 + 1707264q^8 + \dots$$

We will show that Borcherds' basis extends naturally to a basis for $\mathbb{M}_{1/2}$. The construction of this extension relies heavily on the spectral theory of Maass forms.

Theorem 2. *For each $d > 0$ there is a unique mock modular form $f_d(\tau) \in \mathbb{M}_{1/2}$ with shadow $g_d(\tau)$ having a Fourier expansion of the form*

$$(1.12) \quad f_d(\tau) = \sum_{n>0} a(n, d)q^n.$$

These Fourier coefficients $a(n, d)$ satisfy $a(n, d) = a(d, n)$. The set $\{f_d\}_{d \equiv 0, 1 \pmod{4}}$ gives a basis for $\mathbb{M}_{1/2}$.

We have thus defined $a(n, d)$ for all d, n with $n > 0$. We use them to evaluate certain twisted traces, which we now define. Suppose that $D > 0$ is a fundamental discriminant. There is a function $\chi_D : \mathcal{Q}_{dD} \rightarrow \{-1, 1\}$ defined below in (3.2) that restricts to a real character (a genus character) on the group of primitive classes and can be used to define a general twisted trace for dD not a square by

$$(1.13) \quad \text{Tr}_{d,D}(j_m) = \begin{cases} \frac{1}{\sqrt{D}} \sum \chi(Q) |\Gamma_Q|^{-1} j_m(\tau_Q), & \text{if } dD < 0; \\ \frac{1}{2\pi} \sum \chi(Q) \int_{C_Q} j_m(\tau) \frac{d\tau}{Q(\tau, 1)}, & \text{if } dD > 0, \end{cases}$$

each sum being over $Q \in \Gamma \backslash \mathcal{Q}_{dD}$. We have the following evaluation, which generalizes a well-known result of Zagier [49, (25)] to include positive d .

Theorem 3. *Let $a(n, d)$ be the mock modular coefficients defined in (1.11) and (1.12). Suppose that $m \geq 1$. For $d \equiv 0, 1 \pmod{4}$ and fundamental $D > 0$ with dD not a square we have*

$$(1.14) \quad \text{Tr}_{d,D}(j_m) = \sum_{n|m} \left(\frac{D}{m/n} \right) n a(n^2 D, d).$$

Together with Theorem 2, Theorem 3 implies Theorem 1 after we define $\text{Tr}_d(j_1)$ to be equal to $a(d, 1)$ when d is a perfect square. The proof we give uses Poincaré series and a Kloosterman sum identity that generalizes a well-known result of Salié. In particular, for non-square dD with $D > 0$, Theorem 3 gives

$$(1.15) \quad a(D, d) = \begin{cases} \frac{1}{\sqrt{D}} \sum \chi(Q) |\Gamma_Q|^{-1} j_1(\tau_Q), & \text{if } dD < 0; \\ \frac{1}{2\pi} \sum \chi(Q) \int_{C_Q} j_1(\tau) \frac{d\tau}{Q(\tau, 1)}, & \text{if } dD > 0, \end{cases}$$

where each sum is over $Q \in \Gamma \backslash \mathcal{Q}_{dD}$.

Concerning the case $m = 0$, there exists an interesting “second order” mock modular form $\mathbf{Z}_+(\tau)$ of weight $1/2$ that is almost, but not quite, in $\mathbb{M}_{1/2}$ with Fourier expansion

$$(1.16) \quad \mathbf{Z}_+(\tau) = \sum_{d>0} \text{Tr}_d(1) q^d.$$

Here $\text{Tr}_d(1)$ must be defined suitably for square d while for $d > 1$ a fundamental discriminant we have

$$\text{Tr}_d(1) = \pi^{-1} d^{-1/2} h(d) \log \epsilon_d,$$

where $h(d)$ is the narrow class number of $\mathbb{Q}(\sqrt{d})$ and ϵ_d is its smallest unit > 1 of norm 1. A (generalized) shadow of $\mathbf{Z}_+(\tau)$ is the completion of the mock modular form $\mathbf{Z}_-(\tau)$ of weight

3/2 with shadow $\theta(\tau)$ discovered by Zagier in 1975 [47] (see also [20]) whose Fourier expansion is

$$(1.17) \quad \mathbf{Z}_-(\tau) = \sum_{d \leq 0} \text{Tr}_d(1) q^{|d|}.$$

Here for any $d \leq 0$ we have that $\text{Tr}_d(1) = H(|d|)$, the usual Hurwitz class number, whose first few values are given by

$$H(0) = -\frac{1}{12}, \quad H(3) = \frac{1}{3}, \quad H(4) = \frac{1}{2}, \quad H(7) = 1, \quad \dots$$

The completion of $\mathbf{Z}_-(\tau)$, which has weight 3/2 for $\Gamma_0(4)$ is given by

$$(1.18) \quad \widehat{\mathbf{Z}}_-(\tau) = \mathbf{Z}_-(\tau) + \frac{1}{16\pi} \sum_{n \in \mathbb{Z}} \Gamma(-\frac{1}{2}, 4\pi n^2 y) q^{-n^2}.$$

Define for $y > 0$ the special function

$$\alpha(y) = \frac{\sqrt{y}}{4\pi} \int_0^\infty t^{-1/2} \log(1+t) e^{-\pi y t} dt.$$

The next result shows that $\mathbf{Z}_+(\tau)$ from (1.16) has $\widehat{\mathbf{Z}}_-(\tau)$ as a generalized shadow (to be made precise later.)

Theorem 4. *The function $\widehat{\mathbf{Z}}_+(\tau)$ whose Fourier expansion is given by*

$$(1.19) \quad \widehat{\mathbf{Z}}_+(\tau) = \sum_{d > 0} \text{Tr}_d(1) q^d + \frac{\sqrt{y}}{3} + \sum_{d < 0} \frac{\text{Tr}_d(1)}{\sqrt{|d|}} \beta(4|d|y) q^d + \sum_{n \neq 0} \alpha(4n^2 y) q^{n^2} - \frac{1}{4\pi} \log y,$$

has weight 1/2 for $\Gamma_0(4)$.

The automorphic nature of $\widehat{\mathbf{Z}}_+(\tau)$ gives some reason to hope that there might be a connection between the cycle integrals of j and abelian extensions of real quadratic fields. So far this hope has not been realized.

Finally, there is an unexpected connection between mock modular forms of weight 1/2 and modular integrals having rational period functions. Define for each $d \equiv 0, 1 \pmod{4}$

$$(1.20) \quad F_d(\tau) = -\text{Tr}_d(1) - \sum_{m \geq 1} \left(\sum_{n|m} n a(n^2, d) \right) q^m.$$

Note that $F_d(\tau)$ is the derivative of the formal Shimura lift of f_d . When $d < 0$ Borcherds showed that F_d is a meromorphic modular form of weight 2 for Γ having a simple pole with residue $|\Gamma_Q|^{-1}$ at each point $\tau_Q \in \mathcal{H}$ of discriminant d . Thus one has corresponding properties of the infinite product

$$q^{-\text{Tr}_d(1)} \prod_{m \geq 1} (1 - q^m)^{a(m^2, d)}.$$

In case $d = 0$ one finds that this product is $\Delta(\tau)^{1/12}$, and we have that

$$F_0(\tau) = \frac{1}{12} - 2 \sum_{n \geq 1} \sigma(n) q^n = \frac{1}{12} E_2(\tau).$$

This is a holomorphic modular integral of weight 2 with a rational period function:

$$F_0(\tau) - \tau^{-2} F_0(-\frac{1}{\tau}) = -\frac{1}{2\pi i} \tau^{-1}.$$

Theorem 5. *For each $d > 0$ not a square the function F_d defined in (1.20) is a holomorphic modular integral of weight 2 with a rational period function:*

$$(1.21) \quad F_d(\tau) - \tau^{-2}F_d(-\frac{1}{\tau}) = \frac{1}{\pi} \sum_{\substack{c < 0 < a \\ b^2 - 4ac = d}} (a\tau^2 + b\tau + c)^{-1}.$$

The Fourier expansion of $F_d(\tau)$ can be expressed in the form

$$F_d(\tau) = - \sum_{m \geq 0} \text{Tr}_d(j_m) q^m.$$

Note that the period function has simple poles at certain real quadratic integers of discriminant d , in analogy to the behavior of $F_d(\tau)$ when $d < 0$. The existence of a holomorphic F satisfying (1.21) with growth conditions was proved by Knopp [28], [29]. He used a certain Poincaré series built out of cocycles, which was used earlier by Eichler [13], to construct it. However, it appears to be very difficult to compute F explicitly from this construction. At the end of their paper [8], Choie and Zagier raised the problem of explicit construction of a modular integral with a given rational period function. Parson [39] gave a more direct construction in weights $k > 2$ using series of the form

$$\sum_{a > 0} (a\tau^2 + b\tau + c)^{-k/2},$$

which are partial versions of certain hyperbolic Poincaré series studied by Zagier, but they do not converge when $k = 2$. In any case, the expression of the Fourier coefficients as sums of cycle integrals is not immediate from this construction, although it is possible to deduce such expressions this way, at least in higher weights, using methods from this paper. For the rational period functions that occur in (1.21) the modular integral given by $F_d(\tau)$ also gives a real quadratic analogue of (the logarithmic derivative of) the Borcherds product.

TABLE 1. Traces for $d < 0$

| m | $\text{Tr}_{-3}(j_m)$ | $\text{Tr}_{-4}(j_m)$ | $\text{Tr}_{-7}(j_m)$ | $\text{Tr}_{-8}(j_m)$ |
|---|-----------------------|-----------------------|-----------------------|-----------------------|
| 0 | 1/3 | 1/2 | 1 | 1 |
| 1 | -248 | 492 | -4119 | 7256 |
| 2 | 53256 | 287244 | 16572393 | 52255768 |
| 3 | -12288992 | 153540528 | -67515202851 | 377674781024 |

TABLE 2. Traces for $d > 0$

| m | $\text{Tr}_5(j_m)$ | $\text{Tr}_8(j_m)$ | $\text{Tr}_{12}(j_m)$ | $\text{Tr}_{13}(j_m)$ | $\text{Tr}_{17}(j_m)$ | $\text{Tr}_{20}(j_m)$ |
|---|--------------------|--------------------|-----------------------|-----------------------|-----------------------|-----------------------|
| 0 | 0.13700 | 0.19837 | 0.24202 | 0.21095 | 0.32343 | 0.33097 |
| 1 | -5.16163 | -6.76613 | -8.27912 | -6.49263 | -10.65828 | -8.36253 |
| 2 | -11.56343 | -17.92434 | -21.49601 | -19.16428 | -28.46829 | -23.93151 |
| 3 | -14.31225 | -19.50182 | -24.84575 | -21.99742 | -34.17310 | -29.69296 |

It is interesting to examine numerical values of the traces $\text{Tr}_d(j_m)$. We remark that for $d > 1$ fundamental we have the identity

$$(1.22) \quad \text{Tr}_d(j_m) = \sum_{n|m} \left(\frac{d}{m/n}\right) n^{-1} \text{Tr}_{n^2d}(j_1).$$

This result is a consequence of a general identity on Hecke operators proved in [48]. As a numerical check, the reader can verify that the identity (1.22) holds for $d = 5$, $D = 1$ and $m = 2$. For $d = 20$ there are two classes, represented by $[1, 4, -1]$ and the non-primitive form $[2, 2, -2]$, and (1.22) amounts to the curious identity

$$(1.23) \quad \int_{C_{[1,4,-1]}} j_1(\tau) \frac{2d\tau}{\tau^2+4\tau-1} = \int_{C_{[1,1,-1]}} j_2(\tau) \frac{d\tau}{\tau^2+\tau-1}.$$

In general, a comparison shows that the traces for positive discriminants are generally much smaller than those for negative discriminants and appear to exhibit some regular growth behavior in both d and m . For non-square $d > 1$ and $m \in \mathbb{Z}^+$ we will see that we have the conditionally (and slowly) convergent expansion

$$(1.24) \quad \mathrm{Tr}_d(j_m) = -24\sigma_1(m)\mathrm{Tr}_d(1) + d^{-1/2} \sum_{0 < c \equiv 0(4)} S_m(d; c) \sin\left(\frac{4\pi m\sqrt{d}}{c}\right),$$

where

$$S_m(d; c) = \sum_{b^2 \equiv d \pmod{c}} e\left(\frac{2mb}{c}\right).$$

It is natural to expect that the first term dominates as either m or d gets large, which would in particular imply that as $d \rightarrow \infty$ through non-squares we have the asymptotic formula

$$(1.25) \quad \mathrm{Tr}_d(j_m) \sim -24\sigma_1(m) \mathrm{Tr}_d(1).$$

Such a result is known for negative discriminants (see [9]) except that then there is a large main term that grows like $e^{\pi m \sqrt{|d|}}$. If true, this asymptotic would indicate that there is tremendous cancellation in the integrals of j over the cycles when d is large, since $\mathrm{Tr}_d(1) \ll_{\epsilon} d^{\epsilon}$ and j_m has exponential growth in the cusp.

After seeing our paper, Y. Manin and D. Zagier told us that M. Kaneko also performed calculations of cycle integrals of the j function. Kaneko kindly sent us his paper [26], in which he calculated the individual cycle integrals numerically and observed some interesting behaviour in the distribution of their values. This behavior appears to us to be consistent with the conjectured asymptotic given above.

Finally, we remark that there is an obvious similarity between Theorem 3 and the formula of Katok and Sarnak [27] for the traces of a Maass cusp form. The main difference is that there are no Hecke eigenforms in our setting. Nevertheless, the method we use to prove Theorem 3, which involves Poincaré series and identities between Kloosterman sums, can be applied to give another proof of the Katok-Sarnak formula. It would be interesting to approach our results using a regularized theta lift. (See [16], [5] for the use of regularized theta lifts in the negative discriminants case.) Another interesting problem is to understand the nature of the traces on square discriminants.

Acknowledgements: We thank D. Zagier and the referee for numerous and extremely helpful comments on earlier versions of this paper. Duke and Tóth gratefully acknowledge the generous support of the Forschungsinstitut für Mathematik at ETH Zürich for this research. They also thank Marc Burger and the staff of FIM for providing a stimulating and comfortable atmosphere during their visits to Zürich.

2. WEAKLY HARMONIC MODULAR FORMS

We begin by proving Theorem 2 using the theory of weakly harmonic forms. Set $k \in 1/2 + \mathbb{Z}$. If f of weight k for $\Gamma_0(4)$ is smooth, for example, it will have a Fourier expansion in each cusp. For the cusp at $i\infty$ we have *the* Fourier expansion

$$(2.1) \quad f(\tau) = \sum_n a(n; y) e(nx)$$

which, if f is holomorphic, has $a(n; y) = a(n) e(niy)$. Set

$$(2.2) \quad f^e(\tau) = \sum_{n \equiv 0(2)} a(n; \frac{y}{4}) e(\frac{nx}{4}) \quad \text{and} \quad f^o(\tau) = \sum_{n \equiv 1(2)} a(n; \frac{y}{4}) e(\frac{n}{8}) e(\frac{nx}{4}).$$

Suppose that the Fourier coefficients $a(n; y)$ satisfy the *plus space condition*, meaning that they vanish unless $(-1)^{k-1/2} n \equiv 0, 1 \pmod{4}$. An easy extension of arguments given in [33, p.190] shows that such an f satisfies

$$(2.3) \quad \left(\frac{2\tau}{i}\right)^{-k} f\left(-\frac{1}{4\tau}\right) = \alpha f^e(\tau) \quad \text{and} \quad \left(\frac{2\tau+1}{i}\right)^{-k} f\left(\frac{\tau}{2\tau+1}\right) = \alpha f^o(\tau)$$

where

$$\alpha = (-1)^{\lfloor \frac{2k+1}{4} \rfloor} 2^{-k+\frac{1}{2}}.$$

In particular, the behavior of such an f at the cusps 0 and $1/2$ is determined by that at $i\infty$. Thus to check that such a form is weakly holomorphic, meaning it is holomorphic on \mathcal{H} and meromorphic in the cusps, one only needs look at the Fourier expansion at $i\infty$, as we have done in the Introduction. As there, let $M_k^!$ denote the space of all such forms. Let $M_k^+ \subset M_k^!$ denote the subspace of holomorphic forms (having no pole in the cusps) and $S_k^+ \subset M_k^+$ the subspace of cusp forms (having zeros there).

Consider the Maass-type differential operator ξ_k defined for any $k \in \mathbb{R}$ through its action on a differentiable function f on \mathcal{H} by

$$\xi_k(f) = 2iy^k \frac{\partial f}{\partial \bar{\tau}}.$$

This operator is studied in some detail in [6]. It is easily checked that

$$\xi_k((\gamma\tau + \delta)^{-k} f(g\tau)) = (\gamma\tau + \delta)^{k-2} (\xi_k f)(g\tau)$$

for any $g \in \text{PSL}(2, \mathbb{R})$. Thus if $f(\tau)$ has weight k for $\Gamma_0(4)$ then $\xi_k f$ has weight $2 - k$ and $\xi_k f = 0$ if and only if f is holomorphic. Also ξ_k preserves the plus space condition. The weight k Laplacian can be conveniently defined by

$$(2.4) \quad \Delta_k = -\xi_{2-k} \circ \xi_k.$$

Specializing now to $k = 1/2$, suppose that h is a real analytic function on \mathcal{H} of weight $1/2$ for $\Gamma_0(4)$ that is harmonic on \mathcal{H} in the sense that

$$(2.5) \quad \Delta_{1/2} h = 0.$$

By separation of variables every such h has a (unique) Fourier expansion in the cusp at ∞ of the form

$$(2.6) \quad h(\tau) = \sum_n b(n) \mathcal{M}_n(y) e(nx) + \sum_n a(n) \mathcal{W}_n(y) e(nx).$$

The functions $\mathcal{W}_n(y)$ and $\mathcal{M}_n(y)$ in the Fourier expansion (2.6) are defined in terms of the functions $\beta(x)$ and $\mathcal{E}(z)$ from (1.1) and (1.3) by

$$(2.7) \quad \mathcal{W}_n(y) = e^{-2\pi ny} \begin{cases} |n|^{-\frac{1}{2}} \beta(4|n|y) & \text{if } n < 0, \\ -4y^{\frac{1}{2}} & \text{if } n = 0, \\ n^{-\frac{1}{2}} & \text{if } n > 0, \end{cases}$$

$$(2.8) \quad \mathcal{M}_n(y) = e^{-2\pi ny} \begin{cases} 1 - \beta(4|n|y) & \text{if } n < 0, \\ 1 & \text{if } n = 0, \\ 4(ny)^{\frac{1}{2}} \mathcal{E}(-4ny) & \text{if } n > 0. \end{cases}$$

We remark that $\mathcal{W}_n(y)$ and $\mathcal{M}_n(y)$ are special cases of Whittaker functions, (see (2.16)) and we use the notation \mathcal{W} and \mathcal{M} to suggest this relation. More importantly, the definitions (2.7) and (2.8) make possible the complete symmetry of the Fourier coefficients of the basis to be given in the next Proposition. It becomes clear after working with them that one can define the normalization for the Fourier coefficients in different reasonable ways, each with advantages and disadvantages. Note that the function $\mathcal{W}_n(y)$ is exponentially decaying while $\mathcal{M}_n(y)$ is exponentially growing in y (see (A.4)).

Let $H_{1/2}^1$ denote the space of all real analytic functions on \mathcal{H} of weight $1/2$ for $\Gamma_0(4)$ that satisfy (2.5), whose Fourier coefficients at ∞ are supported on integers n with $n \equiv 0, 1 \pmod{4}$ and that have only finitely many non-zero coefficients $b(n)$. As before this is enough to control bad behavior in the other cusps. We will call such an $h \in H_{1/2}^1$ weakly harmonic.² This space was identified by Bruinier and Funke [5] as being interesting arithmetically. It follows easily from its general properties that $\xi_{1/2}$ maps $H_{1/2}^1$ to $M_{3/2}^1$ with kernel $M_{1/2}^1$. This is also directly visible after a calculation from (2.8) and (2.7) yields the formulas

$$(2.9) \quad \xi_{1/2}(\mathcal{M}_n(y)e(nx)) = 2|n|^{\frac{1}{2}}q^{-n} \quad \xi_{1/2}(\mathcal{W}_n(y)e(nx)) = \begin{cases} 0 & \text{if } n > 0 \\ -2q^{|n|} & \text{if } n \leq 0. \end{cases}$$

Given h in (2.6) with $b(n) = 0$ for all n , we infer that $\xi_{1/2}h \in M_{3/2}^+ = \{0\}$. This implies that $h \in S_{1/2}^+ = \{0\}$, and proves the following uniqueness result.

Lemma 1. *If $h \in H_{1/2}^1$ has Fourier expansion*

$$(2.10) \quad h(\tau) = \sum_n b(n)\mathcal{M}_n(y)e(nx) + \sum_n a(n)\mathcal{W}_n(y)e(nx),$$

then $h = 0$ if and only if $b(n) = 0$ for all $n \equiv 0, 1 \pmod{4}$.

It is now easy to explain the relation between mock modular forms and weakly harmonic ones (c.f. [50]). It follows easily from (2.7), (2.8) and (2.9), or directly, that for $g(\tau) \in M_{3/2}^1$

$$\xi_{1/2} g^*(\tau) = -2g(\tau),$$

where $g^*(\tau)$ was defined in (1.4). As a consequence we see that if $f \in \mathbb{M}_{1/2}$ and if $\hat{f} = f + g^*$ is its completion, then $\hat{f} \in H_{1/2}^1$ since $\xi_{1/2} \hat{f}(\tau) = -2g(\tau)$ so $\Delta_{1/2} \hat{f} = 0$. Also $\hat{f}(\tau)$ satisfies the plus space condition. In fact it is easy to see that $f \mapsto \hat{f}$ defines an isomorphism from $\mathbb{M}_{1/2}$ to $H_{1/2}^1$. Given $h \in H_{1/2}^1$ let $g = -\frac{1}{2} \xi_{1/2}(h) \in M_{3/2}^1$ and take $h^+ = h - g^*$. It is easily checked that

²The definition of harmonic weak Maass forms, for example as given in [7] and elsewhere, is more restrictive and does not apply to the non-holomorphic $h \in H_{1/2}^1$, so we use the terminology *weakly harmonic* to avoid confusion.

$h \mapsto h^+$ gives the inverse of $f \mapsto \hat{f}$. Call h^+ the *holomorphic part* of h . In terms of the Fourier expansion (2.6)

$$(2.11) \quad h^+(\tau) = \sum_{n \leq 0} b(n)q^n + \sum_{n > 0} a(n)n^{-1/2}q^n.$$

The next result gives one natural basis for $H_{1/2}^1$.

Proposition 1. *For each $d \equiv 0, 1 \pmod{4}$ there is a unique $h_d \in H_{1/2}^1$ with Fourier expansion of the form*

$$(2.12) \quad h_d(\tau) = \mathcal{M}_d(y)e(dx) + \sum_{n \equiv 0, 1(4)} a_d(n)\mathcal{W}_n(y)e(nx).$$

The set $\{h_d\}_{d \equiv 0, 1(4)}$ forms a basis for $H_{1/2}^1$. The coefficients $a_d(n)$ satisfy the symmetry relation

$$(2.13) \quad a_d(n) = a_n(d)$$

for all integers $n, d \equiv 0, 1 \pmod{4}$. When $d > 0$ we have

$$(2.14) \quad \xi_{1/2} h_d(\tau) = -2d^{1/2} g_d(\tau),$$

where $g_d \in M_{3/2}^1$ has Fourier expansion given in (1.10).

Theorem 2 is an immediate consequence of this proposition. We see that for $d \leq 0$ we have that $h_d = f_d$ from (1.11) and $a(n, d) = n^{-1/2}a_d(n)$ unless $n = d < 0$, in which case $a_d(d) = |d|^{1/2}$. If $d > 0$ let $f_d(\tau) = \sum_{n > 0} a(n, d)q^n$ be the holomorphic part of $d^{-1/2}h_d$. This gives the $f_d(\tau)$ from Theorem 2 and we find that for $n > 0$ we have

$$(2.15) \quad a(n, d) = (dn)^{-1/2}a_d(n).$$

We remark that the fact we quoted from [49] that $\{f_d\}_{d \leq 0}$ from (1.11) gives the Borcherds basis for $M_{1/2}^1$ also follows from the symmetry relation (2.13) and (2.14) of Proposition 1.

We now turn to the construction of h_d . We will give a uniform construction using Poincaré series. Due to some delicate convergence issues that arise from this approach, we will define them through analytic continuation. For fixed s with $\operatorname{Re}(s) > 1/2$ and $n \in \mathbb{Z}$ let

$$\mathcal{M}_n(y, s) = \begin{cases} \Gamma(2s)^{-1}(4\pi|n|y)^{-\frac{1}{4}}M_{\frac{1}{4} \operatorname{sgn} n, s - \frac{1}{2}}(4\pi|n|y) & \text{if } n \neq 0, \\ y^{s - \frac{1}{4}} & \text{if } n = 0 \end{cases}$$

$$\mathcal{W}_n(y, s) = \begin{cases} |n|^{-\frac{3}{4}}\Gamma(s + \frac{\operatorname{sgn} n}{4})^{-1}(4\pi y)^{-\frac{1}{4}}W_{\frac{1}{4} \operatorname{sgn} n, s - \frac{1}{2}}(4\pi|n|y), & \text{if } n \neq 0 \\ \frac{2^{2s - \frac{1}{2}}}{(2s - 1)\Gamma(2s - 1/2)}y^{\frac{3}{4} - s}, & \text{if } n = 0, \end{cases}$$

where M and W are the usual Whittaker functions (see Appendix A). By (A.6) and (A.7), for $n \neq 0$ we have that

$$(2.16) \quad \mathcal{M}_n(y) = \mathcal{M}_n(y, 3/4) \text{ and } \mathcal{W}_n(y) = \mathcal{W}_n(y, 3/4)$$

where $\mathcal{M}_n(y)$ and $\mathcal{W}_n(y)$ were given in (2.8) and (2.7). However, $\mathcal{M}_0(y) = \mathcal{W}_0(y, 3/4)$ and $\mathcal{W}_0(y) = \mathcal{M}_0(y, 3/4)$.³ We also need the usual I and J -Bessel functions, defined for fixed ν and $y > 0$ by (see e.g. [34])

$$(2.17) \quad I_\nu(y) = \sum_{k=0}^{\infty} \frac{(y/2)^{\nu+2k}}{k! \Gamma(\nu + k + 1)} \quad \text{and} \quad J_\nu(y) = \sum_{k=0}^{\infty} \frac{(-1)^k (y/2)^{\nu+2k}}{k! \Gamma(\nu + k + 1)}.$$

³This notational switching is inessential but gives a cleaner statement of Proposition 1 and some other results.

For $m \in \mathbb{Z}$ let

$$(2.18) \quad \psi_m(\tau, s) = \mathcal{M}_m(y, s)e(mx).$$

It follows from (A.3) and (2.4) that

$$\Delta_{1/2}\psi_m(\tau, s) = (s - \frac{1}{4})(\frac{3}{4} - s)\psi_m(\tau, s).$$

Define the Poincaré series

$$P_m(\tau, s) = \sum_{g \in \Gamma_\infty \backslash \Gamma_0(4)} j(g, \tau)^{-1} \psi_m(g\tau, s),$$

where Γ_∞ is the subgroup of translations. By (A.5) this series converges absolutely and uniformly on compacta for $\operatorname{Re} s > 1$. The function $P_0(\tau, s)$ is the usual weight $1/2$ Eisenstein series. It is clear that for $\operatorname{Re}(s) > 1$ and any m the function $P_m(\tau, s)$ has weight $1/2$ and that P_m satisfies

$$\Delta_{1/2}P_m(\tau, s) = (s - \frac{1}{4})(\frac{3}{4} - s)P_m(\tau, s).$$

As in [32], in order to get forms whose Fourier expansions are supported on $n \equiv 0, 1 \pmod{4}$ we will employ the projection operator $\operatorname{pr}^+ = \frac{2}{3}(U_4 \circ W_4) + \frac{1}{3}$, where

$$(U_4 f)(\tau) = \frac{1}{4} \sum_{\nu \pmod{4}} f\left(\frac{\tau + \nu}{4}\right) \quad \text{and} \quad (W_4 f)(\tau) = \left(\frac{2\tau}{i}\right)^{-1/2} f(-1/4\tau).$$

For each $d \equiv 0, 1 \pmod{4}$ and $\operatorname{Re}(s) > 1$ define

$$(2.19) \quad P_d^+(\tau, s) = \operatorname{pr}^+(P_d(\tau, s)).$$

Proposition 2. *For any $d \equiv 0, 1 \pmod{4}$ and $\operatorname{Re}(s) > 1$ the function $P_d^+(\tau, s)$ has weight $1/2$ and satisfies*

$$\Delta_{1/2}P_d^+(\tau, s) = (s - \frac{1}{4})(\frac{3}{4} - s)P_d^+(\tau, s).$$

Its Fourier expansion is given by

$$(2.20) \quad P_d^+(\tau, s) = \mathcal{M}_d(y, s)e(dx) + \sum_{n \equiv 0, 1(4)} b_d(n, s) \mathcal{W}_n(y, s)e(nx), \quad \text{where}$$

$$(2.21) \quad b_d(n, s) = \sum_{0 < c \equiv 0(4)} K^+(d, n; c) \times \begin{cases} 2^{\frac{1}{2}} \pi |dn|^{\frac{1}{4}} c^{-1} I_{2s-1}\left(\frac{4\pi\sqrt{|dn|}}{c}\right) & \text{if } dn < 0; \\ 2^{\frac{1}{2}} \pi |dn|^{\frac{1}{4}} c^{-1} J_{2s-1}\left(\frac{4\pi\sqrt{|dn|}}{c}\right) & \text{if } dn > 0, \\ \pi^{s+\frac{1}{4}} |d+n|^{s-\frac{1}{4}} c^{-2s} & \text{if } dn = 0, d+n \neq 0, \\ 2^{\frac{1}{2}-2s} \pi^{\frac{1}{2}} \Gamma(2s) c^{-2s} & \text{if } d = n = 0. \end{cases}$$

The sum defining each $b_d(n, s)$ is absolutely convergent.

Proof. The first statement is clear. So is the last statement using the trivial bound for $K^+(d, n; c)$ and the definitions (2.17).

For the calculation of the Fourier expansion we employ the following lemma, whose proof is standard and follows from an application of Poisson summation using an integral formula found in [15, p. 176]. See [30, Lemma 2, p. 20] or [32] for the prototype result.

Lemma 2. *Let $\begin{pmatrix} a & b \\ c & d \end{pmatrix} \in SL(2, \mathbb{R})$ have $c > 0$ and suppose that $\operatorname{Re}(s) > 1/2$. Then for ψ_m defined in (2.18) with any $m \in \mathbb{Z}$, we have*

$$\sum_{r \in \mathbb{Z}} (c(\tau + r) + d)^{-1/2} \psi_m \left(\frac{a(\tau + r) + b}{c(\tau + r) + d}, s \right) = 2\pi i^{-1/2} \sum_{n \in \mathbb{Z}} e\left(\frac{am+nd}{c}\right) \mathcal{W}_n(y, s) e(nx) \\ \times \begin{cases} c^{-1} |mn|^{\frac{1}{4}} J_{2s-1}(4\pi \sqrt{|mn|} c^{-1}) & \text{if } mn > 0 \\ c^{-1} |mn|^{\frac{1}{4}} I_{2s-1}(4\pi \sqrt{|mn|} c^{-1}) & \text{if } mn < 0 \\ 2^{-\frac{1}{2}} \pi^{s-\frac{3}{4}} c^{-2s} |m+n|^{s-\frac{1}{4}} & \text{if } mn = 0, m+n \neq 0, \\ \pi^{-\frac{1}{2}} (2c)^{-2s} \Gamma(2s) & \text{if } m = n = 0, \end{cases}$$

where both sides of the identity converge uniformly on compact subsets of \mathcal{H} .

With this Lemma, the computation of the Fourier coefficients parallels so closely that given in [30, pp. 18–27] in the holomorphic case that we will omit the details. \square

It is a well-known consequence of the theory of the resolvent kernel that $P_d(\tau, s)$ has an analytic continuation in s to $\operatorname{Re}(s) > 1/2$ except for possibly finitely many simple poles in $(1/2, 1)$. These poles may only occur at points of the discrete spectrum of $\Delta_{1/2}$ on the Hilbert space consisting of weight $1/2$ functions f on \mathcal{H} that satisfy

$$\int_{\Gamma \backslash \mathcal{H}} |f(\tau)|^2 y d\mu < \infty \quad (d\mu = y^{-2} dx dy),$$

and this space contains the residues.⁴ It is easily seen from (2.19) that $P_d^+(\tau, s)$ also has an analytic continuation to $\operatorname{Re}(s) > 1/2$ with at most finitely many simple poles in $(1/2, 1)$. Actually, such poles can only occur in $(\frac{1}{2}, \frac{3}{4}]$, since by (3.7) and (2.17) the series in (2.21) giving the Fourier coefficient $b_d(n, s)$ converges absolutely for $\operatorname{Re}(s) > 3/4$. Thus for $\operatorname{Re}(s) > 1/2$ away from these poles the function $P_d^+(\tau, s)$ has weight $1/2$ and satisfies

$$\Delta_{1/2} P_d^+(\tau, s) = (s - \frac{1}{4})(\frac{3}{4} - s) P_d^+(\tau, s).$$

Furthermore a residue at $s = 3/4$ is a weight $1/2$ weakly harmonic form $f \in H_{1/2}^1$. In fact, the Fourier expansion of f can be obtained from that of P_d^+ in (2.20) by taking residues term by term, a process that is easily justified using the integral representations for the Fourier coefficients since the convergence is uniform on compacta. This shows that the Fourier expansion of f is supported on n with $n \equiv 0, 1 \pmod{4}$ and that it can have no exponentially growing terms. Another way to see these facts is to observe that f is the projection of the residue of P_d , which comes from the discrete spectrum. Thus by Lemma 1 applied to $f - b(0)\theta$, we obtain the following result.

Lemma 3. *For each d and each $\tau \in \mathcal{H}$ the function $P_d^+(\tau, s)$ has an analytic continuation around $s = 3/4$ with at most a simple pole there with residue*

$$(2.22) \quad \operatorname{res}_{s=3/4} P_d^+(\tau, s) = \rho_d \theta(\tau),$$

where $\rho_d \in \mathbb{C}$.

When $d = 0$ this result is well-known. In fact, $b_0(n, s)$ can be computed in terms of Dirichlet L -functions. We have the following formulas (see e.g. [23]).

⁴See [15, p.179] and its references, especially [40] and [14]. A very clear treatment when the weight is 0 and the multiplier is trivial is given in [36]. In particular, see Satz 6.8 p.60; the case of weight $1/2$ is similar.

Lemma 4. For $m \in \mathbb{Z}^+$ and D a fundamental discriminant we have that

$$(2.23) \quad \sum_{n|m} \left(\frac{D}{n}\right) b_0\left(\frac{Dm^2}{n^2}, s\right) = 2^{2-4s} \pi^{s+\frac{1}{4}} m^{\frac{3}{2}-2s} |D|^{s-\frac{1}{4}} \sigma_{4s-2}(m) \frac{L_D(2s - \frac{1}{2})}{\zeta(4s-1)} \quad \text{and}$$

$$(2.24) \quad b_0(0, s) = \pi^{\frac{1}{2}} 2^{\frac{5}{2}-6s} \Gamma(2s) \frac{\zeta(4s-2)}{\zeta(4s-1)},$$

where L_D is the Dirichlet L -function.

By Möbius inversion, (2.23) gives for $m \neq 0$ the identity

$$(2.25) \quad b_0(Dm^2, s) = 2^{2-4s} \pi^{s+\frac{1}{4}} |D|^{s-\frac{1}{4}} \frac{L_D(2s-1/2)}{\zeta(4s-1)} \sum_{n|m} \mu(m/n) \left(\frac{D}{m/n}\right) n^{\frac{3}{2}-2s} \sigma_{4s-2}(n).$$

This yields a direct proof of Lemma 3 in case $d = 0$. Since $b_0(d, s) = b_d(0, s)$, which is clear from (3.4) and (2.21), a calculation using (2.25) and the (2.24) also gives the constant ρ_d in (2.22):

$$(2.26) \quad \rho_d = \begin{cases} \frac{3}{4\pi} & \text{if } d = 0, \\ \frac{6}{\pi} \sqrt{d} & \text{if } d \text{ is a non-zero square,} \\ 0 & \text{otherwise.} \end{cases}$$

We are finally ready to define the basis functions h_d . For $d \neq 0$ let

$$(2.27) \quad h_d(\tau, s) = P_d^+(\tau, s) - \frac{b_d(0, s)}{b_0(0, s)} P_0^+(\tau, s).$$

It has the Fourier expansion

$$(2.28) \quad h_d(\tau, s) = \mathcal{M}_d(y, s) e(dx) - \frac{b_d(0, s)}{b_0(0, s)} y^{s-\frac{1}{4}} + \sum_{0 \neq n \equiv 0, 1(4)} a_d(n, s) \mathcal{W}_n(y, s) e(nx), \quad \text{where}$$

$$(2.29) \quad a_d(n, s) = b_d(n, s) - \frac{b_d(0, s) b_0(n, s)}{b_0(0, s)}.$$

Lemma 5. For each nonzero $d \equiv 0, 1 \pmod{4}$ the function $h_d(\tau, s)$ defined in (2.27) has an analytic continuation to $s = 3/4$ and

$$h_d(\tau, 3/4) = h_d(\tau) \in H_{1/2}^1.$$

The Fourier expansion of each such h_d at ∞ has the form (2.12), where for each nonzero $n \equiv 0, 1 \pmod{4}$ we have

$$a_d(n) = \lim_{s \rightarrow 3/4^+} a_d(n, s).$$

Furthermore, $a_d(0) = 2\sqrt{d}$ if d is a square and $a_d(0) = 0$ otherwise.

Proof. Observe that $h_d(\tau, s)$ defined in (2.27) is holomorphic at $s = 3/4$, since otherwise by Proposition 3 it would have as residue there a nonzero multiple of $\theta(\tau)$, which cannot happen since (2.28) does not yield the constant term in θ . From (2.28) its Fourier expansion is given by

$$h_d(\tau) = \mathcal{M}_d(y) e(dx) + \sum_{n \equiv 0, 1(4)} a_d(n) \mathcal{W}_n(y) e(nx),$$

where $a_d(n) = \lim_{s \rightarrow 3/4^+} a_d(n, s)$ for $n \neq 0$ and, after recalling the definition of $\mathcal{W}_0(y)$ from (2.7), we have that

$$(2.30) \quad a_d(0) = \lim_{s \rightarrow 3/4^+} \frac{b_d(0, s)}{4b_0(0, s)}.$$

Here again we use the integral representations for the Fourier coefficients and the fact that $h_d(\tau, s) \rightarrow h_d(\tau)$ uniformly on compacta as $s \rightarrow 3/4^+$. Thus $h_d \in H_{1/2}^1$ for all $d \neq 0$. The last statement of Lemma 5 can easily be obtained from (2.30), (2.25) and (2.24). \square

Continuing with the proof of Proposition 1, we next show that the symmetry relation (2.13) holds. By (3.4) and (2.21) we have that $b_d(n, s) = b_n(d, s)$, hence by (2.29)

$$(2.31) \quad a_d(n, s) = a_n(d, s).$$

Now (2.13) follows from Lemma 5 and (2.31), where we use that $h_0 = \theta$ in case $nd = 0$. Note that $a_0(0) = 0$. A direct calculation using (2.9) together with (2.13) yields (2.14). This completes the proof of Proposition 1 and hence of Theorem 2.

3. BINARY QUADRATIC FORMS AND KLOOSTERMAN SUMS

Before turning to the proof of Theorem 3, we need to give some basic results about binary quadratic forms and Kloosterman sums. Recall that \mathcal{Q}_d is the set of integral binary quadratic forms $Q(x, y) = ax^2 + bxy + cy^2 = [a, b, c]$ of discriminant $d = b^2 - 4ac$ that are positive definite if $d < 0$. Let $\mathcal{Q}_d^+ \subset \mathcal{Q}_d$ be those forms with $a > 0$, so that $\mathcal{Q}_d = \mathcal{Q}_d^+$ when $d < 0$. Let $Q \mapsto gQ$ be the usual action of Γ that is compatible with linear fractional action on the roots of $Q(\tau, 1) = 0$. Explicitly,

$$(gQ)(x, y) = Q(\delta x - \beta y, -\gamma x + \alpha y), \quad \text{where } g = \pm \begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix} \in \Gamma.$$

As is well known, the resulting set of classes $\Gamma \backslash \mathcal{Q}_d$ is finite and those classes consisting of primitive forms make up an abelian group under composition. Let $\Gamma_Q = \{g \in \Gamma; gQ = Q\}$ be the group of automorphs of Q . If $d < 0$ then $|\Gamma_Q| = 1$ unless $Q \sim a(x^2 + y^2)$ or $Q \sim a(x^2 + xy + y^2)$, in which case $|\Gamma_Q| = 2$ or 3 , respectively. If $d > 0$ is not a square then Γ_Q is infinite cyclic with a distinguished generator denoted by g_Q , which for primitive Q is given by

$$(3.1) \quad g_Q = \pm \begin{pmatrix} \frac{t+bu}{2} & cu \\ -au & \frac{t-bu}{2} \end{pmatrix}$$

where t, u are the smallest positive integral solutions of $t^2 - du^2 = 4$. If $\delta = \gcd(a, b, c)$ then $g_Q = g_{Q/\delta}$.

Suppose that D is a fundamental discriminant, i.e. the discriminant of $\mathbb{Q}(\sqrt{D})$, and that d is a discriminant. For $Q = [a, b, c]$ with discriminant dD let

$$(3.2) \quad \chi(Q) = \chi_D(Q) = \begin{cases} \left(\frac{D}{r}\right) & \text{if } (a, b, c, D) = 1 \text{ where } Q \text{ represents } r \text{ and } (r, D) = 1, \\ 0, & \text{if } (a, b, c, D) > 1. \end{cases}$$

Here (\cdot) is the Kronecker symbol. It is well-known that χ is well-defined on classes $\Gamma \backslash \mathcal{Q}_{dD}$, that χ restricts to a real character (a genus character) on the group of primitive classes and that all such characters arise this way. We have the identity

$$(3.3) \quad \chi_D(-Q) = (\text{sgn } D)\chi_D(Q).$$

If d is also fundamental we have that $\chi_D = \chi_d$ on $\Gamma \backslash \mathcal{Q}_{dD}$. A good reference for the basic theory of these characters is [19, p. 508].

A crucial ingredient in what follows is an identity connecting the weight $1/2$ Kloosterman sum with a certain exponential sum taken over solutions to a quadratic congruence, a quadratic Weyl sum. In a special case this identity is due to Salié and variants have found many applications in the theory of modular forms. We shall use a general version due essentially to Kohnen [32]. To define the weight $1/2$ Kloosterman sum we need an explicit formula for the theta multiplier in $j(\gamma, \tau)$, which was defined in (1.2). This may be found in [41, p. 447]. As usual, for non-zero $z \in \mathbb{C}$ and $v \in \mathbb{R}$ we define $z^v = |z|^v \exp(iv \arg z)$ with $\arg z \in (-\pi, \pi]$. We have

$$j(\gamma, \tau) = (c\tau + a)^{1/2} \varepsilon_a^{-1} \left(\frac{c}{a} \right) \quad \text{for } \gamma = \pm \begin{pmatrix} * & * \\ c & a \end{pmatrix} \in \Gamma_0(4),$$

where $\left(\frac{c}{a} \right)$ is the extended Kronecker symbol and

$$\varepsilon_a = \begin{cases} 1 & \text{if } a \equiv 1 \pmod{4} \\ i & \text{if } a \equiv 3 \pmod{4}. \end{cases}$$

For $c \in \mathbb{Z}^+$ with $c \equiv 0 \pmod{4}$ and $m, n \in \mathbb{Z}$ let

$$K_{1/2}(m, n; c) = \sum_{a \pmod{c}} \left(\frac{c}{a} \right) \varepsilon_a e \left(\frac{ma + n\bar{a}}{c} \right)$$

be the weight $1/2$ Kloosterman sum. Here $\bar{a} \in \mathbb{Z}$ satisfies

$$a\bar{a} \equiv 1 \pmod{c}.$$

It is convenient to define the modified Kloosterman sum

$$K^+(m, n; c) = (1 - i)K_{1/2}(m, n; c) \times \begin{cases} 1 & \text{if } c/4 \text{ is even} \\ 2 & \text{otherwise.} \end{cases}$$

It is easily checked that

$$(3.4) \quad K^+(m, n; c) = K^+(n, m; c) = \overline{K^+(n, m; c)}.$$

The associated exponential sum is defined for $d \equiv 0, 1 \pmod{4}$ and fundamental D by

$$(3.5) \quad S_m(d, D; c) = \sum_{\substack{b \pmod{c} \\ b^2 \equiv Dd \pmod{c}}} \chi \left(\left[\frac{c}{4}, b, \frac{b^2 - Dd}{c} \right] \right) e \left(\frac{2mb}{c} \right),$$

where χ was defined in (3.2). Clearly

$$S_{-m}(d, D; c) = \overline{S_m(d, D; c)} = S_m(d, D; c).$$

The following identity is proved by a slight modification of the proof given by Kohnen in [32, Prop. 5, p. 259] (see also [9], [25] and [42]).

Proposition 3. *For positive $c \equiv 0 \pmod{4}$, $d, m \in \mathbb{Z}$ with $d \equiv 0, 1 \pmod{4}$ and D a fundamental discriminant, we have*

$$S_m(d, D; c) = \sum_{n | (m, \frac{c}{4})} \left(\frac{D}{n} \right) \sqrt{\frac{n}{c}} K^+ \left(d, \frac{m^2 D}{n^2}; \frac{c}{n} \right).$$

By Möbius inversion in two variables this can be written in the form

$$(3.6) \quad c^{-1/2} K^+(d, m^2 D, c) = \sum_{n | (m, \frac{c}{4})} \mu(n) \left(\frac{D}{n} \right) S_{m/n} \left(d, D; \frac{c}{n} \right).$$

Note that this gives an identity for $K^+(d, d', c)$ for any pair $d, d' \equiv 0, 1 \pmod{4}$. An immediate consequence of (3.6) and the obvious upper bound

$$S_m(d, D; c) \ll_{\epsilon} c^{\epsilon}$$

is the upper bound

$$(3.7) \quad K^+(d, d', c) \ll_{\epsilon} c^{1/2+\epsilon},$$

which holds for any $\epsilon > 0$. Furthermore, since for any $m, n \in \mathbb{Z}$ we have

$$K_{1/2}(m, n; c) = \frac{1}{4}K_{1/2}(4m, 4n; 4c),$$

(3.7) implies that for any $m, n \in \mathbb{Z}$

$$K_{1/2}(m, n, c) \ll_{\epsilon} c^{1/2+\epsilon}.$$

This elementary bound correspond to Weil's bound for the ordinary (weight 0) Kloosterman sum

$$K_0(m, n; c) = \sum_{\substack{a \pmod{c} \\ (a, c) = 1}} e\left(\frac{ma+n\bar{a}}{c}\right),$$

which states that (see [44], [21, Lemma 2])

$$(3.8) \quad K_0(m, n; c) \ll_{\epsilon} (m, n, c)^{1/2} c^{1/2+\epsilon}.$$

4. CYCLE INTEGRALS OF POINCARÉ SERIES

As further preparation for the proof of Theorem 3, in this section we will compute the cycle integrals of certain general Poincaré series, which we will then specialize in order to treat j_m . To begin we need to make some elementary observations about cycle integrals. For $Q \in \mathcal{Q}_d$ with $d > 0$ not a square let S_Q be the oriented semi-circle defined by

$$(4.1) \quad a|\tau|^2 + b \operatorname{Re} \tau + c = 0,$$

directed counterclockwise if $a > 0$ and clockwise if $a < 0$. Clearly

$$(4.2) \quad S_{gQ} = gS_Q,$$

for any $g \in \Gamma$. Given $z \in S_Q$ let C_Q be the directed arc on S_Q from z to $g_Q z$, where g_Q was defined in (3.1). It can easily be checked that C_Q has the same orientation as S_Q . It is convenient to define

$$(4.3) \quad d\tau_Q = \frac{\sqrt{d} d\tau}{Q(\tau, 1)}.$$

If $\tau' = g\tau$ for some $g \in \Gamma$ we have

$$(4.4) \quad d\tau'_{gQ} = d\tau_Q.$$

For any Γ -invariant function f on \mathcal{H} the integral $\int_{C_Q} f(\tau) d\tau_Q$ is both independent of $z \in S_Q$ and is a class invariant. This is an immediate consequence of the following lemma that expresses this cycle integral as a sum of integrals over arcs in a fixed fundamental domain for Γ . This lemma will also be used in the proof of Theorem 5. Let \mathcal{F} be the standard fundamental domain for Γ

$$\mathcal{F} = \left\{ \tau \in \mathcal{H}; -\frac{1}{2} \leq \operatorname{Re} \tau \leq 0, |\tau| \geq 1 \right\} \cup \left\{ \tau \in \mathcal{H}; 0 < \operatorname{Re} \tau < \frac{1}{2}, |\tau| > 1 \right\}.$$

Lemma 6. *Let $Q \in \mathcal{Q}_d$ be a form with $d > 0$ not a square and $\mathcal{F}' = g\mathcal{F}$ be the image of \mathcal{F} under any fixed $g \in \Gamma$. Suppose that f is Γ -invariant and continuous on S_Q . Then for any $z \in S_Q$ we have*

$$(4.5) \quad \int_{C_Q} f(\tau) d\tau_Q = \sum_{q \in (Q)} \int_{S_q \cap \mathcal{F}'} f(\tau) d\tau_q,$$

where (Q) denotes the class of Q .

Proof. Let $\tilde{f}(\tau) = f(\tau)$ if $\tau \in \mathcal{F}'$ and $\tilde{f}(\tau) = 0$ otherwise, so $f(\tau) = \sum_{g \in \Gamma} \tilde{f}(g\tau)$ with only a discrete set of exceptions. Thus

$$\int_{C_Q} f(\tau) d\tau_Q = \int_{C_Q} \sum_{g \in \Gamma} \tilde{f}(g\tau) d\tau_Q = \sum_{g \in \Gamma/\Gamma_Q} \sum_{\sigma \in \Gamma_Q} \int_{C_Q} \tilde{f}(g\sigma\tau) d\tau_Q = \sum_{g \in \Gamma/\Gamma_Q} \int_{S_Q} \tilde{f}(g\tau) d\tau_Q.$$

Take $g\tau$ as a new variable. By (4.2) and (4.4) we get

$$\int_{C_Q} f(\tau) d\tau_Q = \sum_{g \in \Gamma/\Gamma_Q} \int_{S_{gQ}} \tilde{f}(\tau) d\tau_{gQ},$$

which immediately yields (4.5). □

The general Poincaré series are built from a test function $\phi : \mathbb{R}^+ \rightarrow \mathbb{C}$ assumed to be smooth and to satisfy $\phi(y) = O_\varepsilon(y^{1+\varepsilon})$, for any $\varepsilon > 0$. For any $m \in \mathbb{Z}$ let

$$(4.6) \quad G_m(\tau, \phi) = \sum_{g \in \Gamma_\infty \setminus \Gamma} e(m \operatorname{Re} g\tau) \phi(\operatorname{Im} g\tau).$$

This sum converges uniformly on compacta and defines a smooth Γ -invariant function on \mathcal{H} . We will express its cycle integrals in terms of the sum $S_m(d, D; c)$ from (3.5). Define for $t > 0$ the integral transform.

$$\Phi_m(t) = \int_0^\pi \cos(2\pi m t \cos \theta) \phi(t \sin \theta) \frac{d\theta}{\sin \theta}.$$

For ϕ as above we see that this integral converges absolutely and that $\Phi_m(t) = O_\varepsilon(t^{1+\varepsilon})$. As we have seen, we may assume without loss that $d, D > 0$.

Lemma 7. *Suppose that $d, D > 0$ with dD not a square. Then for all $m \in \mathbb{Z}$*

$$\sum_{Q \in \Gamma \setminus \mathcal{Q}_{dD}} \chi(Q) \int_{C_Q} G_m(\tau, \phi) d\tau_Q = \sum_{0 < c \equiv 0(4)} S_m(d, D; c) \Phi_m\left(\frac{2\sqrt{dD}}{c}\right).$$

Proof. For each Q , interchanging the sum defining G_m and the integral yields

$$(4.7) \quad \int_{C_Q} G_m(\tau, \phi) d\tau_Q = \sum_{g \in \Gamma_\infty \setminus \Gamma} \int_{C_Q} e(m \operatorname{Re} g\tau) \phi(\operatorname{Im} g\tau) d\tau_Q.$$

Now Γ_Q , the group of automorphs of Q , acts freely on $\Gamma_\infty \backslash \Gamma$ so we have that

$$\begin{aligned} \sum_{g \in \Gamma_\infty \backslash \Gamma} \int_{C_Q} e(m \operatorname{Re} g\tau) \phi(\operatorname{Im} g\tau) d\tau_Q &= \\ \sum_{g \in \Gamma_\infty \backslash \Gamma / \Gamma_Q} \sum_{\sigma \in \Gamma_Q} \int_{C_Q} e(m \operatorname{Re} g\sigma\tau) \phi(\operatorname{Im} g\sigma\tau) d\tau_Q &= \\ \sum_{g \in \Gamma_\infty \backslash \Gamma / \Gamma_Q} \int_{S_Q} e(m \operatorname{Re} g\tau) \phi(\operatorname{Im} g\tau) d\tau_Q. \end{aligned}$$

Applying (4.4) and (4.2) in the last expression, we get from (4.7) that

$$\int_{C_Q} G_m(\tau, \phi) d\tau_Q = \sum_{g \in \Gamma_\infty \backslash \Gamma / \Gamma_Q} \int_{S_{gQ}} e(m \operatorname{Re} \tau) \phi(\operatorname{Im} \tau) d\tau_{gQ}$$

and hence that

$$\sum_{Q \in \Gamma \backslash \mathcal{Q}_{dD}} \chi(Q) \int_{C_Q} G_m(\tau, \phi) d\tau_Q = \sum_{Q \in \Gamma_\infty \backslash \mathcal{Q}_{dD}} \chi(Q) \int_{S_Q} e(m \operatorname{Re} \tau) \phi(\operatorname{Im} \tau) d\tau_Q.$$

We now need to parameterize the cycle explicitly. Let

$$(4.8) \quad \tau_Q = \frac{-b}{2a} + \frac{i\sqrt{d}}{2|a|},$$

which is easily seen to be the apex of the circle S_Q . We can parameterize S_Q by $\theta \in (0, \pi)$ via

$$\tau = \begin{cases} \operatorname{Re} \tau_Q + e^{i\theta} \operatorname{Im} \tau_Q & \text{if } a > 0 \\ \operatorname{Re} \tau_Q - e^{-i\theta} \operatorname{Im} \tau_Q & \text{if } a < 0. \end{cases}$$

With this parameterization we find that

$$Q(\tau, 1) = \frac{d}{4a} \cdot \begin{cases} e^{2i\theta} - 1 & \text{if } a > 0 \\ e^{-2i\theta} - 1 & \text{if } a < 0 \end{cases}$$

and hence that $d\tau_Q = d\theta / \sin \theta$. Since $\chi(Q) = \chi(-Q)$ we arrive at the identity

$$\sum_{Q \in \Gamma \backslash \mathcal{Q}_{dD}} \chi(Q) \int_{C_Q} G_m(\tau, \phi) d\tau_Q = 2 \sum_{Q \in \Gamma_\infty \backslash \mathcal{Q}_{dD}^+} \chi(Q) e(m \operatorname{Re} \tau_Q) \Phi_m(\operatorname{Im} \tau_Q).$$

The proof of Lemma 7 is thus reduced to the following lemma. □

Lemma 8. *Let ϕ be as above and suppose that dD is not a square. Then for all $m \in \mathbb{Z}$ we have the identity*

$$\sum_{\Gamma_\infty \backslash \mathcal{Q}_{dD}^+} \chi(Q) e(m \operatorname{Re} \tau_Q) \phi(\operatorname{Im} \tau_Q) = \frac{1}{2} \sum_{0 < c \equiv 0(4)} S_m(d, D; c) \phi\left(\frac{2\sqrt{|dD|}}{c}\right),$$

where τ_Q is defined in (4.8).

Proof. Under the growth condition on ϕ both series are absolutely convergent, and can be rearranged at will. Consider the left hand side. For $g = \pm \begin{pmatrix} 1 & k \\ 0 & 1 \end{pmatrix} \in \Gamma_\infty$ and $Q = [a, b, c] \in \mathcal{Q}_{dD}$, $gQ = [a, b - 2ka, *]$ and so the map

$$[a, b, c] \mapsto (a, b \bmod 2a)$$

is Γ_∞ -invariant. Thus

$$\sum_{\Gamma_\infty \backslash \Omega_{dD}^+} \chi(Q) e(m \operatorname{Re} \tau_Q) \phi(\operatorname{Im} \tau_Q) = \sum_{a=1}^{\infty} \phi\left(\frac{\sqrt{|dD|}}{2a}\right) \sum_{b(2a)} \chi\left([a, b, \frac{b^2-dD}{4a}]\right) e\left(-\frac{mb}{2a}\right).$$

The sum in b is restricted to those values for which $\frac{b^2-dD}{4a}$ is an integer. This happens exactly when $b^2 \equiv dD \pmod{4a}$. Thus the inner sum is

$$\sum_{\substack{b(2a) \\ b^2 \equiv dD \pmod{4a}}} \chi\left([a, b, \frac{b^2-dD}{4a}]\right) e\left(-\frac{mb}{2a}\right) = \frac{1}{2} \sum_{\substack{b(4a) \\ b^2 \equiv dD \pmod{4a}}} \chi\left([a, b, \frac{b^2-dD}{4a}]\right) e\left(-\frac{2mb}{4a}\right) = \frac{1}{2} S_m(d, D; 4a).$$

Replace $4a$ by c to finish the proof. \square

We remark that the positive definite version of Lemma 7 is following well-known formula for $dD < 0$:

$$(4.9) \quad \sum_{Q \in \Gamma \backslash \Omega_{dD}} w_Q^{-1} \chi(Q) G_m(\tau, \phi) = \frac{1}{2} \sum_{0 < c \equiv 0(4)} S_m(d, D; c) \phi\left(\frac{2\sqrt{|dD|}}{c}\right).$$

This formula is an immediate consequence of Lemma 8.

Now we will specialize the Poincaré series G_m from (4.6) and construct the modular functions j_m . Let $G_m(\tau, s) = G_m(\tau, \phi_{m,s})$, where

$$\phi_{m,s}(y) = \begin{cases} y^s & \text{if } m = 0 \\ 2\pi|m|^{\frac{1}{2}} y^{\frac{1}{2}} I_{s-\frac{1}{2}}(2\pi|m|y) & \text{if } m \neq 0, \end{cases}$$

with $I_{s-1/2}$ the Bessel function as before. The resulting Γ -invariant function satisfies

$$\Delta_0 G_m(\tau, s) = s(1-s) G_m(\tau, s).$$

The function G_0 is the usual Eisenstein series while G_m for $m \neq 0$ was studied by Neunhöffer [36] and Niebur [37], among others. The required analytic properties of $G_m(\tau, s)$ in s are most easily obtained from their Fourier expansions. For the Eisenstein series we have the well known formulas (see e.g. [24])

$$G_0(\tau, s) = y^s + c_0(0, s) y^{1-s} + \sum_{n \neq 0} c_0(n, s) K_{s-\frac{1}{2}}(2\pi|n|y) e(nx),$$

where $K_{s-\frac{1}{2}}$ is the K -Bessel function (see e.g. [34]),

$$c_0(0, s) = \frac{\xi(2s-1)}{\xi(2s)} \quad \text{and for } n \neq 0 \quad c_0(n, s) = \frac{2y^{1/2}}{\xi(2s)} |n|^{s-1/2} \sigma_{1-2s}(|n|),$$

with $\xi(s) = \pi^{-\frac{s}{2}} \Gamma(s/2) \zeta(s)$. For $m \neq 0$ the Fourier expansion of G_m can be found in [15], and is given by

$$G_m(\tau, s) = 2\pi|m|^{\frac{1}{2}} y^{\frac{1}{2}} I_{s-\frac{1}{2}}(2\pi|m|y) e(mx) + c_m(0, s) y^{1-s} \\ + 4\pi|m|^{1/2} y^{1/2} \sum_{n \neq 0} |n|^{1/2} c_m(n, s) K_{s-\frac{1}{2}}(2\pi|n|y) e(nx),$$

where

$$c_m(0, s) = \frac{4\pi|m|^{1-s} \sigma_{2s-1}(|m|)}{(2s-1)\xi(2s)}$$

and

$$c_m(n; s) = \sum_{c>0} c^{-1} K_0(m, n; c) \cdot \begin{cases} I_{2s-1}(4\pi\sqrt{|mn|}c^{-1}) & \text{if } mn < 0 \\ J_{2s-1}(4\pi\sqrt{|mn|}c^{-1}) & \text{if } mn > 0. \end{cases}$$

Define for $m \in \mathbb{Z}^+$ and $\operatorname{Re}(s) > 1$

$$(4.10) \quad j_m(\tau, s) = G_{-m}(\tau, s) - \frac{2m^{1-s}\sigma_{2s-1}(m)}{\pi^{-(s+\frac{1}{2})}\Gamma(s+\frac{1}{2})\zeta(2s-1)}G_0(\tau, s).$$

It follows from its Fourier expansion, Weil's bound (3.8) and (2.17) that $j_m(\tau, s)$ has an analytic continuation to $\operatorname{Re}(s) > 3/4$. Furthermore, since a bounded harmonic function is constant, for $m \in \mathbb{Z}^+$ we have

$$(4.11) \quad j_m(\tau, 1) = j_m(\tau),$$

where j_m was defined above (1.5) (c.f. [37]). Alternatively, we could apply the theory of the resolvent kernel in weight 0 to get the analytic continuation of $j_m(\tau, s)$ up to $\operatorname{Re} s > 1/2$.

In view of (4.11), in order to compute the traces of $j_m(\tau, s)$ it is enough to compute them for $G_m(\tau, s)$. We have the following identities, which are known when $m = 0$ (Dirichlet/Hecke) and when $dD < 0$ (see e.g. [10], [9], [7]). For the convenience of the reader we will give a uniform proof.

Proposition 4. *Let $\operatorname{Re}(s) > 1$ and $m \in \mathbb{Z}$. Suppose that d and D are not both negative and that dD is not a square. Then, when $dD < 0$ we have*

$$\sum_{Q \in \Gamma \backslash Q_{dD}} \frac{\chi(Q)}{w_Q} G_m(\tau_Q, s) = \begin{cases} \sqrt{2\pi}|m|^{\frac{1}{2}}|dD|^{\frac{1}{4}} \sum_{c \equiv 0(4)} \frac{S_m(d, D; c)}{c^{1/2}} I_{s-\frac{1}{2}}\left(\frac{4\pi\sqrt{m^2|dD|}}{c}\right) & \text{if } m \neq 0 \\ 2^{s-1}|dD|^{\frac{s}{2}} \sum_{c \equiv 0(4)} \frac{S_0(d, D; c)}{c^s} & \text{if } m = 0, \end{cases}$$

while when $dD > 0$ we have

$$\sum_{Q \in \Gamma \backslash Q_{dD}} \frac{\chi(Q)}{B(s)} \int_{C_Q} G_m(\tau, s) d\tau_Q = \begin{cases} \sqrt{2\pi}|m|^{\frac{1}{2}}|dD|^{\frac{1}{4}} \sum_{c \equiv 0(4)} \frac{S_m(d, D; c)}{c^{1/2}} J_{s-\frac{1}{2}}\left(\frac{4\pi\sqrt{m^2|dD|}}{c}\right) & \text{if } m \neq 0 \\ 2^{s-1}|dD|^{\frac{s}{2}} \sum_{c \equiv 0(4)} \frac{S_0(d, D; c)}{c^s} & \text{if } m = 0, \end{cases}$$

where $B(s) = 2^s \Gamma(\frac{s}{2})^2 / \Gamma(s)$.

Proof. By (4.9) the proof of Proposition 4 reduces to the case $dD > 0$. Applying Lemma 7 when $m = 0$ we use the well-known evaluation

$$\int_0^\pi (\sin \theta)^{s-1} d\theta = 2^{s-1} \frac{\Gamma(\frac{s}{2})^2}{\Gamma(s)}.$$

When $m \neq 0$ we need the following not-so-well-known evaluation to finish the proof. □

Lemma 9. *For $\operatorname{Re}(s) > 0$ we have*

$$\int_0^\pi \cos(t \cos \theta) I_{s-\frac{1}{2}}(t \sin \theta) \frac{d\theta}{(\sin \theta)^{1/2}} = 2^{s-1} \frac{\Gamma(\frac{s}{2})^2}{\Gamma(s)} J_{s-1/2}(t).$$

Proof. Denote the left hand side by $L_s(t)$. We use the definition of $I_{s-\frac{1}{2}}$ in (2.17) to get

$$L_s(t) = \sum_{k=0}^{\infty} \frac{(t/2)^{s+2k-1/2}}{k! \Gamma(s+k+\frac{1}{2})} \int_0^\pi \cos(t \cos \theta) (\sin \theta)^{s+2k-1} d\theta.$$

Lommel's integral representation [43, p. 47] gives for $\operatorname{Re} v > -1/2$ that

$$J_\nu(y) = \frac{(y/2)^\nu}{\Gamma(\nu+\frac{1}{2})\Gamma(\frac{1}{2})} \int_0^\pi \cos(y \cos \theta) (\sin \theta)^{2\nu} d\theta.$$

Thus for $\operatorname{Re}(s) > 0$ we have that

$$L_s(t) = \Gamma\left(\frac{1}{2}\right) \sum_{k=0}^{\infty} \frac{\Gamma\left(\frac{s}{2} + k\right)}{k! \Gamma\left(s + k + \frac{1}{2}\right)} (t/2)^{s/2+k} J_{(s-1)/2+k}(t).$$

This Neumann series can be evaluated (see [43, p.143,eq.1]) giving for $\operatorname{Re}(s) > 0$

$$L_s(t) = \frac{\Gamma\left(\frac{1}{2}\right)\Gamma\left(\frac{s}{2}\right)}{\Gamma\left(\frac{s}{2} + \frac{1}{2}\right)} J_{s-1/2}(t).$$

The result follows by the duplication formula for $\Gamma(s)$. \square

5. THE TRACES IN TERMS OF FOURIER COEFFICIENTS

In this section we complete the proofs of Theorems 3 and 4. We need to express the traces of j_m in terms of the Fourier coefficients of our basis h_d . This is first done for $j_m(\tau, s)$ with $\operatorname{Re}(s) > 1$ by applying Proposition 3 to transform the sum of exponential sums in Proposition 4 into a sum of Kloosterman sums, which is then related to the coefficients of $h_d(\tau, s)$. The method of using Kloosterman sums in this way was first applied by Zagier [46] to base change, then by Kohnen [32] to the Shimura lift and more recently to weakly holomorphic forms in [4], [9], [25] and [7].

Theorem 3 follows from Lemma 5, (4.11) and the next result by taking the limit as $s \rightarrow 1^+$ of both sides of (5.1). Also we use the relationship between $a(n, d)$ and $a_d(n)$ given in and above equation (2.15). We remark that we actually get a slightly more general result than Theorem 3 in that we may allow $D < 0$, but the general result is best left in terms of the coefficients $a_d(n)$.

Proposition 5. *Let $m \in \mathbb{Z}^+$ and $\operatorname{Re}(s) > 1$. Suppose that d and D are not both negative and that dD is not a square. Then*

$$(5.1) \quad \sum_{n|m} \left(\frac{D}{n}\right) a_d\left(\frac{m^2 D}{n^2}, \frac{s}{2} + \frac{1}{4}\right) = \begin{cases} \sum_Q \chi(Q) w_Q^{-1} j_m(\tau_Q, s) & \text{if } dD < 0, \\ B(s)^{-1} \sum_Q \chi(Q) \int_{C_Q} j_m(\tau, s) d\tau_Q & \text{if } dD > 0, \end{cases}$$

where each sum on the right hand side is over $Q \in \Gamma \backslash \mathcal{Q}_{dD}$.

Proof. It is convenient to set for any $m \in \mathbb{Z}$

$$T_m(s) = \begin{cases} \sum_Q \chi(Q) w_Q^{-1} G_m(\tau_Q, s) & \text{if } dD < 0, \\ B(s)^{-1} \sum_Q \chi(Q) \int_{C_Q} G_m(\tau, s) d\tau_Q & \text{if } dD > 0, \end{cases}$$

where each sum is over $Q \in \Gamma \backslash \mathcal{Q}_{dD}$. By Propositions 4 and 3 we have for $m \neq 0$ and $\operatorname{Re}(s) > 1$ that

$$T_m(s) = \pi |2m|^{\frac{1}{2}} |dD|^{\frac{1}{4}} \sum_{n|m} \left(\frac{D}{n}\right) n^{-\frac{1}{2}} \sum_{c \equiv 0(4)} c^{-1} K^+(d, \frac{m^2 D}{n^2}; c) \cdot \begin{cases} I_{s-\frac{1}{2}} \left(\frac{4\pi}{c} \sqrt{\frac{m^2}{n^2} |dD|} \right) & \text{if } dD < 0, \\ J_{s-\frac{1}{2}} \left(\frac{4\pi}{c} \sqrt{\frac{m^2}{n^2} |dD|} \right) & \text{if } dD > 0, \end{cases}$$

while when $m = 0$ we have

$$T_0(s) = 2^{s-1} |dD|^{\frac{s}{2}} L_D(s) \sum_{c \equiv 0(4)} c^{-s-1/2} K^+(d, 0; c).$$

Thus by (2.21) of Proposition 2 we derive that

$$(5.2) \quad T_m(s) = \begin{cases} \sum_{n|m} \left(\frac{D}{n}\right) b_d\left(\frac{m^2 D}{n^2}, \frac{s}{2} + \frac{1}{4}\right), & \text{if } m \neq 0 \\ 2^{s-1} \pi^{-\frac{s+1}{2}} |D|^{\frac{s}{2}} L_D(s) b_d\left(0, \frac{s}{2} + \frac{1}{4}\right) & \text{if } m = 0. \end{cases}$$

In view of (4.10), in order to prove Proposition 5 it is enough to show that

$$(5.3) \quad \sum_{n|m} \left(\frac{D}{n}\right) a_d\left(\frac{m^2 D}{n^2}, \frac{s}{2} + \frac{1}{4}\right) = T_m(s) - \frac{2m^{1-s} \sigma_{2s-1}(m)}{\pi^{-(s+\frac{1}{2})} \Gamma(s + \frac{1}{2}) \zeta(2s-1)} T_0(s).$$

By (2.29) and the first formula of (5.2) the left hand side of (5.3) is

$$T_m(s) - \frac{b_d\left(0, \frac{s}{2} + \frac{1}{4}\right)}{b_0\left(0, \frac{s}{2} + \frac{1}{4}\right)} \sum_{n|m} \left(\frac{D}{n}\right) b_0\left(\frac{m^2 D}{n^2}, \frac{s}{2} + \frac{1}{4}\right).$$

Hence by the second formula of (5.2) we are reduced to showing that

$$b_0\left(0, \frac{s}{2} + \frac{1}{4}\right)^{-1} \sum_{n|m} \left(\frac{D}{n}\right) b_0\left(\frac{m^2 D}{n^2}, \frac{s}{2} + \frac{1}{4}\right) = \frac{2^s \pi^{s/2} |D|^{s/2} m^{1-s} \sigma_{2s-1}(m) L_D(s)}{\Gamma(s + \frac{1}{2}) \zeta(2s-1)},$$

which follows by Lemma 4. This finishes the proof of Proposition 5, hence of Theorem 3. \square

We now give a quick proof of Theorem 4. By (2.26) we have

$$\text{Res}_{s=\frac{3}{4}} P_0^+(\tau, s) = \frac{3}{4\pi} \theta(\tau).$$

The function $\widehat{\mathbf{Z}}_+(\tau)$ can now be defined through the limit formula⁵

$$(5.4) \quad \widehat{\mathbf{Z}}_+(\tau) = \frac{1}{3} \lim_{s \rightarrow \frac{3}{4}} \left(P_0^+(\tau, s) - \frac{\frac{3}{4\pi} \theta(\tau)}{s - 3/4} \right).$$

It follows from (5.4) that $\widehat{\mathbf{Z}}_+(\tau)$ has weight $1/2$ and satisfies

$$(5.5) \quad \Delta_{1/2}(\widehat{\mathbf{Z}}_+) = -\frac{1}{8\pi} \theta.$$

Finally, using (5.2) when $m = 0$ and the fact that $G_0(\tau, s)$ has a simple pole at $s = 1$ with residue $3/\pi$, one shows that $\widehat{\mathbf{Z}}_+(\tau)$ has a Fourier expansion of the form (1.19).

The statement that $\widehat{\mathbf{Z}}_+(\tau)$ has generalized shadow $\widehat{\mathbf{Z}}_-(\tau)$ from (1.18) can now be made precise since it follows from (5.5) and the easily established identity

$$\xi_{3/2} \widehat{\mathbf{Z}}_- - \frac{1}{4\pi} \theta,$$

that

$$\xi_{1/2} \widehat{\mathbf{Z}}_+ = -2\widehat{\mathbf{Z}}_-.$$

⁵We remark that a similar limit formula was considered in [11].

6. RATIONAL PERIOD FUNCTIONS

We now prove Theorem 5. First we give a rough bound for the traces in terms of m when $d > 0$ is not a square that is sufficient to show that F_d is holomorphic in \mathcal{H} .

Proposition 6. *For $d > 0$ not a square and $m \in \mathbb{Z}^+$ we have for all $\epsilon > 0$ that*

$$\mathrm{Tr}_d(j_m) \ll_{d,\epsilon} m^{5/4+\epsilon}.$$

Proof. It follows from [22, Thm 1. p. 110] that for fixed d not a square and $x > 0$, we have for all $\epsilon > 0$ that

$$(6.1) \quad \sum_{0 < n < x} S_m(d, 1; 4n) \ll_{d,\epsilon} (mx)^\epsilon (m^{5/4} + x^{3/4}),$$

after replacing d by $4d$ if necessary. For $1 < s < 2$ we have by (3) and the well-known bound (see e.g. [34, pp. 122])

$$J_\nu(y) \ll_\nu y^{-1/2}$$

that

$$\sum_{0 < n \leq m} S_m(d, 1; 4n) \sqrt{\frac{m}{n}} J_{s-\frac{1}{2}}\left(\pi \sqrt{|d|} \frac{m}{n}\right) \ll_{d,\epsilon} m^{1+\epsilon}.$$

By (2.17) we have for $x > m$

$$\sum_{m < n < x} S_m(d, 1; 4n) \sqrt{\frac{m}{n}} J_{s-\frac{1}{2}}\left(\pi \sqrt{|d|} \frac{m}{n}\right) \ll_{d,\epsilon} m^s \left| \sum_{m < n < x} S_m(d, 1; 4n) n^{-s} \right| + m^{1+\epsilon}.$$

Summation by parts and (6.1) give

$$m^s \sum_{m < n < x} S_m(d, 1; 4n) n^{-s} \ll_{d,\epsilon} m^{5/4+\epsilon}.$$

Now Proposition 6 follows by Proposition 4 and (4.10) by taking $s \rightarrow 1^+$ in the resulting uniform inequality

$$\sum_{Q \in \Gamma \backslash \Omega_d} \int_{C_Q} j_m(\tau, s) d\tau_Q \ll_{d,\epsilon} m^{5/4+\epsilon}$$

and using (4.11). \square

It follows from Theorem 3 and Proposition 6 that the function F_d defined in (1.20) for $d > 0$ not a square can be represented by the series

$$(6.2) \quad F_d(\tau) = - \sum_{m \geq 0} \mathrm{Tr}_d(j_m) q^m,$$

which gives a holomorphic function on \mathcal{H} . The basis $\{j_m\}_{m \geq 0}$ has a generating function that goes back to Faber (see e.g. [1]):

$$(6.3) \quad \sum_{m \geq 0} j_m(z) q^m = \frac{j'(\tau)}{j(z) - j(\tau)}, \quad \text{where } j'(\tau) = \frac{1}{2\pi i} \frac{dj}{d\tau}.$$

Note that this formal series converges when $\mathrm{Im}(\tau) > \mathrm{Im}(z)$ and that for fixed τ not a zero of j' it has a simple pole at $z = \tau$ with residue $(2\pi i)^{-1}$. It follows from (6.3) and (6.2) that for $\mathrm{Im}(\tau)$ sufficiently large we have

$$(6.4) \quad F_d(\tau) = \frac{1}{2\pi} \sum_{Q \in \Gamma \backslash \Omega_d} \int_{C_Q} \frac{j'(\tau)}{j(\tau) - j(z)} \frac{d\tau}{Q(\tau, 1)},$$

where we take for C_Q an arc on S_Q , the semi-circle defined in (4.1). Let $\mathcal{F}' = -\mathcal{F}^{-1}$ be the image of the standard fundamental domain under inversion $\tau \mapsto -1/\tau$. By (6.4) and Lemma 6 applied to each class of \mathcal{Q}_d and to each fundamental domain \mathcal{F} and \mathcal{F}' , we can write

$$F_d(\tau) = \frac{1}{4\pi} \sum_{Q \in \mathcal{Q}_d} \left(\int_{S_Q \cap \mathcal{F}} \frac{j'(\tau)}{j(\tau) - j(z)} \frac{d\tau}{Q(\tau, 1)} + \int_{S_Q \cap \mathcal{F}'} \frac{j'(\tau)}{j(\tau) - j(z)} \frac{d\tau}{Q(\tau, 1)} \right).$$

Now it is easily seen that each of these integrals is invariant under $Q \mapsto -Q$, so we may restrict the sum to \mathcal{Q}_d^+ , giving

$$(6.5) \quad F_d(\tau) = \frac{1}{2\pi} \sum_{Q \in \mathcal{Q}_d^+} \left(\int_{S_Q \cap \mathcal{F}} \frac{j'(\tau)}{j(\tau) - j(z)} \frac{d\tau}{Q(\tau, 1)} + \int_{S_Q \cap \mathcal{F}'} \frac{j'(\tau)}{j(\tau) - j(z)} \frac{d\tau}{Q(\tau, 1)} \right).$$

Recall from [8] that an indefinite quadratic form $Q = [a, b, c]$ is called *simple* if $c < 0 < a$. It is easily seen that $Q \in \mathcal{Q}_d$ is simple if and only if $Q \in \mathcal{Q}_d^+$ and S_Q intersects $\mathcal{F}'' = \mathcal{F} \cup \mathcal{F}'$. For simple Q let $A_Q = S_Q \cap \mathcal{F}''$ be the arc in \mathcal{F}'' oriented from right to left. Clearly A_Q must connect the two “vertical” sides of \mathcal{F}'' .⁶ Thus from (6.5) we obtain the identity

$$F_d(\tau) = \frac{1}{2\pi} \sum_{\substack{Q \text{ simple} \\ b^2 - 4ac = d}} \int_{A_Q} \frac{j'(\tau)}{j(\tau) - j(z)} \frac{dz}{Q(z, 1)}.$$

Now we deform each arc A_Q in the sum of integrals to B_Q , which is within \mathcal{F}'' and has the same endpoints as A_Q , but travels above τ . By evaluating each resulting residue at τ , we get the formula

$$F_d(\tau) = \frac{1}{2\pi} \sum_{\substack{Q \text{ simple} \\ b^2 - 4ac = d}} \int_{B_Q} \frac{j'(\tau)}{j(\tau) - j(z)} \frac{dz}{Q(z, 1)} + \frac{1}{2\pi} \sum_{\substack{Q \text{ simple} \\ b^2 - 4ac = d}} Q(\tau, 1)^{-1},$$

which is also valid at $-1/\tau$. A simple calculation now shows that (1.21) holds in a neighborhood of τ , hence for all $\tau \in \mathcal{H}$. Thus Theorem 5 follows.

Finally, for fixed $m \in \mathbb{Z}^+$ the inequality (6.1) can be used to show that the series in Proposition 4 converges when $s = 1$. They yield the formula (1.24) upon using the elementary evaluation

$$J_{1/2}(y) = \sqrt{\frac{2}{\pi y}} \sin y.$$

APPENDIX A. WHITTAKER FUNCTIONS

A standard reference for the theory of Whittaker functions is [45, Chap. 16]. Another good reference is [35]. For the convenience of the reader we will record here some of the properties of these special functions that we need.

For fixed μ, ν with $\operatorname{Re}(\nu \pm \mu + 1/2) > 0$ the Whittaker functions may be defined for $y > 0$ by [35, pp. 311, 313]

$$(A.1) \quad M_{\mu, \nu}(y) = y^{\nu + \frac{1}{2}} e^{\frac{y}{2}} \frac{\Gamma(1+2\nu)}{\Gamma(\nu + \mu + \frac{1}{2}) \Gamma(\nu - \mu + \frac{1}{2})} \int_0^1 t^{\nu + \mu - \frac{1}{2}} (1-t)^{\nu - \mu - \frac{1}{2}} e^{-yt} dt \quad \text{and}$$

$$(A.2) \quad W_{\mu, \nu}(y) = y^{\nu + \frac{1}{2}} e^{\frac{y}{2}} \frac{1}{\Gamma(\nu - \mu + \frac{1}{2})} \int_1^\infty t^{\nu + \mu - \frac{1}{2}} (t-1)^{\nu - \mu - \frac{1}{2}} e^{-yt} dt.$$

⁶For example, when $d = 12$ the simple forms are $[1, 0, -3], [1, -2, -2], [1, 2, -2], [3, 0, -1], [2, 2, -1], [2, -2, -1]$. A diagram showing the corresponding arcs A_Q in this case is given in Figure 1.

Both $M_{\mu,\nu}(y)$ and $W_{\mu,\nu}(y)$ satisfy the second order linear differential equation

$$(A.3) \quad \frac{d^2 w}{dy^2} + \left(-\frac{1}{4} + \mu y^{-1} + \left(\frac{1}{4} - \nu^2\right) y^{-2} \right) w = 0.$$

Their asymptotic behavior as $y \rightarrow \infty$ for fixed μ, ν is easily found from (A.1) and (A.2) by changing variable $t \mapsto t/y$:

$$(A.4) \quad M_{\mu,\nu}(y) \sim \frac{\Gamma(1+2\nu)}{\Gamma(\nu-\mu+\frac{1}{2})} y^{-\mu} e^{y/2} \quad \text{and} \quad W_{\mu,\nu}(y) \sim y^\mu e^{-y/2}.$$

In particular, they are linearly independent. For small y we get directly from (A.1) that

$$(A.5) \quad M_{\mu,\nu}(y) = y^{\nu+\frac{1}{2}} \left(1 + O_{\mu,\nu}(y) \right).$$

It is also apparent from (A.1) and (A.2) that when $\nu - \mu = 1/2$ we have

$$(A.6) \quad M_{\mu,\nu}(y) + (2\mu + 1)W_{\mu,\nu}(y) = \Gamma(2\mu + 2)y^{-\mu} e^{y/2},$$

while when $\nu + \mu = 1/2$ we have from (A.2) that

$$(A.7) \quad W_{\mu,\nu}(y) = y^\mu e^{-y/2}.$$

The I -Bessel and K -Bessel functions are special Whittaker functions [35]:

$$I_\nu(y) = 2^{-2\nu-\frac{1}{2}} \Gamma(\nu + 1)^{-1} y^{-\frac{1}{2}} M_{0,\nu}(2y) \quad \text{and} \quad K_\nu(y) = \sqrt{\frac{\pi}{2y}} W_{0,\nu}(2y).$$

Their asymptotic properties for large y thus follow from (A.4).

REFERENCES

- [1] Asai, T.; Kaneko, M.; Ninomiya, H., Zeros of certain modular functions and an application. *Comment. Math. Univ. St. Paul.* 46 (1997), no. 1, 93–101.
- [2] Borcherds, R. E., Automorphic forms on $O_{s+2,2}(R)$ and infinite products. *Invent. Math.* 120 (1995), no. 1, 161–213.
- [3] Bringmann, K.; Ono, K., Arithmetic properties of coefficients of half-integral weight Maass-Poincaré series. *Math. Ann.* 337 (2007), no. 3, 591–612.
- [4] Bruinier, J.H., Borcherds products and Chern classes of Hirzebruch-Zagier divisors. *Invent. Math.* 138 (1999), no. 1, 51–83.
- [5] Bruinier, J. H.; Funke, J., Traces of CM values of modular functions. *J. Reine Angew. Math.* 594 (2006), 1–33.
- [6] Bruinier, J. H.; Funke, J., On two geometric theta lifts. *Duke Math. J.* 125 (2004), no. 1, 45–90.
- [7] Bruinier and Ono, K., Heegner divisors, L-functions and harmonic weak Maass forms, to appear.
- [8] Choie, Yj.; Zagier, D., Rational period functions for $PSL(2, \mathbb{Z})$. A tribute to Emil Grosswald: number theory and related analysis, 89–108, *Contemp. Math.*, 143, Amer. Math. Soc., Providence, RI, 1993.
- [9] Duke, W., Modular functions and the uniform distribution of CM points. *Math. Ann.* 334 (2006), no. 2, 241–252.
- [10] Duke, W.; Friedlander, J. B.; Iwaniec, H. Equidistribution of roots of a quadratic congruence to prime moduli. *Ann. of Math. (2)* 141 (1995), no. 2, 423–441.
- [11] Duke, W.; Imamoglu, ., A converse theorem and the Saito-Kurokawa lift. *Internat. Math. Res. Notices* 1996, no. 7, 347–355.
- [12] Duke, W.; Jenkins, P., Integral traces of singular values of Maass forms, Preprint 2007
- [13] Eichler, M., Grenzkreisgruppen und kettenbruchartige Algorithmen. (German) *Acta Arith.* 11 1965 169–180.
- [14] Elstrodt, J; Die Resolvente zum Eigenwertproblem der automorphen Formen in der hyperbolischen Ebene. I *Math. Ann.* 203 (1973), 295–300, II *Math. Z.* 132 (1973), 99–134, III *Math. Ann.* 208 (1974), 99–132.
- [15] Fay, J. D., Fourier coefficients of the resolvent for a Fuchsian group, *J. Reine Angew. Math.* **293/294** (1977), 143–203.
- [16] Funke, J., Heegner divisors and nonholomorphic modular forms. *Compositio Math.* 133 (2002), no. 3, 289–321.

- [17] Goldfeld, D.; Hoffstein, J., Eisenstein series of $1/2$ -integral weight and the mean value of real Dirichlet L -series. *Invent. Math.* 80 (1985), no. 2, 185–208.
- [18] Gradshteyn, I.S.; Ryzhik, I.M., *Table of Integrals, Series and Products*, 5th Edition, Academic Press
- [19] Gross, B.; Kohlen, W.; Zagier, D., Heegner points and derivatives of L -series. II. *Math. Ann.* 278 (1987), no. 1-4, 497–562.
- [20] Hirzebruch, F.; Zagier, D., Intersection numbers of curves on Hilbert modular surfaces and modular forms of Nebentypus. *Invent. Math.* 36 (1976), 57–113.
- [21] Hooley, C., An asymptotic formula in the theory of numbers. *Proc. London Math. Soc.* (3) 7 (1957), 396–413.
- [22] Hooley, C., On the number of divisors of a quadratic polynomial. *Acta Math.* 110 1963 97–114.
- [23] Ibukiyama, T.; Saito, H., On zeta functions associated to symmetric matrices. II. MPI Preprint.
- [24] Iwaniec, H.; Kowalski, E., *Analytic number theory*. American Mathematical Society Colloquium Publications, 53. American Mathematical Society, Providence, RI, 2004.
- [25] Jenkins, P., Kloosterman sums and traces of singular moduli. *J. Number Theory* 117 (2006), no. 2, 301–314.
- [26] Observations on the "values" of the elliptic modular function $j(\tau)$ at real quadratics, <http://arxiv.org/abs/0905.3013>
- [27] Katok, S.; Sarnak, P., Heegner points, cycles and Maass forms. *Israel J. Math.* 84 (1993), no. 1-2, 193–227.
- [28] Knopp, M. I., Some new results on the Eichler cohomology of automorphic forms. *Bull. Amer. Math. Soc.* 80 (1974), 607–632.
- [29] Knopp, M. I., Rational period functions of the modular group. With an appendix by Georges Grinstein. *Duke Math. J.* 45 (1978), no. 1, 47–62.
- [30] Kohlen, Beziehungen zwischen Modulformen halbganzen Gewichts und Modulformen ganzen Gewichts, *Bonner mathematische Schrifte*, 131, Bonn, 1981.
- [31] Kohlen, W., Modular forms of half-integral weight on $\Gamma_0(4)$. *Math. Ann.* 248 (1980), no. 3, 249–266.
- [32] Kohlen, W., Fourier coefficients of modular forms of half-integral weight. *Math. Ann.* 271 (1985), no. 2, 237–268.
- [33] Kohlen, W.; Zagier, D., Values of L -series of modular forms at the center of the critical strip. *Invent. Math.* 64 (1981), no. 2, 175–198.
- [34] Lebedev, N. N. , *Special functions and their applications*. Revised edition, translated from the Russian and edited by Richard A. Silverman. Unabridged and corrected republication. Dover Publications, Inc., New York, 1972.
- [35] Magnus, W.; Oberhettinger, F.; Soni, R. P., *Formulas and theorems for the special functions of mathematical physics*. Third enlarged edition. *Die Grundlehren der mathematischen Wissenschaften, Band 52* Springer-Verlag New York, Inc., New York 1966
- [36] Neunhöffer, H., Über die analytische Fortsetzung von Poincarreihen. (German) *S.-B. Heidelberger Akad. Wiss. Math.-Natur. Kl.* 1973, 33–90.
- [37] Niebur, D., A class of nonanalytic automorphic functions. *Nagoya Math. J.* 52 (1973), 133–145.
- [38] Ono, K. , *Unearthing the visions of a master: harmonic Maass forms and number theory*, Harvard-MIT Current Developments in Mathematics 2008, International Press, (preliminary version).
- [39] Parson, L. A., Modular integrals and indefinite binary quadratic forms. A tribute to Emil Grosswald: number theory and related analysis, 513–523, *Contemp. Math.*, 143, Amer. Math. Soc., Providence, RI, 1993.
- [40] Roelcke, W., Das Eigenwertproblem der automorphen Formen in der hyperbolischen Ebene. I, II. (German) *Math. Ann.* 167 (1966), 292–337; *ibid.* 168 1966 261–324.
- [41] Shimura, G., On modular forms of half integral weight. *Ann. of Math.* (2) 97 (1973), 440–481.
- [42] Tóth, Á., On the evaluation of Sali sums. *Proc. Amer. Math. Soc.* 133 (2005), no. 3, 643–645.
- [43] Watson, G. N., *A Treatise on the Theory of Bessel Functions*. Cambridge University Press, Cambridge, England; The Macmillan Company, New York, 1944
- [44] Weil, A., On some exponential sums. *Proc. Nat. Acad. Sci. U. S. A.* 34, (1948). 204–207.
- [45] Whittaker, E. T.; Watson, G. N., *A course of modern analysis. An introduction to the general theory of infinite processes and of analytic functions: with an account of the principal transcendental functions*. Fourth edition. Reprinted Cambridge University Press, New York 1962
- [46] Zagier, D., Modular forms associated to real quadratic fields. *Invent. Math.* 30 (1975), no. 1, 1–46.
- [47] Zagier, D., Nombres de classes et formes modulaires de poids $3/2$. (French) *C. R. Acad. Sci. Paris Sr. A-B* 281 (1975), no. 21, A883–A886.

- [48] Zagier, D., Eisenstein series and the Riemann zeta function. Automorphic forms, representation theory and arithmetic (Bombay, 1979), pp. 275–301, Tata Inst. Fund. Res. Studies in Math., 10, Tata Inst. Fundamental Res., Bombay, 1981
- [49] Zagier, D., Traces of singular moduli. Motives, polylogarithms and Hodge theory, Part I (Irvine, CA, 1998), 211–244, Int. Press Lect. Ser., 3, I, Int. Press, Somerville, MA, 2002.
- [50] Zagier, D. Ramanujan’s mock theta functions and their applications [d’après Zwegers and Bringmann-Ono], Seminaire Bourbaki, 986
- [51] Zwegers, S. P., Mock Theta Functions, Utrecht PhD thesis, (2002) ISBN 90-393-3155-3
- [52] Zwegers, S. P. Mock θ -functions and real analytic modular forms. q -series with applications to combinatorics, number theory, and physics (Urbana, IL, 2000), 269–277, Contemp. Math., 291, Amer. Math. Soc., Providence, RI, 2001.

UCLA MATHEMATICS DEPARTMENT, BOX 951555, LOS ANGELES, CA 90095-1555
E-mail address: `wdduke@ucla.edu`

ETH, MATHEMATICS DEPT., CH-8092, ZÜRICH, SWITZERLAND
E-mail address: `ozlem@math.ethz.ch`

EOTVOS LORAND UNIVERSITY, DEPARTMENT OF ANALYSIS, BUDAPEST, HUNGARY
E-mail address: `toth@cs.elte.hu`